

# Package ‘ALUES’

May 6, 2026

**Type** Package

**Title** Agricultural Land Use Evaluation System

**Version** 0.2.1

**Date** 2022-01-10

**Maintainer** Al-Ahmadgaid B. Asaad <alahmadgaid@gmail.com>

**Author** Al-Ahmadgaid B. Asaad [aut, cre] (ORCID:  
<<https://orcid.org/0000-0003-3784-8593>>),  
Arnold R. Salvacion [aut] (ORCID:  
<<https://orcid.org/0000-0001-8868-2226>>),  
Bui Tan Yen [aut]

**Description** Evaluates land suitability for different crops production. The package is based on the Food and Agriculture Organization (FAO) and the International Rice Research Institute (IRRI) methodology for land evaluation. Development of ALUES is inspired by similar tool for land evaluation, Land Use Suitability Evaluation Tool (LUSSET). The package uses fuzzy logic approach to evaluate land suitability of a particular area based on inputs such as rainfall, temperature, topography, and soil properties. The membership functions used for fuzzy modeling are the following: Triangular, Trapezoidal and Gaussian. The methods for computing the overall suitability of a particular area are also included, and these are the Minimum, Maximum and Average. Finally, ALUES is a highly optimized library with core algorithms written in C++.

**License** MIT + file LICENSE

**LazyData** true

**LinkingTo** Rcpp

**Encoding** UTF-8

**Depends** R (>= 3.5.0), Rcpp (>= 0.10.6)

**Suggests** testthat, markdown, knitr, microbenchmark, ggmap, raster, reshape2

**RoxygenNote** 7.1.1

**VignetteBuilder** knitr

**URL** <https://github.com/alstat/ALUES/>

**BugReports** <https://github.com/alstat/ALUES/issues/>

**NeedsCompilation** yes

**Repository** CRAN

**Date/Publication** 2022-01-12 21:12:44 UTC

## Contents

ALUES-package . . . . .	7
ALFALFASoil . . . . .	7
ALFALFATemp . . . . .	8
ALFALFATerrain . . . . .	9
ALFALFAWater . . . . .	9
AVOCADOSoil . . . . .	10
AVOCADOTemp . . . . .	11
AVOCADOTerrain . . . . .	11
AVOCADOWater . . . . .	12
BAMBOOSoil . . . . .	12
BAMBOOTemp . . . . .	13
BAMBOOTerrain . . . . .	13
BAMBOOWater . . . . .	14
BANANASoil . . . . .	14
BANANATemp . . . . .	15
BANANATerrain . . . . .	16
BANANAWater . . . . .	16
BARLEYSoil . . . . .	17
BARLEYTemp . . . . .	18
BARLEYTerrain . . . . .	18
BARLEYWater . . . . .	19
BEANCAsoil . . . . .	19
BEANCATemp . . . . .	20
BEANCATerrain . . . . .	21
BEANCAWater . . . . .	21
BEANSSoil . . . . .	22
BEANSTemp . . . . .	23
BEANSTerrain . . . . .	23
BEANSWater . . . . .	24
CABBAGESoil . . . . .	24
CABBAGETemp . . . . .	25
CABBAGETerrain . . . . .	26
CABBAGEWater . . . . .	26
CARROTSSoil . . . . .	27
CARROTSTemp . . . . .	28
CARROTSTerrain . . . . .	28
CARROTSWater . . . . .	29
CASHEWSoil . . . . .	29
CASHEWTemp . . . . .	30
CASHEWTerrain . . . . .	31

CASHEWWater . . . . .	31
CASSAVASoil . . . . .	32
CASSAVATemp . . . . .	33
CASSAVATerrain . . . . .	33
CASSAWater . . . . .	34
CHICKPEASoil . . . . .	34
CHICKPEATemp . . . . .	35
CHICKPEATerrain . . . . .	36
CHICKPEAWater . . . . .	36
CINNAMONSoil . . . . .	37
CINNAMONTemp . . . . .	38
CINNAMONTerrain . . . . .	38
CINNAMONWater . . . . .	39
CITRUSSoil . . . . .	39
CITRUSTemp . . . . .	40
CITRUSTerrain . . . . .	41
CITRUSWater . . . . .	41
COCOASoil . . . . .	42
COCOATemp . . . . .	43
COCOATerrain . . . . .	43
COCOAWater . . . . .	44
COCONUTSoil . . . . .	44
COCONUTTemp . . . . .	45
COCONUTTerrain . . . . .	46
COCONUTWater . . . . .	46
COFFEEARSoil . . . . .	47
COFFEEARTemp . . . . .	48
COFFEEARTerrain . . . . .	48
COFFEEARWater . . . . .	49
COFFEEROSoil . . . . .	49
COFFEEROTemp . . . . .	50
COFFEEROTerrain . . . . .	51
COFFEEROWater . . . . .	51
COTTONSoil . . . . .	52
COTTONTemp . . . . .	53
COTTONTerrain . . . . .	53
COTTONWater . . . . .	54
COWPEASoil . . . . .	55
COWPEATemp . . . . .	55
COWPEATerrain . . . . .	56
COWPEAWater . . . . .	57
CUCUMBERSoil . . . . .	57
CUCUMBERTemp . . . . .	58
CUCUMBERTerrain . . . . .	59
CUCUMBERWater . . . . .	59
GROUNDNUTSSoil . . . . .	60
GROUNDNUTSTemp . . . . .	61
GROUNDNUTSTerrain . . . . .	61

GROUNDNUTSWater . . . . .	62
GUAVASoil . . . . .	62
GUAVATemp . . . . .	63
GUAVATerrain . . . . .	64
GUAVAWater . . . . .	64
LaoCaiLT . . . . .	65
LaoCaiTemp . . . . .	65
LaoCaiWater . . . . .	66
LONGANSoil . . . . .	67
LONGANTemp . . . . .	68
LONGANTerrain . . . . .	68
LONGANWater . . . . .	69
MAIZESoil . . . . .	69
MAIZETemp . . . . .	70
MAIZETerrain . . . . .	71
MAIZEWater . . . . .	71
MANGOSoil . . . . .	72
MANGOTemp . . . . .	73
MANGOTerrain . . . . .	73
MANGOWater . . . . .	74
MarinduqueLT . . . . .	74
MarinduqueTemp . . . . .	75
MarinduqueWater . . . . .	76
MILLETSSoil . . . . .	76
MILLETSTemp . . . . .	77
MILLETSTerrain . . . . .	78
MILLETSWater . . . . .	78
OILPALMSoil . . . . .	79
OILPALMTemp . . . . .	80
OILPALMTerrain . . . . .	80
OILPALMWater . . . . .	81
OLIVESSoil . . . . .	81
OLIVESTemp . . . . .	82
OLIVESTerrain . . . . .	83
OLIVESWater . . . . .	83
ONIONSoil . . . . .	84
ONIONTemp . . . . .	85
ONIONTerrain . . . . .	85
ONIONWater . . . . .	86
overall_suit . . . . .	86
PAPAYASoil . . . . .	88
PAPAYATemp . . . . .	88
PAPAYATerrain . . . . .	89
PAPAYAWater . . . . .	89
PEACHSoil . . . . .	90
PEACHTemp . . . . .	91
PEACHTerrain . . . . .	91
PEACHWater . . . . .	92

PEARSoil . . . . .	92
PEARTemp . . . . .	93
PEARTerrain . . . . .	93
PEARWater . . . . .	94
PEASoil . . . . .	95
PEATemp . . . . .	95
PEATerrain . . . . .	96
PEAWater . . . . .	97
PEPPERGRSoil . . . . .	97
PEPPERGRTemp . . . . .	98
PEPPERGRTerrain . . . . .	98
PEPPERGRWater . . . . .	99
PERSIMMONSoil . . . . .	100
PERSIMMONTemp . . . . .	101
PERSIMMONTerrain . . . . .	101
PERSIMMONWater . . . . .	102
PINEAPPLESoil . . . . .	102
PINEAPPLETemp . . . . .	103
PINEAPPLETerrain . . . . .	103
PINEAPPLEWater . . . . .	104
PLUMSoil . . . . .	105
PLUMTemp . . . . .	106
PLUMTerrain . . . . .	106
PLUMWater . . . . .	107
POTATOSoil . . . . .	107
POTATOSWSoil . . . . .	108
POTATOSWTemp . . . . .	109
POTATOSWTerrain . . . . .	109
POTATOSWWater . . . . .	110
POTATOTemp . . . . .	110
POTATOTerrain . . . . .	111
POTATOWater . . . . .	112
RICEBRSoil . . . . .	112
RICEBRTemp . . . . .	113
RICEBRTerrain . . . . .	114
RICEBRWater . . . . .	114
RICEIWSoil . . . . .	115
RICEIWTemp . . . . .	116
RICEIWTerrain . . . . .	116
RICEIWWater . . . . .	117
RICENFSoil . . . . .	118
RICENFTemp . . . . .	119
RICENFTerrain . . . . .	119
RICENFWater . . . . .	120
RICEURSoil . . . . .	121
RICEURTemp . . . . .	121
RICEURTerrain . . . . .	122
RICEURWater . . . . .	123

RUBBERSoil . . . . .	123
RUBBERTemp . . . . .	124
RUBBERTerrain . . . . .	125
RUBBERWater . . . . .	125
SAFFLOWERSoil . . . . .	126
SAFFLOWERTemp . . . . .	126
SAFFLOWERTerrain . . . . .	127
SAFFLOWERWater . . . . .	127
SESAMESoil . . . . .	128
SESAMETemp . . . . .	129
SESAMETerrain . . . . .	129
SESAMEWater . . . . .	130
SORGHUMSoil . . . . .	130
SORGHUMTemp . . . . .	131
SORGHUMTerrain . . . . .	132
SORGHUMWater . . . . .	132
SOYASoil . . . . .	133
SOYATemp . . . . .	134
SOYATerrain . . . . .	134
SOYAWater . . . . .	135
SUGARCANESoil . . . . .	136
SUGARCANETemp . . . . .	137
SUGARCANETerrain . . . . .	137
SUGARCANEWater . . . . .	138
suit . . . . .	138
suitability . . . . .	141
SUNFLOWERSoil . . . . .	142
SUNFLOWERTemp . . . . .	143
SUNFLOWERTerrain . . . . .	144
SUNFLOWERWater . . . . .	144
TEASoil . . . . .	145
TEATemp . . . . .	146
TEATerrain . . . . .	146
TEAWater . . . . .	147
TOBACCSOIL . . . . .	148
TOBACCOTemp . . . . .	149
TOBACCOTerrain . . . . .	149
TOBACCOWater . . . . .	150
TOMATOSoil . . . . .	150
TOMATOTemp . . . . .	151
TOMATOTerrain . . . . .	152
TOMATOWater . . . . .	152
WATERMELONSoil . . . . .	153
WATERMELONTemp . . . . .	154
WATERMELONTerrain . . . . .	154
WATERMELONWater . . . . .	155
WHEATSoil . . . . .	155
WHEATTemp . . . . .	156

WHEATTerrain . . . . .	157
WHEATWater . . . . .	157

<b>Index</b>	<b>159</b>
--------------	------------

---

ALUES-package	<i>Agricultural Land Use Evaluation System</i>
---------------	--

---

### Description

Agricultural Land Use Evaluation System (ALUES) is a package that evaluates land suitability for different crop production. The package is based on the Food and Agriculture Organization (FAO) and the International Rice Research Institute (IRRI) methodology for land evaluation. Development of ALUES is inspired by similar tool for land evaluation, Land Use Suitability Evaluation Tool (LUSSET). The package uses fuzzy logic approach to evaluate land suitability of a particular area based on inputs such as rainfall, temperature, topography, and soil properties. The membership functions used for fuzzy modeling are the following: Triangular, Trapezoidal, Gaussian, Sigmoidal and custom models with functions that can be defined by the user. The package also aims on complicated methods like considering more than one fuzzy membership function on different suitability class. The methods for computing the overall suitability of a particular area are also included, and these are the Minimum, Maximum, Product, Sum, Average, Exponential and Gamma. Finally, ALUES utilizes the power of Rcpp library for efficient computation.

### Author(s)

Al-Ahmadgaid B. Asaad <alahmadgaid@gmail.com> (maintainer)

Arnold R. Salvacion <arsalvacion@gmail.com>

Bui Tan Yen

---

ALFALFASoil	<i>Alfalfa soil requirement for land evaluation</i>
-------------	---

---

### Description

A dataset containing the soil characteristics of the crop requirements for farming Alfalfa.

### Format

A data frame with 12 rows and 8 columns

**Details**

The following are the factors for evaluation:

- SoilTe - 12 classes of soil texture (Soil Taxonomy)
- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

ALFALFATemp

*Alfalfa temp requirement for land evaluation*

---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Alfalfa.

**Format**

A data frame with 1 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

ALFALFATerrain      *Alfalfa terrain requirement for land evaluation*

---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Alfalfa.

**Format**

A data frame with 6 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization. )
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

ALFALFAWater      *Alfalfa water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Alfalfa.

**Format**

A data frame with 3 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CropLen - Length of growing period (days)
- WgAv - Precipitation of growing cycle (mm)
- WghAv - Relative humidity growing cycle (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

AVOCADOSoil

*Avocado soil requirement for land evaluation*

---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Avocado.

**Format**

A data frame with 10 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 AVOCADOTemp

*Avocado temp requirement for land evaluation*


---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Avocado.

**Format**

A data frame with 2 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TmMinXm - Avarage minimum temperature of coldest month ( C )

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 AVOCADOTerrain

*Avocado terrain requirement for land evaluation*


---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Avocado.

**Format**

A data frame with 5 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of managemnet with full mechanization. )
- Slope3 - Slope (%) (3. Low level of managemnet animal traction or handwork.)
- Flood - Flooding
- Drainage - Drainage

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

AVOCADOWater

*Avocado water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Avocado.

**Format**

A data frame with 2 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)
- WmDryLen - Length dry season (months :  $P < 1/2$  PET)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

BAMBOOSoil

*Bamboo soil requirement for land evaluation*

---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Bamboo.

**Format**

A data frame with 4 rows and 8 columns

**Details**

The following are the factors for evaluation:

- SoilDpt - Soil depth (cm)
- OC - Organic carbon (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)
- pHH2O - pH H2O

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

BAMBOOTemp

*Bamboo temp requirement for land evaluation*

---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Bamboo.

**Format**

A data frame with 1 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TyAv - Mean annual temperature (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

BAMBOOTerrain

*Bamboo terrain requirement for land evaluation*

---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Bamboo.

**Format**

A data frame with 1 rows and 8 columns

**Details**

The following are the factors for evaluation:

- SlopeD - Slope (degree, 6 classes)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

BAMBOOWater

*Bamboo water requirement for land evaluation*

---

### **Description**

A dataset containing the water characteristics of the crop requirements for farming Bamboo.

### **Format**

A data frame with 1 rows and 8 columns

### **Details**

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)

### **See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

BANANASoil

*Banana soil requirement for land evaluation*

---

### **Description**

A dataset containing the soil characteristics of the crop requirements for farming Banana.

### **Format**

A data frame with 12 rows and 8 columns

### **Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)

- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

BANANATemp

*Banana temp requirement for land evaluation*

---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Banana.

**Format**

A data frame with 3 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TyMaxAv - Mean annual maximum temperature (°C)
- TmMinXm - Avarage minimum temperature of coldest month ( C )
- TmMinXmAb - Absolute min temp. coldest month (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

BANANATerrain

*Banana terrain requirement for land evaluation*

---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Banana.

**Format**

A data frame with 6 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization. )
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage - Drainage
- SlopeD - Slope (degree, 6 classes)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

BANANAWater*Banana water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Banana.

**Format**

A data frame with 2 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)
- WmDryLen - Length dry season (months :  $P < 1/2$  PET)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

BARLEYSoil

*Barley soil requirement for land evaluation*

---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Barley.

**Format**

A data frame with 13 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragn - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC6 - Organic carbon (%) - Kaolinitic materials
- OC7 - Organic carbon (%) - Non Kaolinitic, Non calcareous materials
- OC8 - Organic carbon (%) - Calcareous materials
- ECedS - ECe (dS/m)
- ESP - ESP (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 BARLEYTemp

*Barley temp requirement for land evaluation*


---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Barley.

**Format**

A data frame with 3 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TmAv2 - Mean temp. crop development stage (2nd month) (°C)
- TmAv3 - Mean temp. of the flowering stage (°C)
- TmAv4 - Mean temp. of the ripening stage (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 BARLEYTerrain

*Barley terrain requirement for land evaluation*


---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Barley.

**Format**

A data frame with 6 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization. )
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

BARLEYWater

*Barley water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Barley.

**Format**

A data frame with 3 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WmAv2 - Mean precipitation of second month (mm)
- WmAv3 - Mean precipitation of third month (mm)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

BEANCASoil

*Castor Beans soil requirement for land evaluation*

---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Castor Beans.

**Format**

A data frame with 9 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

BEANCATemp

*Castor Beans temp requirement for land evaluation*

---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Castor Beans.

**Format**

A data frame with 2 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TmAv1 - Mean temp. of the initial stage( C )

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

BEANCATerrain

*Castor Beans terrain requirement for land evaluation*

---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Castor Beans.

**Format**

A data frame with 6 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization. )
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

BEANCAWater

*Castor Beans water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Castor Beans.

**Format**

A data frame with 2 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WmhAv3 - Relative humidity of maturation Stage (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

BEANSSoil

*Beans soil requirement for land evaluation*

---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Beans.

**Format**

A data frame with 12 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragment - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

BEANSTemp

*Beans temp requirement for land evaluation*

---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Beans.

**Format**

A data frame with 2 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TgMinAv - Mean min. temp. of growing cycle (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

BEANSTerrain

*Beans terrain requirement for land evaluation*

---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Beans.

**Format**

A data frame with 7 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of managemnet with full mechanization. )
- Slope3 - Slope (%) (3. Low level of managemnet animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- SlopeD - Slope (degree, 6 classes)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 BEANSWater

*Beans water requirement for land evaluation*


---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Beans.

**Format**

A data frame with 5 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WmhAv2 - Relative humidity of devel. Stage (%)
- WmhAv3 - Relative humidity of maturation Stage (%)
- WmnN2 - n/N develop. Stage (2nd month)
- WmnN4 - n/N maturation stage (4th month)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 CABBAGESoil

*Cabbage soil requirement for land evaluation*


---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Cabbage.

**Format**

A data frame with 12 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragment - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

CABBAGETemp

*Cabbage temp requirement for land evaluation*

---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Cabbage.

**Format**

A data frame with 3 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TmAv0 - Mean temp. at germination (°C) (1st month)
- TdDiff - Temp difference day/night ( C )

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

CABBAGETerrain      *Cabbage terrain requirement for land evaluation*

---

### **Description**

A dataset containing the terrain characteristics of the crop requirements for farming Cabbage.

### **Format**

A data frame with 7 rows and 8 columns

### **Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization. )
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- SlopeD - Slope (degree, 6 classes)

### **See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

CABBAGEWater      *Cabbage water requirement for land evaluation*

---

### **Description**

A dataset containing the water characteristics of the crop requirements for farming Cabbage.

### **Format**

A data frame with 2 rows and 8 columns

### **Details**

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WghAv - Relative humidity growing cycle (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

CARROTSSoil

*Carrots soil requirement for land evaluation*

---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Carrots.

**Format**

A data frame with 11 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 CARROTSTemp

*Carrots temp requirement for land evaluation*


---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Carrots.

**Format**

A data frame with 2 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TmAv0 - Mean temp. at germination (°C) (1st month)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 CARROTSTerrain

*Carrots terrain requirement for land evaluation*


---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Carrots.

**Format**

A data frame with 6 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization. )
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

CARROTSWater

*Carrots water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Carrots.

**Format**

A data frame with 2 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WghAv - Relative humidity growing cycle (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

CASHEWSoil

*Cashew soil requirement for land evaluation*

---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Cashew.

**Format**

A data frame with 8 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

CASHEWTemp

*Cashew temp requirement for land evaluation*

---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Cashew.

**Format**

A data frame with 2 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TyMaxAv - Mean annual maximum temperature (°C)
- TmMinXm - Avarage minimum temperature of coldest month ( C )

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

CASHEWTerrain

*Cashew terrain requirement for land evaluation*

---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Cashew.

**Format**

A data frame with 8 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization. )
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- SlopeD - Slope (degree, 6 classes)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

CASHEWWater

*Cashew water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Cashew.

**Format**

A data frame with 2 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)
- WmDryLen - Length dry season (months :  $P < 1/2$  PET)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

CASSAVASoil

*Cassava soil requirement for land evaluation*

---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Cassava.

**Format**

A data frame with 12 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragm1 - Coarse fragment in surface (Vol.%)
- CFragm2 - Coarse fragment in depth (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

CASSAVATemp

*Cassava temp requirement for land evaluation*

---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Cassava.

**Format**

A data frame with 3 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TyMaxAv - Mean annual maximum temperature (°C)
- TmMinXmAb - Absolute min temp. coldest month (°C)
- TgAv - Mean temperature of the growing cycle (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

CASSAVATerrain

*Cassava terrain requirement for land evaluation*

---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Cassava.

**Format**

A data frame with 6 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization. )
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage - Drainage
- SlopeD - Slope (degree, 6 classes)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 CASSAVAWater

*Cassava water requirement for land evaluation*


---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Cassava.

**Format**

A data frame with 3 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)
- WmDryLen - Length dry season (months :  $P < 1/2$  PET)
- WmnN5 - n/N of the 5 dryest months

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 CHICKPEASoil

*Chickpea soil requirement for land evaluation*


---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Chickpea.

**Format**

A data frame with 11 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

CHICKPEATemp

*Chickpea temp requirement for land evaluation*

---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Chickpea.

**Format**

A data frame with 1 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

CHICKPEATerrain

*Chickpea terrain requirement for land evaluation*

---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Chickpea.

**Format**

A data frame with 8 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization. )
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- SlopeD - Slope (degree, 6 classes)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

CHICKPEAWater

*Chickpea water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Chickpea.

**Format**

A data frame with 3 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WmAv3 - Mean precipitation of third month (mm)
- WmAv4 - Mean precipitation of fourth month (mm)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

CINNAMONSoil

*Cinnamon soil requirement for land evaluation*

---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Cinnamon.

**Format**

A data frame with 9 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

CINNAMONTemp	<i>Cinnamon temp requirement for land evaluation</i>
--------------	--

---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Cinnamon.

**Format**

A data frame with 1 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

CINNAMONTerrain	<i>Cinnamon terrain requirement for land evaluation</i>
-----------------	---

---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Cinnamon.

**Format**

A data frame with 2 rows and 8 columns

**Details**

The following are the factors for evaluation:

- SlopeD - Slope (degree, 6 classes)
- Flood - Flooding

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

CINNAMONWater

*Cinnamon water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Cinnamon.

**Format**

A data frame with 2 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WghAv - Relative humidity growing cycle (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

CITRUSSoil

*Citrus soil requirement for land evaluation*

---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Citrus.

**Format**

A data frame with 12 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)

- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

#### See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

CITRUSTemp

*Citrus temp requirement for land evaluation*

---

#### Description

A dataset containing the temp characteristics of the crop requirements for farming Citrus.

#### Format

A data frame with 8 rows and 8 columns

#### Details

The following are the factors for evaluation:

- TyAv - Mean annual temperature (°C)
- TmMax38 - No of months with mean temp. > 38 °C
- TmMin13 - No of months with mean temp. < 13 °C
- TyMinAb - Absolute minimum temperature (°C)
- TyMinAb - Absolute minimum temperature (°C)
- TyMinAb - Absolute minimum temperature (°C)
- TyMinAb - Absolute minimum temperature (°C)
- TmAv3 - Mean temp. of the flowering stage (°C)

#### See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

CITRUSTerrain

*Citrus terrain requirement for land evaluation*

---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Citrus.

**Format**

A data frame with 6 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization. )
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage - Drainage
- SlopeD - Slope (degree, 6 classes)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

CITRUSWater

*Citrus water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Citrus.

**Format**

A data frame with 5 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)
- WmDryLen - Length dry season (months :  $P < 1/2$  PET)
- WmhColdXm - Relative humidity of coldest month if frost (%)
- WmhAv4 - Relative humidity at harvest stage (%)
- WmhAv3 - Relative humidity of maturation Stage (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

COCOASoil

*Cocoa soil requirement for land evaluation*

---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Cocoa.

**Format**

A data frame with 10 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragment - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pH<sub>H2O</sub> - pH H<sub>2O</sub>
- OC - Organic carbon (%)
- EC<sub>dS</sub> - EC<sub>e</sub> (dS/m)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

COCOATemp

*Cocoa temp requirement for land evaluation*

---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Cocoa.

**Format**

A data frame with 3 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TyAv - Mean annual temperature (°C)
- TyMaxAv - Mean annual maximum temperature (°C)
- TyMaxAv - Mean annual maximum temperature (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

COCOATerrain

*Cocoa terrain requirement for land evaluation*

---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Cocoa.

**Format**

A data frame with 5 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization. )
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage - Drainage

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 COCOAWater

*Cocoa water requirement for land evaluation*


---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Cocoa.

**Format**

A data frame with 4 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)
- WmDryLen - Length dry season (months :  $P < 1/2$  PET)
- WmhDryXm - Mean rel. humidity dryest month (%)
- WmhDryXm - Mean rel. humidity dryest month (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 COCONUTSoil

*Coconut soil requirement for land evaluation*


---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Coconut.

**Format**

A data frame with 6 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- OC - Organic carbon (%)
- ECemh - ECe (mmhos/cm)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

COCONUTTemp

*Coconut temp requirement for land evaluation*

---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Coconut.

**Format**

A data frame with 1 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TyAv - Mean annual temperature (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

COCONUTTerrain      *Coconut terrain requirement for land evaluation*

---

### **Description**

A dataset containing the terrain characteristics of the crop requirements for farming Coconut.

### **Format**

A data frame with 5 rows and 8 columns

### **Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization. )
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage - Drainage

### **See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

COCONUTWater      *Coconut water requirement for land evaluation*

---

### **Description**

A dataset containing the water characteristics of the crop requirements for farming Coconut.

### **Format**

A data frame with 3 rows and 8 columns

### **Details**

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)
- WmDryLen - Length dry season (months :  $P < 1/2$  PET)
- WyhAv - Mean annual rel. humidity (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

COFFEEARSoil

*Arabica Coffee soil requirement for land evaluation*

---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Arabica Coffee.

**Format**

A data frame with 11 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 COFFEEARTemp

*Arabica Coffee temp requirement for land evaluation*


---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Arabica Coffee.

**Format**

A data frame with 3 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TyMaxAv - Mean annual maximum temperature (°C)
- TdMinXm - Mean daily minimum temperature of coldest month (°C)
- TyAv - Mean annual temperature (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 COFFEEARTerrain

*Arabica Coffee terrain requirement for land evaluation*


---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Arabica Coffee.

**Format**

A data frame with 6 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization. )
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage - Drainage
- SlopeD - Slope (degree, 6 classes)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

COFFEEARWater

*Arabica Coffee water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Arabica Coffee.

**Format**

A data frame with 4 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)
- WmDryLen - Length dry season (months :  $P < 1/2$  PET)
- WmhDryXm - Mean rel. humidity dryest month (%)
- WmnN5 - n/N of the 5 dryest months

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

COFFEEROSoil

*Robusta Coffee soil requirement for land evaluation*

---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Robusta Coffee.

**Format**

A data frame with 10 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

COFFEEROTemp

*Robusta Coffee temp requirement for land evaluation*

---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Robusta Coffee.

**Format**

A data frame with 3 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TyAv - Mean annual temperature (°C)
- TyMaxAv - Mean annual maximum temperature (°C)
- TdMinXm - Mean daily minimum temperature of coldest month (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

COFFEEROTerrain	<i>Robusta Coffee terrain requirement for land evaluation</i>
-----------------	---

---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Robusta Coffee.

**Format**

A data frame with 5 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization. )
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage - Drainage

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

COFFEEROWater	<i>Robusta Coffee water requirement for land evaluation</i>
---------------	---

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Robusta Coffee.

**Format**

A data frame with 4 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)
- WmDryLen - Length dry season (months :  $P < 1/2$  PET)
- WmhDryXm - Mean rel. humidity dryest month (%)
- WmnN5 - n/N of the 5 dryest months

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

COTTONSoil

*Cotton soil requirement for land evaluation*

---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Cotton.

**Format**

A data frame with 13 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragn - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC6 - Organic carbon (%) - Kaolinitic materials
- OC7 - Organic carbon (%) - Non Kaolinitic, Non calcareous materials
- OC8 - Organic carbon (%) - Calcareous materials
- ECemh - ECe (mmhos/cm)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

COTTONTemp

*Cotton temp requirement for land evaluation*

---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Cotton.

**Format**

A data frame with 6 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TgMaxAv - Mean max temp. of growing cycle (°C)
- TmMaxXm - Average max. temp. warmest month (°C)
- TmAv2 - Mean temp. crop development stage (2nd month) (°C)
- TdAvg3 - Mean DAY temp. of flowering stage (°C)
- TdMinN3 - Mean Night temp. of flowering stage (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

COTTONTerrain

*Cotton terrain requirement for land evaluation*

---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Cotton.

**Format**

A data frame with 7 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization. )
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- Slope - nan

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

COTTONWater

*Cotton water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Cotton.

**Format**

A data frame with 6 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WmAv1 - Mean precipitation of first month (mm)
- WmAv2 - Mean precipitation of second month (mm)
- WmAv5 - Mean precipitation of fifth month (mm)
- WmAv6 - Precipitation of ripening stage (mm)(6th month)
- WmhAv3 - Relative humidity of maturation Stage (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

COWPEASoil

*Cowpea soil requirement for land evaluation*

---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Cowpea.

**Format**

A data frame with 11 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragn - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

COWPEATemp

*Cowpea temp requirement for land evaluation*

---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Cowpea.

**Format**

A data frame with 3 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TmAv0 - Mean temp. at germination (°C) (1st month)
- TyMinAv - Mean annual minimum temperature (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

COWPEATerrain

*Cowpea terrain requirement for land evaluation*

---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Cowpea.

**Format**

A data frame with 7 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization. )
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- SlopeD - Slope (degree, 6 classes)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

COWPEAWater

*Cowpea water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Cowpea.

**Format**

A data frame with 6 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WmAv1 - Mean precipitation of first month (mm)
- WmAv2 - Mean precipitation of second month (mm)
- WmAv3 - Mean precipitation of third month (mm)
- WmAv4 - Mean precipitation of fourth month (mm)
- WmhAv4 - Relative humidity at harvest stage (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

CUCUMBERSoil

*Cucumber soil requirement for land evaluation*

---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Cucumber.

**Format**

A data frame with 10 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

CUCUMBERTemp

*Cucumber temp requirement for land evaluation*

---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Cucumber.

**Format**

A data frame with 1 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

CUCUMBERTerrain      *Cucumber terrain requirement for land evaluation*

---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Cucumber.

**Format**

A data frame with 7 rows and 8 columns

**Details**

The following are the factors for evaluation:

- SlopeD - Slope (degree, 6 classes)
- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization. )
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

CUCUMBERWater      *Cucumber water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Cucumber.

**Format**

A data frame with 2 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WghAv - Relative humidity growing cycle (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

GROUNDNUTSSoil

*Groundnuts soil requirement for land evaluation*

---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Groundnuts.

**Format**

A data frame with 15 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECs - Apparent CEC Soil (cmol (+)/kg soil)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC6 - Organic carbon (%) - Kaolinitic materials
- OC7 - Organic carbon (%) - Non Kaolinitic, Non calcareous materials
- OC8 - Organic carbon (%) - Calcareous materials
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- OC - Organic carbon (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

GROUNDNUTSTemp

*Groundnuts temp requirement for land evaluation*

---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Groundnuts.

**Format**

A data frame with 3 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TgMaxAv - Mean max temp. of growing cycle (°C)
- TgMinAv - Mean min. temp. of growing cycle (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

GROUNDNUTSTerrain

*Groundnuts terrain requirement for land evaluation*

---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Groundnuts.

**Format**

A data frame with 7 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization. )
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- SlopeD - Slope (degree, 6 classes)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 GROUNDNUTSWater

*Groundnuts water requirement for land evaluation*


---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Groundnuts.

**Format**

A data frame with 6 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WmAv1 - Mean precipitation of first month (mm)
- WmAv2 - Mean precipitation of second month (mm)
- WmAv3 - Mean precipitation of third month (mm)
- WmAv4 - Mean precipitation of fourth month (mm)
- WghAv - Relative humidity growing cycle (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 GUAVASoil

*Guava soil requirement for land evaluation*


---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Guava.

**Format**

A data frame with 9 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

GUAVATemp

*Guava temp requirement for land evaluation*

---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Guava.

**Format**

A data frame with 1 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TyAv - Mean annual temperature (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 GUAVATerrain

*Guava terrain requirement for land evaluation*


---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Guava.

**Format**

A data frame with 5 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of managemnet with full mechanization. )
- Slope3 - Slope (%) (3. Low level of managemnet animal traction or handwork.)
- Flood - Flooding
- Drainage - Drainage

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 GUAVAWater

*Guava water requirement for land evaluation*


---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Guava.

**Format**

A data frame with 1 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

LaoCaiLT

*Land and Terrain Characteristics of Lao Cai, Vietnam*

---

**Description**

A dataset containing the land and terrain characteristics of the land units in Lao Cai, Vietnam.

**Format**

A data frame with 2928 rows and 10 columns

**Details**

- SlopeD - Slope (degree, 6 classes);
- CFragm - Coarse fragment (Vol.%);
- SoilDpt - Soil depth (cm);
- SoilTe - 12 classes of soil texture (Soil Taxonomy);
- CECc - Apparent CEC Clay (cmol (+)/kg clay);
- SumBCs - Sum of basic cations (cmol (+)/kg soil);
- pHH2O - pH H2O;
- BS - Base Saturation (%);
- OC - Organic carbon (%);
- Flood - Flooding;

---

LaoCaiTemp

*Temperature Characteristics of Lao Cai, Vietnam*

---

**Description**

A dataset containing the temperature characteristics of the land units in Lao Cai, Vietnam.

**Format**

A data frame with 2928 rows and 12 columns

**Details**

- Jan - January;
- Feb - February;
- Mar - March;
- Apr - April;
- May - May;
- Jun - June;
- Jul - July;
- Aug - August;
- Sep - September;
- Oct - October;
- Nov - November;
- Dec - December;

---

LaoCaiWater

*Water Characteristics of Lao Cai, Vietnam*

---

**Description**

A dataset containing the water characteristics of the land units in Lao Cai, Vietnam.

**Format**

A data frame with 2928 rows and 12 columns

**Details**

- Jan - January;
- Feb - February;
- Mar - March;
- Apr - April;
- May - May;
- Jun - June;
- Jul - July;
- Aug - August;
- Sep - September;
- Oct - October;
- Nov - November;
- Dec - December;

---

LONGANSoil

*Longan soil requirement for land evaluation*

---

### **Description**

A dataset containing the soil characteristics of the crop requirements for farming Longan.

### **Format**

A data frame with 12 rows and 8 columns

### **Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

### **See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

LONGANTemp

*Longan temp requirement for land evaluation*


---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Longan.

**Format**

A data frame with 3 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TyAv - Mean annual temperature (°C)
- TdMinXm - Mean daily minimum temperature of coldest month (°C)
- TyMinAv - Mean annual minimum temperature (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

LONGANTerrain

*Longan terrain requirement for land evaluation*


---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Longan.

**Format**

A data frame with 6 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization. )
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage - Drainage
- SlopeD - Slope (degree, 6 classes)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

LONGANWater

*Longan water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Longan.

**Format**

A data frame with 1 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

MAIZESoil

*Maize soil requirement for land evaluation*

---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Maize.

**Format**

A data frame with 15 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC6 - Organic carbon (%) - Kaolinitic materials
- OC7 - Organic carbon (%) - Non Kaolinitic, Non calcareous materials
- OC8 - Organic carbon (%) - Calcareous materials
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- OC - Organic carbon (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

MAIZETemp

*Maize temp requirement for land evaluation*

---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Maize.

**Format**

A data frame with 2 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TgMinAv - Mean min. temp. of growing cycle (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

MAIZETerrain

*Maize terrain requirement for land evaluation*

---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Maize.

**Format**

A data frame with 7 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization. )
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- SlopeD - Slope (degree, 6 classes)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

MAIZEWATER

*Maize water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Maize.

**Format**

A data frame with 9 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WmAv1 - Mean precipitation of first month (mm)
- WmAv2 - Mean precipitation of second month (mm)
- WmAv3 - Mean precipitation of third month (mm)
- WmAv4 - Mean precipitation of fourth month (mm)
- WmhAv2 - Relative humidity of devel. Stage (%)
- WmhAv3 - Relative humidity of maturation Stage (%)
- WmnN2 - n/N develop. Stage (2nd month)
- WmnN4 - n/N maturation stage (4th month)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

MANGOSoil

*Mango soil requirement for land evaluation*


---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Mango.

**Format**

A data frame with 12 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCc - Sum of basic cations (cmol (+)/kg of clay)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

MANGOTemp

*Mango temp requirement for land evaluation***Description**

A dataset containing the temp characteristics of the crop requirements for farming Mango.

**Format**

A data frame with 2 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TyAv - Mean annual temperature (°C)
- TyMinAv - Mean annual minimum temperature (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

MANGOTerrain

*Mango terrain requirement for land evaluation***Description**

A dataset containing the terrain characteristics of the crop requirements for farming Mango.

**Format**

A data frame with 6 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1) Basin furrow irrigation
- Slope2 - Slope (%) (2) Mechanized, high management level
- Slope3 - Slope (%) (3) Manual, low management level
- Flood - Flooding
- Drainage - Drainage
- SlopeD - 6 classes of slope (Degree)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

MANGOWater

*Mango water requirement for land evaluation*


---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Mango.

**Format**

A data frame with 5 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)
- WmDryLen - Length dry season (months :  $P < 1/2$  PET)
- WmAvDry - Monthly precipitation during dry season (mm)
- WyhAv - Mean annual rel. humidity (%)
- WmnN4 - n/N maturation stage (4th month)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

MarinduqueLT

*Land and Terrain Characteristics of Marinduque, Philippines*


---

**Description**

A dataset containing the land and terrain characteristics of the land units in Marinduque, Philippines.

**Format**

A data frame with 881 rows and 6 columns

**Details**

- Lat - Latitude of Land Units;
- Lon - Longitude of Land Units;
- CECc - Apparent CEC Clay (cmol (+)/kg clay);
- pHH2O - pH H2O;
- CFragm - Coarse fragment (Vol.%);
- SoilTe - 12 classes of soil texture (Soil Taxonomy);

---

MarinduqueTemp

*Temperature Characteristics of Marinduque, Philippines*

---

**Description**

A dataset containing the temperature characteristics of the land units in Marinduque, Philippines.

**Format**

A data frame with 881 rows and 14 columns

**Details**

- Lat - Latitude of Land Units;
- Lon - Longitude of Land Units;
- Jan - January;
- Feb - February;
- Mar - March;
- Apr - April;
- May - May;
- Jun - June;
- Jul - July;
- Aug - August;
- Sep - September;
- Oct - October;
- Nov - November;
- Dec - December;

---

MarinduqueWater

*Water Characteristics of Marinduque, Philippines*

---

**Description**

A dataset containing the water characteristics of the land units in Marinduque, Philippines.

**Format**

A data frame with 881 rows and 14 columns

**Details**

- Lat - Latitude of Land Units;
- Lon - Longitude of Land Units;
- Jan - January;
- Feb - February;
- Mar - March;
- Apr - April;
- May - May;
- Jun - June;
- Jul - July;
- Aug - August;
- Sep - September;
- Oct - October;
- Nov - November;
- Dec - December;

---

MILLETSSoil

*Millets soil requirement for land evaluation*

---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Millets.

**Format**

A data frame with 11 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC6 - Organic carbon (%) - Kaolinitic materials
- ECedS - ECe (dS/m)
- ESP - ESP (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

MILLETSTemp

*Millets temp requirement for land evaluation*


---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Millets.

**Format**

A data frame with 3 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TgMaxAv - Mean max temp. of growing cycle (°C)
- TgMinAv - Mean min. temp. of growing cycle (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

MILLETSTerrain

*Millets terrain requirement for land evaluation*

---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Millets.

**Format**

A data frame with 6 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization. )
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.
- 

MILLETWater

*Millets water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Millets.

**Format**

A data frame with 5 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WmAv1 - Mean precipitation of first month (mm)
- WmAv2 - Mean precipitation of second month (mm)
- WmAv4 - Mean precipitation of fourth month (mm)
- WmhAv4 - Relative humidity at harvest stage (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

OILPALMSoil

*Oil Palm soil requirement for land evaluation*

---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Oil Palm.

**Format**

A data frame with 10 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragm2 - Coarse fragment in depth (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 OILPALMTemp

*Oil Palm temp requirement for land evaluation*


---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Oil Palm.

**Format**

A data frame with 3 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TyMaxAv - Mean annual maximum temperature (°C)
- TdMinXm - Mean daily minimum temperature of coldest month (°C)
- TyAv - Mean annual temperature (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 OILPALMTerrain

*Oil Palm terrain requirement for land evaluation*


---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Oil Palm.

**Format**

A data frame with 4 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope - Slope
- Flood - Flooding
- Drainage4 - Drainage (4) Medium & fine textured soils
- Drainage5 - Drainage (5) Coarse textured soils

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

OILPALMWater

*Oil Palm water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Oil Palm.

**Format**

A data frame with 3 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)
- WmDryLen - Length dry season (months :  $P < 1/2$  PET)
- WynN - Mean annual n/N

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

OLIVESSoil

*Olives soil requirement for land evaluation*

---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Olives.

**Format**

A data frame with 10 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragment - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)

- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pH<sub>H2O</sub> - pH H<sub>2</sub>O
- OC - Organic carbon (%)
- EC<sub>dS</sub> - EC<sub>e</sub> (dS/m)
- ESP - ESP (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

OLIVESTemp

*Olives temp requirement for land evaluation*

---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Olives.

**Format**

A data frame with 2 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Ty<sub>Av</sub> - Mean annual temperature (°C)
- Tm<sub>MinXm</sub> - Average minimum temperature of coldest month ( C )

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

OLIVESTerrain

*Olives terrain requirement for land evaluation*

---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Olives.

**Format**

A data frame with 6 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization. )
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage - Drainage
- SlopeD - Slope (degree, 6 classes)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

OLIVESWater

*Olives water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Olives.

**Format**

A data frame with 3 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)
- WmSpecial1 - Monthly rainfall during the sclerification of stone (mm) - August (N hem) February (S hem.)
- WmSpecial2 - Monthly rainfall during the sclerification of stone (mm) - September (N hem) March (S hem.)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

ONIONSoil

*Onion soil requirement for land evaluation*

---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Onion.

**Format**

A data frame with 11 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragment - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 ONIONTemp

*Onion temp requirement for land evaluation*


---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Onion.

**Format**

A data frame with 2 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TmAv0 - Mean temp. at germination (°C) (1st month)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 ONIONTerrain

*Onion terrain requirement for land evaluation*


---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Onion.

**Format**

A data frame with 6 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization. )
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 ONIONWater

*Onion water requirement for land evaluation*


---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Onion.

**Format**

A data frame with 2 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- TmAvDlen3 - Daylength (h) during yield form. Period

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 overall\_suit

*Overall Suitability Scores/Class of the Land Units*


---

**Description**

This function computes the overall suitability scores and class of the land units.

**Usage**

```
overall_suit(suit, method = NULL, interval = NULL)
```

**Arguments**

suit	an object of class suitability.
method	a character for the method for computing the overall suitability, choices are: "minimum", "maximum", and "average". If NULL, method is set to "minimum".
interval	if NULL, the interval of the suitability class are the following: 0% - 25% (Not suitable, N), 25% - 50% (Marginally Suitable, S3), 50% - 75% (Moderately Suitable, S2), and 75% - 100% (Highly Suitable, S1). But users can assign custom intervals by specifying the values of the end points of the intervals. Say for intervals: 0% - 20% (Not suitable, N), 20% - 50% (Marginally Suitable, S3), 50% - 80% (Moderately Suitable, S2), and 80% - 100% (Highly Suitable, S1), is equivalent to interval = c(0, 0.2, 0.5, 0.8, 1).

**Value**

A data frame with columns:

- Score - the overall suitability scores
- Class - the overall suitability classes

**See Also**

[suit](https://alstat.github.io/ALUES/), <https://alstat.github.io/ALUES/>

**Examples**

```
library(ALUES)
out <- suit("ricebr", terrain=MarinduqueLT, water=MarinduqueWater, temp=MarinduqueTemp, sow_month=1)
out[["terrain"]]

# Soil Overall Suitability
head(overall_suit(out[["soil"]]))
head(overall_suit(out[["soil"]], "average"))
head(overall_suit(out[["soil"]], "maximum"))
head(overall_suit(out[["soil"]], "average", c(0, 0.3, 0.35, 0.6, 1.0)))

# Water Overall Suitability
head(overall_suit(out[["water"]], "average"))
head(overall_suit(out[["water"]], "maximum"))
head(overall_suit(out[["water"]], "average", c(0, 0.3, 0.35, 0.6, 1.0)))

# Temperature Overall Suitability
head(overall_suit(out[["temp"]], "average"))
head(overall_suit(out[["temp"]], "maximum"))
head(overall_suit(out[["temp"]], "average", c(0, 0.3, 0.35, 0.6, 1.0)))
```

---

PAPAYASoil

*Papaya soil requirement for land evaluation*

---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Papaya.

**Format**

A data frame with 9 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

PAPAYATemp

*Papaya temp requirement for land evaluation*

---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Papaya.

**Format**

A data frame with 2 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TyAv - Mean annual temperature (°C)
- TyMinAv - Mean annual minimum temperature (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

PAPAYATerrain

*Papaya terrain requirement for land evaluation*

---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Papaya.

**Format**

A data frame with 0 rows and 8 columns

**Details**

The following are the factors for evaluation:

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

PAPAYAWater

*Papaya water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Papaya.

**Format**

A data frame with 3 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)
- WynN - Mean annual n/N
- WyhAv - Mean annual rel. humidity (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

PEACHSoil

*Peach soil requirement for land evaluation*

---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Peach.

**Format**

A data frame with 12 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

PEACHTemp

*Peach temp requirement for land evaluation*

---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Peach.

**Format**

A data frame with 1 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TyAv - Mean annual temperature (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

PEACHTerrain

*Peach terrain requirement for land evaluation*

---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Peach.

**Format**

A data frame with 6 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of managemnet with full mechanization. )
- Slope3 - Slope (%) (3. Low level of managemnet animal traction or handwork.)
- Flood - Flooding
- Drainage - Drainage
- SlopeD - Slope (degree, 6 classes)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

PEACHWater

*Peach water requirement for land evaluation*

---

### **Description**

A dataset containing the water characteristics of the crop requirements for farming Peach.

### **Format**

A data frame with 1 rows and 8 columns

### **Details**

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)

### **See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

PEARSoil

*Pear soil requirement for land evaluation*

---

### **Description**

A dataset containing the soil characteristics of the crop requirements for farming Pear.

### **Format**

A data frame with 12 rows and 8 columns

### **Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)

- pH<sub>H2O</sub> - pH H<sub>2</sub>O
- OC - Organic carbon (%)
- EC<sub>dS</sub> - EC<sub>e</sub> (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

#### See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

PEARTemp

*Pear temp requirement for land evaluation*

---

#### Description

A dataset containing the temp characteristics of the crop requirements for farming Pear.

#### Format

A data frame with 1 rows and 8 columns

#### Details

The following are the factors for evaluation:

- TyAv - Mean annual temperature (°C)

#### See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

PEARTerrain

*Pear terrain requirement for land evaluation*

---

#### Description

A dataset containing the terrain characteristics of the crop requirements for farming Pear.

#### Format

A data frame with 6 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization. )
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage - Drainage
- SlopeD - Slope (degree, 6 classes)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

PEARWater

*Pear water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Pear.

**Format**

A data frame with 1 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

PEASoil

*Pea soil requirement for land evaluation*

---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Pea.

**Format**

A data frame with 11 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragn - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

PEATemp

*Pea temp requirement for land evaluation*

---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Pea.

**Format**

A data frame with 2 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TmMinAv0 - Mean min. temp. at germination (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

PEATerrain

*Pea terrain requirement for land evaluation*

---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Pea.

**Format**

A data frame with 7 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of managemnet with full mechanization. )
- Slope3 - Slope (%) (3. Low level of managemnet animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- SlopeD - Slope (degree, 6 classes)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

PEAWater

*Pea water requirement for land evaluation*

---

### **Description**

A dataset containing the water characteristics of the crop requirements for farming Pea.

### **Format**

A data frame with 1 rows and 8 columns

### **Details**

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)

### **See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

PEPPERGRSoil

*Green Pepper soil requirement for land evaluation*

---

### **Description**

A dataset containing the soil characteristics of the crop requirements for farming Green Pepper.

### **Format**

A data frame with 11 rows and 8 columns

### **Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic caions (cmol (+)/kg soil)

- pHH<sub>2</sub>O - pH H<sub>2</sub>O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

#### See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

PEPPERGRTemp

*Green Pepper temp requirement for land evaluation*

---

#### Description

A dataset containing the temp characteristics of the crop requirements for farming Green Pepper.

#### Format

A data frame with 2 rows and 8 columns

#### Details

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TmAv0 - Mean temp. at germination (°C) (1st month)

#### See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

PEPPERGRTerrain

*Green Pepper terrain requirement for land evaluation*

---

#### Description

A dataset containing the terrain characteristics of the crop requirements for farming Green Pepper.

#### Format

A data frame with 7 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization. )
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- Slope - nan

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

PEPPERGRWater

*Green Pepper water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Green Pepper.

**Format**

A data frame with 1 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

PERSIMMONSoil

*Persimmon soil requirement for land evaluation*

---

### **Description**

A dataset containing the soil characteristics of the crop requirements for farming Persimmon.

### **Format**

A data frame with 12 rows and 8 columns

### **Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

### **See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

PERSIMMONTemp      *Persimmon temp requirement for land evaluation*

---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Persimmon.

**Format**

A data frame with 1 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

PERSIMMONTerrain      *Persimmon terrain requirement for land evaluation*

---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Persimmon.

**Format**

A data frame with 6 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of managemnet with full mechanization. )
- Slope3 - Slope (%) (3. Low level of managemnet animal traction or handwork.)
- Flood - Flooding
- Drainage - Drainage
- SlopeD - Slope (degree, 6 classes)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

PERSIMMONWater      *Persimmon water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Persimmon.

**Format**

A data frame with 1 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

PINEAPPLESoil      *Pineapple soil requirement for land evaluation*

---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Pineapple.

**Format**

A data frame with 12 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic caions (cmol (+)/kg soil)

- pH<sub>H2O</sub> - pH H<sub>2</sub>O
- OC - Organic carbon (%)
- EC<sub>dS</sub> - EC<sub>e</sub> (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

#### See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

PINEAPPLETemp

*Pineapple temp requirement for land evaluation*

---

#### Description

A dataset containing the temp characteristics of the crop requirements for farming Pineapple.

#### Format

A data frame with 1 rows and 8 columns

#### Details

The following are the factors for evaluation:

- TyAv - Mean annual temperature (°C)

#### See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

PINEAPPLETerrain

*Pineapple terrain requirement for land evaluation*

---

#### Description

A dataset containing the terrain characteristics of the crop requirements for farming Pineapple.

#### Format

A data frame with 7 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization. )
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- SlopeD - Slope (degree, 6 classes)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

PINEAPPLEWater

*Pineapple water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Pineapple.

**Format**

A data frame with 2 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)
- WyhAv - Mean annual rel. humidity (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

PLUMSoil

*Plum soil requirement for land evaluation*

---

### **Description**

A dataset containing the soil characteristics of the crop requirements for farming Plum.

### **Format**

A data frame with 12 rows and 8 columns

### **Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

### **See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

PLUMTemp                      *Plum temp requirement for land evaluation*

---

### Description

A dataset containing the temp characteristics of the crop requirements for farming Plum.

### Format

A data frame with 1 rows and 8 columns

### Details

The following are the factors for evaluation:

- TyAv - Mean annual temperature (°C)

### See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

PLUMTerrain                      *Plum terrain requirement for land evaluation*

---

### Description

A dataset containing the terrain characteristics of the crop requirements for farming Plum.

### Format

A data frame with 6 rows and 8 columns

### Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of managemnet with full mechanization. )
- Slope3 - Slope (%) (3. Low level of managemnet animal traction or handwork.)
- Flood - Flooding
- Drainage - Drainage
- SlopeD - Slope (degree, 6 classes)

### See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

PLUMWater

*Plum water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Plum.

**Format**

A data frame with 1 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

POTATOSoil

*Potato soil requirement for land evaluation*

---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Potato.

**Format**

A data frame with 12 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt1 - Surface Soil Depth (cm)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)

- pH<sub>H2O</sub> - pH H<sub>2</sub>O
- OC - Organic carbon (%)
- EC<sub>dS</sub> - EC<sub>e</sub> (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

#### See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

POTATOSWSoil

*Sweet Potato soil requirement for land evaluation*

---

#### Description

A dataset containing the soil characteristics of the crop requirements for farming Sweet Potato.

#### Format

A data frame with 12 rows and 8 columns

#### Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO<sub>3</sub> - CaCO<sub>3</sub> (%)
- Gyps - Gypsum (%)
- CEC<sub>c</sub> - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pH<sub>H2O</sub> - pH H<sub>2</sub>O
- OC - Organic carbon (%)
- EC<sub>dS</sub> - EC<sub>e</sub> (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

#### See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 POTATOSWTemp

*Sweet Potato temp requirement for land evaluation*


---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Sweet Potato.

**Format**

A data frame with 2 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TgMinAv - Mean min. temp. of growing cycle (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 POTATOSWTerrain

*Sweet Potato terrain requirement for land evaluation*


---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Sweet Potato.

**Format**

A data frame with 7 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of managemnet with full mechanization. )
- Slope3 - Slope (%) (3. Low level of managemnet animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- SlopeD - Slope (degree, 6 classes)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 POTATOSWater

*Sweet Potato water requirement for land evaluation*


---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Sweet Potato.

**Format**

A data frame with 5 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)
- WmDryLen - Length dry season (months :  $P < 1/2$  PET)
- WmhAv4 - Relative humidity at harvest stage (%)
- WmnN2 - n/N develop. Stage (2nd month)
- WmnN4 - n/N maturation stage (4th month)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 POTATOTemp

*Potato temp requirement for land evaluation*


---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Potato.

**Format**

A data frame with 4 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TmMinAv1 - Average absolut Min. temperature of the first month (°C)
- TmMinAv4 - Average absolut Min. temperature of other months (°C)
- TdAvgDiff - Average Temperature difference between day-night (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 POTATOTerrain

*Potato terrain requirement for land evaluation*


---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Potato.

**Format**

A data frame with 7 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of managemnet with full mechanization. )
- Slope3 - Slope (%) (3. Low level of managemnet animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- SlopeD - Slope (degree, 6 classes)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

POTATOWater

*Potato water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Potato.

**Format**

A data frame with 5 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WmAv1 - Mean precipitation of first month (mm)
- WmAv2 - Mean precipitation of second month (mm)
- WmAv3 - Mean precipitation of third month (mm)
- WmAv4 - Mean precipitation of fourth month (mm)
- TgAvDlen - Average daylength growing cycle (h)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

RICEBRSoil

*Rainfed Bunded Rice soil requirement for land evaluation*

---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Rainfed Bunded Rice.

**Format**

A data frame with 6 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

RICEBRTemp

*Rainfed Banded Rice temp requirement for land evaluation*

---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Rainfed Banded Rice.

**Format**

A data frame with 4 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TmMaxXm - Average max. temp. warmest month (°C)
- TmAv2 - Mean temp. crop development stage (2nd month) (°C)
- TmMinAv4 - Average absolut Min. temperature of other months (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

RICEBRTerrain      *Rainfed Bunded Rice terrain requirement for land evaluation*

---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Rainfed Bunded Rice.

**Format**

A data frame with 3 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Flood - Flooding
- Drainage - Drainage

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

RICEBRWater      *Rainfed Bunded Rice water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Rainfed Bunded Rice.

**Format**

A data frame with 7 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WmAv1 - Mean precipitation of first month (mm)
- WmAv2 - Mean precipitation of second month (mm)
- WmAv3 - Mean precipitation of third month (mm)
- WmAv4 - Mean precipitation of fourth month (mm)

- WmhAv2 - Relative humidity of devel. Stage (%)
- WmhAv4 - Relative humidity at harvest stage (%)
- WynN - Mean annual n/N

#### See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 RICEIWSoil

*Irrigated Rice soil requirement for land evaluation*


---

#### Description

A dataset containing the soil characteristics of the crop requirements for farming Irrigated Rice.

#### Format

A data frame with 12 rows and 8 columns

#### Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- pHH2O - pH H2O
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

#### See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

RICEIWTemp

*Irrigated Rice temp requirement for land evaluation*


---

### Description

A dataset containing the temp characteristics of the crop requirements for farming Irrigated Rice.

### Format

A data frame with 4 rows and 8 columns

### Details

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TmMaxXm - Average max. temp. warmest month (°C)
- TmAv2 - Mean temp. crop development stage (2nd month) (°C)
- TmMinAv4 - Average absolut Min. temperature of other months (°C)

### See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

RICEIWTerrain

*Irrigated Rice terrain requirement for land evaluation*


---

### Description

A dataset containing the terrain characteristics of the crop requirements for farming Irrigated Rice.

### Format

A data frame with 4 rows and 8 columns

### Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Flood - Flooding
- Drainage - Drainage
- SlopeD - Slope (degree, 6 classes)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

RICEIWWater

*Irrigated Rice water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Irrigated Rice.

**Format**

A data frame with 7 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WmAv1 - Mean precipitation of first month (mm)
- WmAv2 - Mean precipitation of second month (mm)
- WmAv3 - Mean precipitation of third month (mm)
- WmAv4 - Mean precipitation of fourth month (mm)
- WmhAv2 - Relative humidity of devel. Stage (%)
- WmhAv4 - Relative humidity at harvest stage (%)
- WynN - Mean annual n/N

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

RICENFSoil	<i>Rice Cultivation under Natural Floods soil requirement for land evaluation</i>
------------	---

---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Rice Cultivation under Natural Floods.

**Format**

A data frame with 11 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

RICENFTemp	<i>Rice Cultivation under Natural Floods temp requirement for land evaluation</i>
------------	---

---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Rice Cultivation under Natural Floods.

**Format**

A data frame with 4 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TmMaxXm - Average max. temp. warmest month (°C)
- TmAv2 - Mean temp. crop development stage (2nd month) (°C)
- TmMinAv4 - Average absolut Min. temperature of other months (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

RICENFTerrain	<i>Rice Cultivation under Natural Floods terrain requirement for land evaluation</i>
---------------	--

---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Rice Cultivation under Natural Floods.

**Format**

A data frame with 2 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Flood - Flooding
- Drainage - Drainage

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

RICENFWater

*Rice Cultivation under Natural Floods water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Rice Cultivation under Natural Floods.

**Format**

A data frame with 7 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WmAv1 - Mean precipitation of first month (mm)
- WmAv2 - Mean precipitation of second month (mm)
- WmAv3 - Mean precipitation of third month (mm)
- WmAv4 - Mean precipitation of fourth month (mm)
- WmhAv2 - Relative humidity of devel. Stage (%)
- WmhAv4 - Relative humidity at harvest stage (%)
- WynN - Mean annual n/N

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

RICEURSoil

*Rainfed Upland Rice soil requirement for land evaluation*

---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Rainfed Upland Rice.

**Format**

A data frame with 8 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

RICEURTemp

*Rainfed Upland Rice temp requirement for land evaluation*

---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Rainfed Upland Rice.

**Format**

A data frame with 4 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TmMaxXm - Average max. temp. warmest month (°C)
- TmAv2 - Mean temp. crop development stage (2nd month) (°C)
- TmMinAv4 - Average absolut Min. temperature of other months (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

RICEURTerrain

*Rainfed Upland Rice terrain requirement for land evaluation*

---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Rainfed Upland Rice.

**Format**

A data frame with 6 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of managemnet with full mechanization. )
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- SlopeD - Slope (degree, 6 classes)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

RICEURWater

*Rainfed Upland Rice water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Rainfed Upland Rice.

**Format**

A data frame with 7 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WmAv1 - Mean precipitation of first month (mm)
- WmAv2 - Mean precipitation of second month (mm)
- WmAv3 - Mean precipitation of third month (mm)
- WmAv4 - Mean precipitation of fourth month (mm)
- WmhAv2 - Relative humidity of devel. Stage (%)
- WmhAv4 - Relative humidity at harvest stage (%)
- WynN - Mean annual n/N

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

RUBBERSoil

*Rubber soil requirement for land evaluation*

---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Rubber.

**Format**

A data frame with 10 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

RUBBERTemp

*Rubber temp requirement for land evaluation*

---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Rubber.

**Format**

A data frame with 3 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TyAv - Mean annual temperature (°C)
- TyMaxAv - Mean annual maximum temperature (°C)
- TdMinXm - Mean daily minimum temperature of coldest month (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

RUBBERTerrain      *Rubber terrain requirement for land evaluation*

---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Rubber.

**Format**

A data frame with 4 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope2 - Slope (%) (2. High level of management with full mechanization. )
- Flood - Flooding
- Drainage - Drainage
- Slope - nan

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

RUBBERWater      *Rubber water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Rubber.

**Format**

A data frame with 3 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)
- WmER - Months of excessive rain (x)
- WmDryLen - Length dry season (months :  $P < 1/2$  PET)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 SAFFLOWERSoil

*Safflower soil requirement for land evaluation*


---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Safflower.

**Format**

A data frame with 8 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 SAFFLOWERTemp

*Safflower temp requirement for land evaluation*


---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Safflower.

**Format**

A data frame with 2 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TmAv1 - Mean temp. of the initial stage( C )
- TmAv2 - Mean temp. crop development stage (2nd month) (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

SAFFLOWERTerrain

*Safflower terrain requirement for land evaluation*

---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Safflower.

**Format**

A data frame with 6 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization. )
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

SAFFLOWERWater

*Safflower water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Safflower.

**Format**

A data frame with 6 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WmAv1 - Mean precipitation of first month (mm)
- WmAv2 - Mean precipitation of second month (mm)
- WmAv3 - Mean precipitation of third month (mm)
- WmAv4 - Mean precipitation of fourth month (mm)
- WmhAv3 - Relative humidity of maturation Stage (%)
- WmhAv4 - Relative humidity at harvest stage (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

SESAMESoil

*Sesame soil requirement for land evaluation*

---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Sesame.

**Format**

A data frame with 7 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 SESAMETemp

*Sesame temp requirement for land evaluation*


---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Sesame.

**Format**

A data frame with 3 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TgMaxAv - Mean max temp. of growing cycle (°C)
- TgMinAv - Mean min. temp. of growing cycle (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 SESAMETerrain

*Sesame terrain requirement for land evaluation*


---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Sesame.

**Format**

A data frame with 6 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization. )
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 SESAMEWater

*Sesame water requirement for land evaluation*


---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Sesame.

**Format**

A data frame with 4 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WmAv1 - Mean precipitation of first month (mm)
- WmAv2 - Mean precipitation of second month (mm)
- WmAv3 - Mean precipitation of third month (mm)
- WghAv - Relative humidity growing cycle (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 SORGHUMSoil

*Sorghum soil requirement for land evaluation*


---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Sorghum.

**Format**

A data frame with 12 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragment - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC6 - Organic carbon (%) - Kaolinitic materials
- OC7 - Organic carbon (%) - Non Kaolinitic, Non calcareous materials
- ECedS - ECe (dS/m)
- ESP - ESP (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

SORGHUMTemp

*Sorghum temp requirement for land evaluation*

---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Sorghum.

**Format**

A data frame with 3 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TgMaxAv - Mean max temp. of growing cycle (°C)
- TgMinAv - Mean min. temp. of growing cycle (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

SORGHUMTerrain      *Sorghum terrain requirement for land evaluation*

---

### Description

A dataset containing the terrain characteristics of the crop requirements for farming Sorghum.

### Format

A data frame with 6 rows and 8 columns

### Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization. )
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)

### See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

SORGHUMWater      *Sorghum water requirement for land evaluation*

---

### Description

A dataset containing the water characteristics of the crop requirements for farming Sorghum.

### Format

A data frame with 3 rows and 8 columns

### Details

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WmAv4 - Mean precipitation of fourth month (mm)
- WghAv - Relative humidity growing cycle (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

SOYASoil

*Soya soil requirement for land evaluation*

---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Soya.

**Format**

A data frame with 11 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragment - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 SOYATemp

*Soya temp requirement for land evaluation*


---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Soya.

**Format**

A data frame with 2 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TgMinAv - Mean min. temp. of growing cycle (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 SOYATerrain

*Soya terrain requirement for land evaluation*


---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Soya.

**Format**

A data frame with 7 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of managemnet with full mechanization. )
- Slope3 - Slope (%) (3. Low level of managemnet animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- SlopeD - Slope (degree, 6 classes)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

SOYAWater

*Soya water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Soya.

**Format**

A data frame with 9 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WmAv1 - Mean precipitation of first month (mm)
- WmAv2 - Mean precipitation of second month (mm)
- WmAv3 - Mean precipitation of third month (mm)
- WmAv4 - Mean precipitation of fourth month (mm)
- WmhAv2 - Relative humidity of devel. Stage (%)
- WmhAv3 - Relative humidity of maturation Stage (%)
- WmnN2 - n/N develop. Stage (2nd month)
- WmnN4 - n/N maturation stage (4th month)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

SUGARCANESoil

*Sugar Cane soil requirement for land evaluation*

---

### **Description**

A dataset containing the soil characteristics of the crop requirements for farming Sugar Cane.

### **Format**

A data frame with 15 rows and 8 columns

### **Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC6 - Organic carbon (%) - Kaolinitic materials
- OC7 - Organic carbon (%) - Non Kaolinitic, Non calcareous materials
- OC8 - Organic carbon (%) - Calcareous materials
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- OC - Organic carbon (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

### **See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 SUGARCANETemp

*Sugar Cane temp requirement for land evaluation*


---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Sugar Cane.

**Format**

A data frame with 4 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TdAvg0 - Mean day temperature at germination stage(°C)
- TdAvg1 - Mean day temperature for tillage stage (°C)
- TdAvg2 - Mean day temperature for vegetative stage (°C)
- Tcoef -  $(T_{max}-T_{min})/T_{mean}$  maturation stage

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 SUGARCANETerrain

*Sugar Cane terrain requirement for land evaluation*


---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Sugar Cane.

**Format**

A data frame with 7 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of managemnet with full mechanization. )
- Slope3 - Slope (%) (3. Low level of managemnet animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- SlopeD - Slope (degree, 6 classes)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 SUGARCANEWater

*Sugar Cane water requirement for land evaluation*


---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Sugar Cane.

**Format**

A data frame with 4 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Wd10 - 10 days of rainfall (mm)
- SunH - Sunshine : hours/year
- WynN - Mean annual n/N
- WmhAv3 - Relative humidity of maturation Stage (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 suit

*Suitability Scores/Class of the Land Units*


---

**Description**

This function calculates the suitability scores and class of the land units.

**Usage**

```
suit(
  crop,
  terrain = NULL,
  water = NULL,
  temp = NULL,
  mf = "triangular",
  sow_month = NULL,
  minimum = NULL,
  maximum = "average",
  interval = NULL,
  sigma = NULL
)
```

**Arguments**

crop	a string for the name of the crop;
terrain	a data frame for the terrain characteristics of the input land units;
water	a data frame for the water characteristics of the input land units;
temp	a data frame for the temperature characteristics of the input land units;
mf	membership function with default assigned to "triangular" fuzzy model. Other fuzzy models included are "trapezoidal" and "gaussian".
sow_month	sowing month of the crop. Takes integers from 1 to 12 (inclusive), representing the twelve months of the year. So if sets to 1, the function assumes sowing month to be January.
minimum	factor's minimum value. If NULL (default), minimum is set to 0. But if numeric of length one, say 0.5, then minimum is set to 0.5, for all factors. To set multiple minimums for multiple factors, simply concatenate these into a numeric vector, the length of this vector should be equal to the number of factors in input land units parameters. However, it can also be set to "average", please refer to the online documentation for more, link in the "See Also" section below.
maximum	maximum value for factors. To set multiple maximums for multiple factors, simply concatenate these into a numeric vector, the length of this vector should be equal to the number of factors in input land units parameters. However, it can also be set to "average", please refer to the online documentation for more, link in the "See Also" section below.
interval	domains for every suitability class (S1, S2, S3). If fixed (NULL), the interval would be 0 to 25% for N (Not Suitable), 25% to 50% for S3 (Marginally Suitable), 50% to 75% for S2 (Moderately Suitable), and 75% to 100% for (Highly Suitable). If "unbias", the package will take into account the shape of the membership function, and provide the appropriate suitability class intervals. However, it can also be customized by specifying the limits of the suitability classes. Please refer to the online documentation for more, link in the "See Also" section below.
sigma	If mf = "gaussian", then sigma represents the constant sigma in the Gaussian formula.

**Value**

A list of outputs of target characteristics, with the following components:

- "terrain" - a list of outputs for terrain characteristics
- "soil" - a list of outputs for soil characteristics
- "water" - a list of outputs for water characteristics
- "temp" - a list of outputs for temperature characteristics

These components are only available when specified as the target characteristics in either of the arguments above, that is, if terrain argument is specified above, then the "terrain" and "soil" components will be available in the output list. This is also true if water and temp are specified in the arguments above.

Each of the components returned above contains a list of outputs as well with the following components:

- "Factors Evaluated" - a character of factors that matched between the input land units factor and the targetted crop requirement factor
- "Suitability Score" - a data frame of suitability scores for each of the matched factors
- "Suitability Class" - a data frame of suitability classes for each of the matched factors
- "Factors' Minimum Values" - a numeric of minimum values used in the membership function for computing the suitability scores
- "Factors' Maximum Values" - a numeric of maximum values used in the membership function for computing the suitability scores
- "Factors' Weights" - a numeric of weights of the factors specified in the input crop requirements
- "Crop Evaluated" - a character of the name of the targetted crop requirement dataset

**See Also**

<https://alstat.github.io/ALUES/>

**Examples**

```
library(ALUES)

rice_suit <- suit("ricebr", water=MarinduqueWater, temp=MarinduqueTemp, sow_month = 1)
lapply(rice_suit[["water"]], function(x) head(x)) # access results for water suitability
lapply(rice_suit[["temp"]], function(x) head(x)) # access results for temperature suitability
rice_suit <- suit("ricebr", terrain=MarinduqueLT)
lapply(rice_suit[["terrain"]], function(x) head(x))
lapply(rice_suit[["soil"]], function(x) head(x))
```

---

 suitability

*Suitability Scores/Class of the Land Units*


---

### Description

This function calculates the suitability scores and class of the land units.

### Usage

```
suitability(
  x,
  y,
  mf = "triangular",
  sow_month = NULL,
  minimum = NULL,
  maximum = "average",
  interval = NULL,
  sigma = NULL
)
```

### Arguments

x	a data frame consisting the properties of the land units;
y	a data frame consisting the requirements of a given characteristics (terrain, soil, water and temperature) for a given crop (e.g. coconut, cassava, etc.);
mf	membership function with default assigned to "triangular" fuzzy model. Other fuzzy models included are "trapezoidal" and "gaussian".
sow_month	sowing month of the crop. Takes integers from 1 to 12 (inclusive), representing the twelve months of the year. So if sets to 1, the function assumes sowing month to be January.
minimum	factor's minimum value. If NULL (default), minimum is set to 0. But if numeric of length one, say 0.5, then minimum is set to 0.5, for all factors. To set multiple minimums for multiple factors, simply concatenate these into a numeric vector, the length of this vector should be equal to the number of factors in input land units parameters. However, it can also be set to "average", please refer to the online documentation for more, link in the "See Also" section below.
maximum	maximum value for factors. To set multiple maximums for multiple factors, simply concatenate these into a numeric vector, the length of this vector should be equal to the number of factors in input land units parameters. However, it can also be set to "average", please refer to the online documentation for more, link in the "See Also" section below.
interval	domains for every suitability class (S1, S2, S3). If fixed (NULL), the interval would be 0 to 25% for N (Not Suitable), 25% to 50% for S3 (Marginally Suitable), 50% to 75% for S2 (Moderately Suitable), and 75% to 100% for (Highly

Suitable). If "unbias", the package will take into account the shape of the membership function, and provide the appropriate suitability class intervals. However, it can also be customized by specifying the limits of the suitability classes. Please refer to the online documentation for more, link in the "See Also" section below.

sigma If mf = "gaussian", then sigma represents the constant sigma in the Gaussian formula.

### Value

A list with the following components:

- "Factors Evaluated" - a character of factors that matched between the input land units factor and the targetted crop requirement factor
- "Suitability Score" - a data frame of suitability scores for each of the matched factors
- "Suitability Class" - a data frame of suitability classes for each of the matched factors
- "Factors' Minimum Values" - a numeric of minimum values used in the membership function for computing the suitability scores
- "Factors' Maximum Values" - a numeric of maximum values used in the membership function for computing the suitability scores
- "Factors' Weights" - a numeric of weights of the factors specified in the input crop requirements
- "Crop Evaluated" - a character of the name of the targetted crop requirement dataset

#' @seealso <https://alstat.github.io/ALUES/>

---

SUNFLOWERSoil

*Sunflower soil requirement for land evaluation*

---

### Description

A dataset containing the soil characteristics of the crop requirements for farming Sunflower.

### Format

A data frame with 11 rows and 8 columns

### Details

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)

- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pH<sub>H2O</sub> - pH H<sub>2</sub>O
- OC - Organic carbon (%)
- EC<sub>dS</sub> - EC<sub>e</sub> (dS/m)
- ESP - ESP (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

SUNFLOWERTemp

*Sunflower temp requirement for land evaluation*

---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Sunflower.

**Format**

A data frame with 1 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

SUNFLOWERTerrain

*Sunflower terrain requirement for land evaluation*

---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Sunflower.

**Format**

A data frame with 6 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization. )
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

SUNFLOWERWater

*Sunflower water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Sunflower.

**Format**

A data frame with 6 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WmAv1 - Mean precipitation of first month (mm)
- WmAv2 - Mean precipitation of second month (mm)
- WmAv4 - Mean precipitation of fourth month (mm)
- WmAv5 - Mean precipitation of fifth month (mm)
- WghAv - Relative humidity growing cycle (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 TEASoil

*Tea soil requirement for land evaluation*


---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Tea.

**Format**

A data frame with 12 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 TEATemp

*Tea temp requirement for land evaluation*


---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Tea.

**Format**

A data frame with 5 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TyAv - Mean annual temperature (°C)
- TmMinAv - Mean min. temp. of warmest month (°C)
- TmMinXm - Avarage minimum temperature of coldest month ( C )
- TmMaxXm - Average max. temp. warmest month (°C)
- TmAv4Xm - Mean temp. of 4 warmest month (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 TEATerrain

*Tea terrain requirement for land evaluation*


---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Tea.

**Format**

A data frame with 4 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope2 - Slope (%) (2. High level of management with full mechanization. )
- Flood - Flooding
- Drainage - Drainage
- SlopeD - Slope (degree, 6 classes)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

TEAWater

*Tea water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Tea.

**Format**

A data frame with 4 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WyAv - Annual precipitation (mm)
- WmDryLen - Length dry season (months :  $P < 1/2$  PET)
- WmDryLen - Length dry season (months :  $P < 1/2$  PET)
- WyhAv - Mean annual rel. humidity (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

TOBACCOSoil

*Tobacco soil requirement for land evaluation*

---

### **Description**

A dataset containing the soil characteristics of the crop requirements for farming Tobacco.

### **Format**

A data frame with 12 rows and 8 columns

### **Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

### **See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 TOBACCOTemp

*Tobacco temp requirement for land evaluation*


---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Tobacco.

**Format**

A data frame with 1 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 TOBACCOTerrain

*Tobacco terrain requirement for land evaluation*


---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Tobacco.

**Format**

A data frame with 7 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of managemnet with full mechanization. )
- Slope3 - Slope (%) (3. Low level of managemnet animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- SlopeD - Slope (degree, 6 classes)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 TOBACCOWater

*Tobacco water requirement for land evaluation*


---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Tobacco.

**Format**

A data frame with 3 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WymN - Mean annual n/N
- WghAv - Relative humidity growing cycle (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

 TOMATOSoil

*Tomato soil requirement for land evaluation*


---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Tomato.

**Format**

A data frame with 12 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyps - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- SoilTe - 12 classes of soil texture (Soil Taxonomy)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

TOMATOTemp

*Tomato temp requirement for land evaluation*


---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Tomato.

**Format**

A data frame with 4 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TmAv0 - Mean temp. at germination (°C) (1st month)
- TmAv3 - Mean temp. of the flowering stage (°C)
- TdDiff3 - Temp. diff day/night flowering stage (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

TOMATOTerrain                      *Tomato terrain requirement for land evaluation*

---

### **Description**

A dataset containing the terrain characteristics of the crop requirements for farming Tomato.

### **Format**

A data frame with 7 rows and 8 columns

### **Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization. )
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- SlopeD - Slope (degree, 6 classes)

### **See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

TOMATOWater                      *Tomato water requirement for land evaluation*

---

### **Description**

A dataset containing the water characteristics of the crop requirements for farming Tomato.

### **Format**

A data frame with 2 rows and 8 columns

### **Details**

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WghAv - Relative humidity growing cycle (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

WATERMELONSoil

*Watermelon soil requirement for land evaluation*

---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Watermelon.

**Format**

A data frame with 10 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragm - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- ECedS - ECe (dS/m)
- ESP - ESP (%)
- SoilTe - nan

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

WATERMELONTemp      *Watermelon temp requirement for land evaluation*

---

### Description

A dataset containing the temp characteristics of the crop requirements for farming Watermelon.

### Format

A data frame with 1 rows and 8 columns

### Details

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)

### See Also

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

WATERMELONTerrain      *Watermelon terrain requirement for land evaluation*

---

### Description

A dataset containing the terrain characteristics of the crop requirements for farming Watermelon.

### Format

A data frame with 7 rows and 8 columns

### Details

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of managemnet with full mechanization. )
- Slope3 - Slope (%) (3. Low level of managemnet animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)
- SlopeD - Slope (degree, 6 classes)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

WATERMELONWater

*Watermelon water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Watermelon.

**Format**

A data frame with 2 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WghAv - Relative humidity growing cycle (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

WHEATSoil

*Wheat soil requirement for land evaluation*

---

**Description**

A dataset containing the soil characteristics of the crop requirements for farming Wheat.

**Format**

A data frame with 14 rows and 8 columns

**Details**

The following are the factors for evaluation:

- CFragn - Coarse fragment (Vol.%)
- SoilDpt - Soil depth (cm)
- CaCO3 - CaCO3 (%)
- Gyph - Gypsum (%)
- CECc - Apparent CEC Clay (cmol (+)/kg clay)
- BS - Base Saturation (%)
- SumBCs - Sum of basic cations (cmol (+)/kg soil)
- pHH2O - pH H2O
- OC - Organic carbon (%)
- OC6 - Organic carbon (%) - Kaolinitic materials
- OC7 - Organic carbon (%) - Non Kaolinitic, Non calcareous materials
- OC8 - Organic carbon (%) - Calcareous materials
- ECedS - ECe (dS/m)
- ESP - ESP (%)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

WHEATTemp

*Wheat temp requirement for land evaluation*

---

**Description**

A dataset containing the temp characteristics of the crop requirements for farming Wheat.

**Format**

A data frame with 4 rows and 8 columns

**Details**

The following are the factors for evaluation:

- TgAv - Mean temperature of the growing cycle (°C)
- TmAv2 - Mean temp. crop development stage (2nd month) (°C)
- TmAv3 - Mean temp. of the flowering stage (°C)
- TmAv4 - Mean temp. of the ripening stage (°C)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

WHEATTerrain

*Wheat terrain requirement for land evaluation*

---

**Description**

A dataset containing the terrain characteristics of the crop requirements for farming Wheat.

**Format**

A data frame with 6 rows and 8 columns

**Details**

The following are the factors for evaluation:

- Slope1 - Slope (%) (1. Irrigated agriculture, basin furrow irrigation)
- Slope2 - Slope (%) (2. High level of management with full mechanization. )
- Slope3 - Slope (%) (3. Low level of management animal traction or handwork.)
- Flood - Flooding
- Drainage4 - Drainage (Medium and fine textured soils)
- Drainage5 - Drainage (Coarse textured soils - Sandy families)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

---

WHEATWater

*Wheat water requirement for land evaluation*

---

**Description**

A dataset containing the water characteristics of the crop requirements for farming Wheat.

**Format**

A data frame with 4 rows and 8 columns

**Details**

The following are the factors for evaluation:

- WgAv - Precipitation of growing cycle (mm)
- WmAv2 - Mean precipitation of second month (mm)
- WmAv3 - Mean precipitation of third month (mm)
- WmAv4 - Mean precipitation of fourth month (mm)

**See Also**

- Yen, B. T., Pheng, K. S., and Hoanh, C. T. (2006). *LUSET: Land Use Suitability Evaluation Tool User's Guide*. International Rice Research Institute.

# Index

## \* dataset

ALFALFASoil, 7  
ALFALFATemp, 8  
ALFALFATerrain, 9  
ALFALFAWater, 9  
AVOCADOSoil, 10  
AVOCADOTemp, 11  
AVOCADOTerrain, 11  
AVOCADOWater, 12  
BAMBOOSoil, 12  
BAMBOOTemp, 13  
BAMBOOTerrain, 13  
BAMBOOWater, 14  
BANANASoil, 14  
BANANATemp, 15  
BANANATerrain, 16  
BANANAWater, 16  
BARLEYSoil, 17  
BARLEYTemp, 18  
BARLEYTerrain, 18  
BARLEYWater, 19  
BEANCAsoil, 19  
BEANCATemp, 20  
BEANCATerrain, 21  
BEANCAWater, 21  
BEANSSoil, 22  
BEANSTemp, 23  
BEANSTerrain, 23  
BEANSWater, 24  
CABBAGESoil, 24  
CABBAGETemp, 25  
CABBAGETerrain, 26  
CABBAGEWater, 26  
CARROTSSoil, 27  
CARROTSTemp, 28  
CARROTSTerrain, 28  
CARROTSWater, 29  
CASHEWSoil, 29  
CASHEWTemp, 30  
CASHEWTerrain, 31  
CASHEWWater, 31  
CASSAVASoil, 32  
CASSAVATemp, 33  
CASSAVATerrain, 33  
CASSAVAWater, 34  
CHICKPEASoil, 34  
CHICKPEATemp, 35  
CHICKPEATerrain, 36  
CHICKPEAWater, 36  
CINNAMONSoil, 37  
CINNAMONTemp, 38  
CINNAMONTerrain, 38  
CINNAMONWater, 39  
CITRUSSoil, 39  
CITRUSTemp, 40  
CITRUSTerrain, 41  
CITRUSWater, 41  
COCOASoil, 42  
COCOATemp, 43  
COCOATerrain, 43  
COCOAWater, 44  
COCONUTSoil, 44  
COCONUTTemp, 45  
COCONUTTerrain, 46  
COCONUTWater, 46  
COFFEEARSoil, 47  
COFFEEARTemp, 48  
COFFEEARTerrain, 48  
COFFEEARWater, 49  
COFFEEEROSoil, 49  
COFFEEEROTemp, 50  
COFFEEEROTerrain, 51  
COFFEEEROWater, 51  
COTTONSoil, 52  
COTTONTemp, 53  
COTTONTerrain, 53  
COTTONWater, 54  
COWPEASoil, 55

- COWPEATemp, [55](#)
- COWPEATerrain, [56](#)
- COWPEAWater, [57](#)
- CUCUMBERSoil, [57](#)
- CUCUMBERTemp, [58](#)
- CUCUMBERTerrain, [59](#)
- CUCUMBERWater, [59](#)
- GROUNDNUTSoil, [60](#)
- GROUNDNUTSTemp, [61](#)
- GROUNDNUTSTerrain, [61](#)
- GROUNDNUTSWater, [62](#)
- GUAVASoil, [62](#)
- GUAVATemp, [63](#)
- GUAVATerrain, [64](#)
- GUAVAWater, [64](#)
- LaoCaiLT, [65](#)
- LaoCaiTemp, [65](#)
- LaoCaiWater, [66](#)
- LONGANSoil, [67](#)
- LONGANTemp, [68](#)
- LONGANTerrain, [68](#)
- LONGANWater, [69](#)
- MAIZESoil, [69](#)
- MAIZETemp, [70](#)
- MAIZETerrain, [71](#)
- MAIZEWater, [71](#)
- MANGOSoil, [72](#)
- MANGOTemp, [73](#)
- MANGOTerrain, [73](#)
- MANGOWater, [74](#)
- MarinduqueLT, [74](#)
- MarinduqueTemp, [75](#)
- MarinduqueWater, [76](#)
- MILLETSoil, [76](#)
- MILLETSTemp, [77](#)
- MILLETSTerrain, [78](#)
- MILLETSWater, [78](#)
- OILPALMSoil, [79](#)
- OILPALMTemp, [80](#)
- OILPALMTerrain, [80](#)
- OILPALMWater, [81](#)
- OLIVESSoil, [81](#)
- OLIVESTemp, [82](#)
- OLIVESTerrain, [83](#)
- OLIVESWater, [83](#)
- ONIONSoil, [84](#)
- ONIONTemp, [85](#)
- ONIONTerrain, [85](#)
- ONIONWater, [86](#)
- PAPAYASoil, [88](#)
- PAPAYATemp, [88](#)
- PAPAYATerrain, [89](#)
- PAPAYAWater, [89](#)
- PEACHSoil, [90](#)
- PEACHTemp, [91](#)
- PEACHTerrain, [91](#)
- PEACHWater, [92](#)
- PEARSoil, [92](#)
- PEARTemp, [93](#)
- PEARTerrain, [93](#)
- PEARWater, [94](#)
- PEASoil, [95](#)
- PEATemp, [95](#)
- PEATerrain, [96](#)
- PEAWater, [97](#)
- PEPPERGRSoil, [97](#)
- PEPPERGRTemp, [98](#)
- PEPPERGRTerrain, [98](#)
- PEPPERGRWater, [99](#)
- PERSIMMONSoil, [100](#)
- PERSIMMONTemp, [101](#)
- PERSIMMONTerrain, [101](#)
- PERSIMMONWater, [102](#)
- PINEAPPLESoil, [102](#)
- PINEAPPLETemp, [103](#)
- PINEAPPLETerrain, [103](#)
- PINEAPPLEWater, [104](#)
- PLUMSoil, [105](#)
- PLUMTemp, [106](#)
- PLUMTerrain, [106](#)
- PLUMWater, [107](#)
- POTATOSoil, [107](#)
- POTATOSWSoil, [108](#)
- POTATOSWTemp, [109](#)
- POTATOSWTerrain, [109](#)
- POTATOSWWater, [110](#)
- POTATOTemp, [110](#)
- POTATOTerrain, [111](#)
- POTATOWater, [112](#)
- RICEBRSoil, [112](#)
- RICEBRTemp, [113](#)
- RICEBRTerrain, [114](#)
- RICEBRWater, [114](#)
- RICEIWSoil, [115](#)
- RICEIWTemp, [116](#)
- RICEIWTerrain, [116](#)

- RICEIWater, 117
- RICENSoil, 118
- RICENTemp, 119
- RICENTerrain, 119
- RICENWater, 120
- RICEURSoil, 121
- RICEURTemp, 121
- RICEURTerrain, 122
- RICEURWater, 123
- RUBBERSoil, 123
- RUBBERTemp, 124
- RUBBERTerrain, 125
- RUBBERWater, 125
- SAFFLOWERSoil, 126
- SAFFLOWERTemp, 126
- SAFFLOWERTerrain, 127
- SAFFLOWERWater, 127
- SESAMESoil, 128
- SESAMETemp, 129
- SESAMETerrain, 129
- SESAMEWater, 130
- SORGHUMSoil, 130
- SORGHUMTemp, 131
- SORGHUMTerrain, 132
- SORGHUMWater, 132
- SOYASoil, 133
- SOYATemp, 134
- SOYATerrain, 134
- SOYAWater, 135
- SUGARCANESoil, 136
- SUGARCANETemp, 137
- SUGARCANETerrain, 137
- SUGARCANEWater, 138
- SUNFLOWERSoil, 142
- SUNFLOWERTemp, 143
- SUNFLOWERTerrain, 144
- SUNFLOWERWater, 144
- TEASoil, 145
- TEATemp, 146
- TEATerrain, 146
- TEAWater, 147
- TOBACCSoil, 148
- TOBACCOTemp, 149
- TOBACCOTerrain, 149
- TOBACOWater, 150
- TOMATOSoil, 150
- TOMATOTemp, 151
- TOMATOTerrain, 152
- TOMATOWater, 152
- WATERMELONSoil, 153
- WATERMELONTemp, 154
- WATERMELONTerrain, 154
- WATERMELONWater, 155
- WHEATSoil, 155
- WHEATTemp, 156
- WHEATTerrain, 157
- WHEATWater, 157
  
- ALFALFASoil, 7
- ALFALFATemp, 8
- ALFALFATerrain, 9
- ALFALFAWater, 9
- ALUES-package, 7
- AVOCADOSoil, 10
- AVOCADOTemp, 11
- AVOCADOTerrain, 11
- AVOCADOWater, 12
  
- BAMBOOSoil, 12
- BAMBOOTemp, 13
- BAMBOOTerrain, 13
- BAMBOOWater, 14
- BANANASoil, 14
- BANANATemp, 15
- BANANATerrain, 16
- BANANAWater, 16
- BARLEYSoil, 17
- BARLEYTemp, 18
- BARLEYTerrain, 18
- BARLEYWater, 19
- BEANCASoil, 19
- BEANCATemp, 20
- BEANCATerrain, 21
- BEANCAWater, 21
- BEANSSoil, 22
- BEANSTemp, 23
- BEANSTerrain, 23
- BEANSWater, 24
  
- CABBAGESoil, 24
- CABBAGETemp, 25
- CABBAGETerrain, 26
- CABBAGEWater, 26
- CARROTSSoil, 27
- CARROTSTemp, 28
- CARROTSTerrain, 28
- CARROTSWater, 29

- CASHEWSoil, 29  
 CASHEWTemp, 30  
 CASHEWTerrain, 31  
 CASHEWWater, 31  
 CASSAVASoil, 32  
 CASSAVATemp, 33  
 CASSAVATerrain, 33  
 CASSAVAWater, 34  
 CHICKPEASoil, 34  
 CHICKPEATemp, 35  
 CHICKPEATerrain, 36  
 CHICKPEAWater, 36  
 CINNAMONSoil, 37  
 CINNAMONTemp, 38  
 CINNAMONTerrain, 38  
 CINNAMONWater, 39  
 CITRUSSoil, 39  
 CITRUSTemp, 40  
 CITRUSTerrain, 41  
 CITRUSWater, 41  
 COCOASoil, 42  
 COCOATemp, 43  
 COCOATerrain, 43  
 COCOAWater, 44  
 COCONUTSoil, 44  
 COCONUTTemp, 45  
 COCONUTTerrain, 46  
 COCONUTWater, 46  
 COFFEEARSoil, 47  
 COFFEEARTemp, 48  
 COFFEEARTerrain, 48  
 COFFEEARWater, 49  
 COFFEEROSoil, 49  
 COFFEEROTemp, 50  
 COFFEEROTerrain, 51  
 COFFEEROWater, 51  
 COTTONSoil, 52  
 COTTONTemp, 53  
 COTTONTerrain, 53  
 COTTONWater, 54  
 COWPEASoil, 55  
 COWPEATemp, 55  
 COWPEATerrain, 56  
 COWPEAWater, 57  
 CUCUMBERSoil, 57  
 CUCUMBERTemp, 58  
 CUCUMBERTerrain, 59  
 CUCUMBERWater, 59  
 GROUNDNUTSoil, 60  
 GROUNDNUTTemp, 61  
 GROUNDNUTTerrain, 61  
 GROUNDNUTWater, 62  
 GUAVASoil, 62  
 GUAVATemp, 63  
 GUAVATerrain, 64  
 GUAVAWater, 64  
 LaoCaiLT, 65  
 LaoCaiTemp, 65  
 LaoCaiWater, 66  
 LONGANSoil, 67  
 LONGANTemp, 68  
 LONGANTerrain, 68  
 LONGANWater, 69  
 MAIZESoil, 69  
 MAIZETemp, 70  
 MAIZETerrain, 71  
 MAIZEWater, 71  
 MANGOSoil, 72  
 MANGOTemp, 73  
 MANGOTerrain, 73  
 MANGOWater, 74  
 MarinduqueLT, 74  
 MarinduqueTemp, 75  
 MarinduqueWater, 76  
 MILLETSoil, 76  
 MILLETSTemp, 77  
 MILLETSTerrain, 78  
 MILLETWater, 78  
 OILPALMSoil, 79  
 OILPALMTemp, 80  
 OILPALMTerrain, 80  
 OILPALMWater, 81  
 OLIVESSoil, 81  
 OLIVESTemp, 82  
 OLIVESTerrain, 83  
 OLIVESWater, 83  
 ONIONSoil, 84  
 ONIONTemp, 85  
 ONIONTerrain, 85  
 ONIONWater, 86  
 overall\_suit, 86  
 PAPAYASoil, 88  
 PAPAYATemp, 88

- PAPAYATerrain, 89  
PAPAYAWater, 89  
PEACHSoil, 90  
PEACHTemp, 91  
PEACHTerrain, 91  
PEACHWater, 92  
PEARSoil, 92  
PEARTemp, 93  
PEARTerrain, 93  
PEARWater, 94  
PEASoil, 95  
PEATemp, 95  
PEATerrain, 96  
PEAWater, 97  
PEPPERGRSoil, 97  
PEPPERGRTemp, 98  
PEPPERGRTerrain, 98  
PEPPERGRWater, 99  
PERSIMMONSoil, 100  
PERSIMMONTemp, 101  
PERSIMMONTerrain, 101  
PERSIMMONWater, 102  
PINEAPPLESoil, 102  
PINEAPPLETemp, 103  
PINEAPPLETerrain, 103  
PINEAPPLEWater, 104  
PLUMSoil, 105  
PLUMTemp, 106  
PLUMTerrain, 106  
PLUMWater, 107  
POTATOSoil, 107  
POTATOSWSoil, 108  
POTATOSWTemp, 109  
POTATOSWTerrain, 109  
POTATOSWWater, 110  
POTATOTemp, 110  
POTATOTerrain, 111  
POTATOWater, 112
- RICEBRSoil, 112  
RICEBRTemp, 113  
RICEBRTerrain, 114  
RICEBRWater, 114  
RICEIWSoil, 115  
RICEIWTemp, 116  
RICEIWTerrain, 116  
RICEIWWater, 117  
RICENFSoil, 118  
RICENFTemp, 119
- RICENFTerrain, 119  
RICENFWater, 120  
RICEURSoil, 121  
RICEURTemp, 121  
RICEURTerrain, 122  
RICEURWater, 123  
RUBBERSoil, 123  
RUBBERTemp, 124  
RUBBERTerrain, 125  
RUBBERWater, 125
- SAFFLOWERSoil, 126  
SAFFLOWERTemp, 126  
SAFFLOWERTerrain, 127  
SAFFLOWERWater, 127  
SESAMESoil, 128  
SESAMETemp, 129  
SESAMETerrain, 129  
SESAMEWater, 130  
SORGHUMSoil, 130  
SORGHUMTemp, 131  
SORGHUMTerrain, 132  
SORGHUMWater, 132  
SOYASoil, 133  
SOYATemp, 134  
SOYATerrain, 134  
SOYAWater, 135  
SUGARCANESoil, 136  
SUGARCANETemp, 137  
SUGARCANETerrain, 137  
SUGARCANEWater, 138  
suit, 87, 138  
suitability, 141  
SUNFLOWERSoil, 142  
SUNFLOWERTemp, 143  
SUNFLOWERTerrain, 144  
SUNFLOWERWater, 144
- TEASoil, 145  
TEATemp, 146  
TEATerrain, 146  
TEAWater, 147  
TOBACCSOIL, 148  
TOBACCOTemp, 149  
TOBACCOTerrain, 149  
TOBACCOWater, 150  
TOMATOSoil, 150  
TOMATOTemp, 151  
TOMATOTerrain, 152

TOMATOWater, [152](#)

WATERMELONSoil, [153](#)

WATERMELONTemp, [154](#)

WATERMELONTerrain, [154](#)

WATERMELONWater, [155](#)

WHEATSoil, [155](#)

WHEATTemp, [156](#)

WHEATTerrain, [157](#)

WHEATWater, [157](#)