

Package ‘REAT’

May 7, 2026

Type Package

Title Regional Economic Analysis Toolbox

Version 3.0.3

Date 2021-08-24

Author Thomas Wieland

Maintainer Thomas Wieland <thomas.wieland.geo@gmail.com>

Description

Collection of models and analysis methods used in regional and urban economics and (quantitative) economic geography, e.g. measures of inequality, regional disparities and convergence, regional specialization as well as accessibility and spatial interaction models.

License GPL (>= 2)

NeedsCompilation no

Repository CRAN

Date/Publication 2021-09-03 08:00:11 UTC

Contents

REAT-package	3
atkinson	4
Automotive	5
betaconv.nls	6
betaconv.ols	9
betaconv.speed	12
conc	14
converse	15
coulter	16
curvefit	17
cv	19
dalton	21
disp	22
dist.buf	23
dist.calc	25

dist.mat	26
durpug	27
ellison.a	28
ellison.a2	29
ellison.c	30
ellison.c2	31
EU28.emp	33
FK2014_EGC	34
Freiburg	35
G.counties.gdp	36
G.regions.emp	39
G.regions.industries	40
gifpro	41
gifpro.tbs	44
gini	46
gini.conc	50
gini.spec	52
gini2	55
Goettingen	57
GoettingenHealth1	59
GoettingenHealth2	60
growth	61
hansen	62
herf	65
hoover	67
howard.cl	68
howard.xcl	69
howard.xcl2	70
huff	71
krugman.conc	74
krugman.conc2	75
krugman.spec	77
krugman.spec2	78
litzenberger	79
litzenberger2	80
lm.beta	82
locq	83
locq.growth	85
locq2	87
lorenz	89
mean2	92
mssd	93
portfolio	94
rca	96
reilly	101
ripley	103
sd2	105
shift	107

shift.growth	109
shiftd	111
shifti	114
shiftd	116
shiftp	118
sigmaconv	121
sigmaconv.t	124
spec	127
theil	128
to.dummy	129
var2	130
williamson	131

Index	134
--------------	------------

REAT-package

Regional Economic Analysis Toolbox

Description

In regional and urban economics and economic geography, very frequent research fields are the existence and evolution of agglomerations due to (internal and external) agglomeration economies, regional economic growth and regional disparities, where these concepts and relationships are closely related to each other (Capello/Nijkamp 2009, Dinc 2015, Farhauer/Kroell 2013, McCann/van Oort 2009). Also accessibility and spatial interaction modeling is mostly regarded as related to these disciplines (Aoyama et al. 2011, Guessefeldt 1999). The group of the related analysis methods is sometimes summarized by the term *regional analysis* or *regional economic analysis* (Dinc 2015, Guessefeldt 1999, Isard 1960).

This package contains a collection of models and analysis methods used in regional and urban economics and (quantitative) economic geography. The functions in this package can be divided into seven groups:

- (1) Inequality, concentration and dispersion, including *Gini coefficient*, *Lorenz curve*, *Herfindahl-Hirschman-coefficient*, *Theil coefficient*, *Hoover coefficient* and *(weighted) coefficient of variation*
- (2) Specialization of regions and spatial concentration of industries, including *location quotient*, *spatial Gini coefficients* for regional specialization and industry concentration and *Krugman coefficients* for regional specialization and industry concentration
- (3) Regional disparities and regional convergence, especially analysis of *beta and sigma convergence* for cross-sectional data
- (4) Regional growth, including *portfolio matrix*, several types of *shift-share analysis* and *commercial area prognosis* ("GIFPRO")
- (5) Spatial interaction and accessibility models, including *Huff model* and *Hansen accessibility*
- (6) Proximity analysis, including calculation of *distance matrices* and *buffers*
- (7) Additional tools for data preparation und visualization, such as for creating *dummy variables* and calculating *standardized regression coefficients*. The package also contains data examples.

Author(s)

Thomas Wieland

Maintainer: Thomas Wieland <thomas.wieland.geo@googlemail.com>

References

Aoyama, Y./Murphy, J. T./Hanson, S. (2011): “Key Concepts in Economic Geography”. London: SAGE.

Capello, R./Nijkamp, P. (2009): “Introduction: regional growth and development theories in the twenty-first century - recent theoretical advances and future challenges”. In: Capello, R./Nijkamp, P. (eds.): *Handbook of Regional Growth and Development Theories*. Cheltenham: Elgar. p. 1-16.

Dinc, M. (2015): “Introduction to Regional Economic Development. Major Theories and Basic Analytical Tools”. Cheltenham: Elgar.

Farhauer, O./Kroell, A. (2013): “Standorttheorien: Regional- und Stadtoekonomik in Theorie und Praxis”. Wiesbaden: Springer.

Guessefeldt, J. (1999): “Regionalanalyse”. Muenchen: Oldenbourg.

Isard, W. (1960): “Methods of Regional Analysis: an Introduction to Regional Science”. Cambridge: M.I.T. Press.

McCann, P./van Oort, F. (2009): “Theories of agglomeration and regional economic growth: a historical review”. In: Capello, R./Nijkamp, P. (eds.): *Handbook of Regional Growth and Development Theories*. Cheltenham: Elgar. p. 19-32.

atkinson

Atkinson Inequality Index

Description

Calculating the Atkinson Inequality Index e.g. with respect to regional income

Usage

```
atkinson(x, epsilon = 0.5, na.rm = TRUE)
```

Arguments

x	A numeric vector (e.g. dataset of regional income)
epsilon	A single value of the ϵ weighting coefficient (default: <code>at.epsilon = 0.5</code>)
na.rm	logical argument that indicates whether NA values should be excluded before computing results

Details

The *Atkinson Inequality Index (AI)* varies between 0 (no inequality/concentration) and 1 (complete inequality/concentration). It can be used for economic inequality and/or regional disparities (Portnov/Felsenstein 2010).

Value

A single numeric value of the *Atkinson Inequality Index* ($0 < AI < 1$).

Author(s)

Thomas Wieland

References

Portnov, B.A./Felsenstein, D. (2010): "On the suitability of income inequality measures for regional analysis: Some evidence from simulation analysis and bootstrapping tests". In: *Socio-Economic Planning Sciences*, **44**, 4, p. 212-219.

See Also

[cv](#), [gini](#), [gini2](#), [herf](#), [theil](#), [hoover](#), [coulter](#), [dalton](#), [disp](#)

Examples

```
atkinson(c(100,0,0,0), epsilon = 0.8)
```

```
atkinson(c(100,100,100,100), epsilon = 0.8)
```

Automotive

Automotive industry data

Description

Top 20 automotive industry companies, including their manufacturing quantity and turnovers (Table from wikipedia)

Usage

```
data("Automotive")
```

Format

A data frame with 20 observations on the following 8 variables.

Rank Rank of the company

Company Name of the company (German)

Country Origin county of the company (German)

Quantity2014 Quantity of produced vehicles in 2014

Quantity2014_car Quantity of produced cars in 2014

Turnover2008 Annual turnover 2008 (in billion dollars)

Turnover2012 Annual turnover 2012 (in billion dollars)

Turnover2013 Annual turnover 2013 (in billion dollars)

Source

Wikipedia (2018): “Automobilindustrie — Wikipedia, Die freie Enzyklopaedie”. <https://de.wikipedia.org/wiki/Automobilindustrie> (accessed October 14, 2018). Own postprocessing.

References

Wikipedia (2018): “Automobilindustrie — Wikipedia, Die freie Enzyklopaedie”. <https://de.wikipedia.org/wiki/Automobilindustrie> (accessed October 14, 2018).

Examples

```
# Market concentration in automotive industry

data(Automotive)

gini(Automotive$Turnover2008, lsize=1, lc=TRUE, le.col = "black",
lc.col = "orange", lcx = "Shares of companies", lcy = "Shares of turnover / cars",
lctitle = "Automotive industry: market concentration",
lcg = TRUE, lcg.n = TRUE, lcg.caption = "Turnover 2008:", lcg.lab.x = 0, lcg.lab.y = 1)
# Gini coefficient and Lorenz curve for turnover 2008

gini(Automotive$Turnover2013, lsize=1, lc = TRUE, add.lc = TRUE, lc.col = "red",
lcg = TRUE, lcg.n = TRUE, lcg.caption = "Turnover 2013:", lcg.lab.x = 0, lcg.lab.y = 0.85)
# Adding Gini coefficient and Lorenz curve for turnover 2013

gini(Automotive$Quantity2014_car, lsize=1, lc = TRUE, add.lc = TRUE, lc.col = "blue",
lcg = TRUE, lcg.n = TRUE, lcg.caption = "Cars 2014:", lcg.lab.x = 0, lcg.lab.y = 0.7)
# Adding Gini coefficient and Lorenz curve for cars 2014
```

betaconv.nls

Analysis of regional beta convergence using nonlinear regression

Description

This function provides the analysis of absolute and conditional regional economic beta convergence for cross-sectional data using a nonlinear least squares (NLS) technique.

Usage

```
betaconv.nls(gdp1, time1, gdp2, time2, conditions = NULL, conditions.formula = NULL,
conditions.startval = NULL, beta.plot = FALSE, beta.plotPSize = 1,
beta.plotPCol = "black", beta.plotLine = FALSE, beta.plotLineCol = "red",
beta.plotX = "Ln (initial)", beta.plotY = "Ln (growth)",
beta.plotTitle = "Beta convergence", beta.bgCol = "gray95", beta.bgrid = TRUE,
beta.bgridCol = "white", beta.bgridSize = 2, beta.bgridType = "solid",
print.results = TRUE)
```

Arguments

<code>gdp1</code>	A numeric vector containing the GDP per capita (or another economic variable) at time t
<code>time1</code>	A single value of time t (= the initial year)
<code>gdp2</code>	A numeric vector containing the GDP per capita (or another economic variable) at time $t+1$ or a data frame containing the GDPs per capita (or another economic variable) at time $t+1, t+2, t+3, \dots, t+n$
<code>time2</code>	A single value of time $t+1$ or t_n , respectively
<code>conditions</code>	A data frame containing the conditions for conditional beta convergence
<code>conditions.formula</code>	A formula for the functional linkage of the conditions in the case of conditional beta convergence
<code>conditions.startval</code>	Starting values for the parameters of the conditions in the case of conditional beta convergence
<code>beta.plot</code>	Boolean argument that indicates if a plot of beta convergence has to be created
<code>beta.plotPSize</code>	If <code>beta.plot = TRUE</code> : Point size in the beta convergence plot
<code>beta.plotPCol</code>	If <code>beta.plot = TRUE</code> : Point color in the beta convergence plot
<code>beta.plotLine</code>	If <code>beta.plot = TRUE</code> : Logical argument that indicates if a regression line has to be added to the plot
<code>beta.plotLineColor</code>	If <code>beta.plot = TRUE</code> and <code>beta.plotLine = TRUE</code> : Line color of regression line
<code>beta.plotX</code>	If <code>beta.plot = TRUE</code> : Name of the X axis
<code>beta.plotY</code>	If <code>beta.plot = TRUE</code> : Name of the Y axis
<code>beta.plotTitle</code>	If <code>beta.plot = TRUE</code> : Plot title
<code>beta.bgCol</code>	If <code>beta.plot = TRUE</code> : Plot background color
<code>beta.bgrid</code>	If <code>beta.plot = TRUE</code> : Logical argument that indicates if the plot contains a grid
<code>beta.bgridCol</code>	If <code>beta.plot = TRUE</code> and <code>beta.bgrid = TRUE</code> : Color of the grid
<code>beta.bgridSize</code>	If <code>beta.plot = TRUE</code> and <code>beta.bgrid = TRUE</code> : Size of the grid
<code>beta.bgridType</code>	If <code>beta.plot = TRUE</code> and <code>beta.bgrid = TRUE</code> : Type of the grid
<code>print.results</code>	Logical argument that indicates if the function shows the results or not

Details

From the regional economic perspective (in particular the neoclassical growth theory), regional disparities are expected to decline. This *convergence* can have different meanings: *Sigma convergence* (σ) means a harmonization of regional economic output or income over time, while *beta convergence* (β) means a decline of dispersion because poor regions have a stronger economic growth than rich regions (Capello/Nijkamp 2009). Regardless of the theoretical assumptions of a harmonization in reality, the related analytical framework allows to analyze both types of convergence for cross-sectional data (GDP p.c. or another economic variable, y , for i regions and two points in time, t and $t + T$), or one starting point (t) and the average growth within the following n years

($t + 1, t + 2, \dots, t + n$), respectively. Beta convergence can be calculated either in a linearized OLS regression model or in a nonlinear regression model. When no other variables are integrated in this model, it is called *absolute* beta convergence. Implementing other region-related variables (conditions) into the model leads to *conditional* beta convergence. If there is beta convergence ($\beta < 0$), it is possible to calculate the *speed of convergence*, λ , and the so-called *Half-Life* H , while the latter is the time taken to reduce the disparities by one half (Allington/McCombie 2007, Goecke/Huether 2016). There is *sigma convergence*, when the dispersion of the variable (σ), e.g. calculated as standard deviation or coefficient of variation, reduces from t to $t + T$. This can be measured using ANOVA for two years or trend regression with respect to several years (Furceri 2005, Goecke/Huether 2016).

This function calculates absolute and/or conditional beta convergence using a nonlinear least squares approach for estimation. It needs at least two vectors (GDP p.c. or another economic variable, y , for i regions) and the related two points in time (t and $t + T$). If the beta coefficient is negative (using OLS) or positive (using NLS), there is beta convergence.

Value

A list containing the following objects:

regdata	A data frame containing the regression data, including the \ln -transformed economic variables
abeta	A list containing the estimates of the absolute beta convergence regression model, including lambda and half-life
cbeta	If conditions are stated: a list containing the estimates of the conditional beta convergence regression model, including lambda and half-life

Author(s)

Thomas Wieland

References

- Allington, N. F. B./McCombie, J. S. L. (2007): "Economic growth and beta-convergence in the East European Transition Economies". In: Arestis, P./Baddely, M./McCombie, J. S. L. (eds.): *Economic Growth. New Directions in Theory and Policy*. Cheltenham: Elgar. p. 200-222.
- Capello, R./Nijkamp, P. (2009): "Introduction: regional growth and development theories in the twenty-first century - recent theoretical advances and future challenges". In: Capello, R./Nijkamp, P. (eds.): *Handbook of Regional Growth and Development Theories*. Cheltenham: Elgar. p. 1-16.
- Dapena, A. D./Vazquez, E. F./Morollon, F. R. (2016): "The role of spatial scale in regional convergence: the effect of MAUP in the estimation of beta-convergence equations". In: *The Annals of Regional Science*, **56**, 2, p. 473-489.
- Furceri, D. (2005): "Beta and sigma-convergence: A mathematical relation of causality". In: *Economics Letters*, **89**, 2, p. 212-215.
- Goecke, H./Huether, M. (2016): "Regional Convergence in Europe". In: *Intereconomics*, **51**, 3, p. 165-171.
- Young, A. T./Higgins, M. J./Levy, D. (2008): "Sigma Convergence versus Beta Convergence: Evidence from U.S. County-Level Data". In: *Journal of Money, Credit and Banking*, **40**, 5, p. 1083-1093.

See Also

[rca](#), [betaconv.ols](#), [betaconv.speed](#), [sigmaconv](#), [sigmaconv.t](#), [cv](#), [sd2](#), [var2](#)

Examples

```
data (G.counties.gdp)
# Loading GDP data for Germany (counties = Landkreise)
betaconv.nls (G.counties.gdp$gdppc2010, 2010, G.counties.gdp$gdppc2011, 2011,
conditions = NULL, print.results = TRUE)
# Two years, no conditions (Absolute beta convergence)
```

betaconv.ols

Analysis of regional beta convergence using OLS regression

Description

This function provides the analysis of absolute and conditional regional economic beta convergence for cross-sectional data using ordinary least squares (OLS) technique.

Usage

```
betaconv.ols(gdp1, time1, gdp2, time2, conditions = NULL, beta.plot = FALSE,
beta.plotPSize = 1, beta.plotPCol = "black", beta.plotLine = FALSE,
beta.plotLineCol = "red", beta.plotX = "Ln (initial)", beta.plotY = "Ln (growth)",
beta.plotTitle = "Beta convergence", beta.bgCol = "gray95", beta.bgrid = TRUE,
beta.bgridCol = "white", beta.bgridSize = 2, beta.bgridType = "solid",
print.results = FALSE)
```

Arguments

gdp1	A numeric vector containing the GDP per capita (or another economic variable) at time t
time1	A single value of time t (= the initial year)
gdp2	A numeric vector containing the GDP per capita (or another economic variable) at time $t+1$ or a data frame containing the GDPs per capita (or another economic variable) at time $t+1, t+2, t+3, \dots, t+n$
time2	A single value of time $t+1$ or t_n , respectively
conditions	A data frame containing the conditions for conditional beta convergence
beta.plot	Boolean argument that indicates if a plot of beta convergence has to be created
beta.plotPSize	If beta.plot = TRUE: Point size in the beta convergence plot
beta.plotPCol	If beta.plot = TRUE: Point color in the beta convergence plot
beta.plotLine	If beta.plot = TRUE: Logical argument that indicates if a regression line has to be added to the plot
beta.plotLineCol	If beta.plot = TRUE and beta.plotLine = TRUE: Line color of regression line

beta.plotX	If beta.plot = TRUE: Name of the X axis
beta.plotY	If beta.plot = TRUE: Name of the Y axis
beta.plotTitle	If beta.plot = TRUE: Plot title
beta.bgCol	If beta.plot = TRUE: Plot background color
beta.bgrid	If beta.plot = TRUE: Logical argument that indicates if the plot contains a grid
beta.bgridCol	If beta.plot = TRUE and beta.bgrid = TRUE: Color of the grid
beta.bgridSize	If beta.plot = TRUE and beta.bgrid = TRUE: Size of the grid
beta.bgridType	If beta.plot = TRUE and beta.bgrid = TRUE: Type of the grid
print.results	Logical argument that indicates if the function shows the results or not

Details

From the regional economic perspective (in particular the neoclassical growth theory), regional disparities are expected to decline. This *convergence* can have different meanings: *Sigma convergence* (σ) means a harmonization of regional economic output or income over time, while *beta convergence* (β) means a decline of dispersion because poor regions have a stronger economic growth than rich regions (Capello/Nijkamp 2009). Regardless of the theoretical assumptions of a harmonization in reality, the related analytical framework allows to analyze both types of convergence for cross-sectional data (GDP p.c. or another economic variable, y , for i regions and two points in time, t and $t + T$), or one starting point (t) and the average growth within the following n years ($t + 1, t + 2, \dots, t + n$), respectively. Beta convergence can be calculated either in a linearized OLS regression model or in a nonlinear regression model. When no other variables are integrated in this model, it is called *absolute* beta convergence. Implementing other region-related variables (conditions) into the model leads to *conditional* beta convergence. If there is beta convergence ($\beta < 0$), it is possible to calculate the *speed of convergence*, λ , and the so-called *Half-Life* H , while the latter is the time taken to reduce the disparities by one half (Allington/McCombie 2007, Goecke/Huether 2016). There is *sigma convergence*, when the dispersion of the variable (σ), e.g. calculated as standard deviation or coefficient of variation, reduces from t to $t + T$. This can be measured using ANOVA for two years or trend regression with respect to several years (Furceri 2005, Goecke/Huether 2016).

This function calculates absolute and/or conditional beta convergence using ordinary least squares regression (OLS) for estimation. It needs at least two vectors (GDP p.c. or another economic variable, y , for i regions) and the related two points in time (t and $t + T$). If the beta coefficient is negative (using OLS) or positive (using NLS), there is beta convergence.

Value

A list containing the following objects:

regdata	A data frame containing the regression data, including the \ln -transformed economic variables
abeta	A list containing the estimates of the absolute beta convergence regression model, including lambda and half-life
cbeta	If conditions are stated: a list containing the estimates of the conditional beta convergence regression model, including lambda and half-life

Author(s)

Thomas Wieland

References

- Allington, N. F. B./McCombie, J. S. L. (2007): “Economic growth and beta-convergence in the East European Transition Economies”. In: Arestis, P./Baddely, M./McCombie, J. S. L. (eds.): *Economic Growth. New Directions in Theory and Policy*. Cheltenham: Elgar. p. 200-222.
- Capello, R./Nijkamp, P. (2009): “Introduction: regional growth and development theories in the twenty-first century - recent theoretical advances and future challenges”. In: Capello, R./Nijkamp, P. (eds.): *Handbook of Regional Growth and Development Theories*. Cheltenham: Elgar. p. 1-16.
- Dapena, A. D./Vazquez, E. F./Morollon, F. R. (2016): “The role of spatial scale in regional convergence: the effect of MAUP in the estimation of beta-convergence equations”. In: *The Annals of Regional Science*, **56**, 2, p. 473-489.
- Furceri, D. (2005): “Beta and sigma-convergence: A mathematical relation of causality”. In: *Economics Letters*, **89**, 2, p. 212-215.
- Goecke, H./Huether, M. (2016): “Regional Convergence in Europe”. In: *Intereconomics*, **51**, 3, p. 165-171.
- Young, A. T./Higgins, M. J./Levy, D. (2008): “Sigma Convergence versus Beta Convergence: Evidence from U.S. County-Level Data”. In: *Journal of Money, Credit and Banking*, **40**, 5, p. 1083-1093.

See Also

[rca](#), [betaconv.nls](#), [betaconv.speed](#), [sigmaconv](#), [sigmaconv.t](#), [cv](#), [sd2](#), [var2](#)

Examples

```
data (G.counties.gdp)

betaconv.ols (G.counties.gdp$gdppc2010, 2010, G.counties.gdp$gdppc2011, 2011,
conditions = NULL, print.results = TRUE)
# Two years, no conditions (Absolute beta convergence)

regionaldummies <- to.dummy(G.counties.gdp$regional)
# Creating dummy variables for West/East
G.counties.gdp$West <- regionaldummies[,2]
G.counties.gdp$East <- regionaldummies[,1]
# Adding dummy variables to data

betaconv.ols (G.counties.gdp$gdppc2010, 2010, G.counties.gdp$gdppc2011, 2011,
conditions = G.counties.gdp[c(70,71)], print.results = TRUE)
# Two years, with condition (dummy for West/East)
# (Absolute and conditional beta convergence)

betaconverg1 <- betaconv.ols (G.counties.gdp$gdppc2010, 2010, G.counties.gdp$gdppc2011, 2011,
conditions = G.counties.gdp[c(70,71)], print.results = TRUE)
# Store results in object
betaconverg1$cbeta$estimates
```

```
# Addressing estimates for the conditional beta model

betaconv.ols (G.counties.gdp$gdppc2010, 2010, G.counties.gdp[65:66], 2012,
conditions = NULL, print.results = TRUE)
# Three years (2010-2012), no conditions (Absolute beta convergence)

betaconv.ols (G.counties.gdp$gdppc2010, 2010, G.counties.gdp[65:66], 2012,
conditions = G.counties.gdp[c(70,71)], print.results = TRUE)
# Three years (2010-2012), with conditions (Absolute and conditional beta convergence)

betaconverg2 <- betaconv.ols (G.counties.gdp$gdppc2010, 2010, G.counties.gdp[65:66],
2012, conditions = G.counties.gdp[c(70,71)], print.results = TRUE)
# Store results in object
betaconverg2$beta$estimates
# Addressing estimates for the conditional beta model
```

betaconv.speed	<i>Regional beta convergence: Convergence speed and half-life</i>
----------------	---

Description

This function calculates the beta convergence speed and half-life based on a given beta value and time interval.

Usage

```
betaconv.speed(beta, tinterval, print.results = TRUE)
```

Arguments

beta	Beta value
tinterval	Time interval (in time units, such as years)
print.results	Logical argument that indicates if the function shows the results or not

Details

From the regional economic perspective (in particular the neoclassical growth theory), regional disparities are expected to decline. This *convergence* can have different meanings: *Sigma convergence* (σ) means a harmonization of regional economic output or income over time, while *beta convergence* (β) means a decline of dispersion because poor regions have a stronger economic growth than rich regions (Capello/Nijkamp 2009). Regardless of the theoretical assumptions of a harmonization in reality, the related analytical framework allows to analyze both types of convergence for cross-sectional data (GDP p.c. or another economic variable, y , for i regions and two points in time, t and $t + T$), or one starting point (t) and the average growth within the following n years ($t + 1, t + 2, \dots, t + n$), respectively. Beta convergence can be calculated either in a linearized OLS regression model or in a nonlinear regression model. When no other variables are integrated in this model, it is called *absolute* beta convergence. Implementing other region-related variables

(conditions) into the model leads to *conditional* beta convergence. If there is beta convergence ($\beta < 0$), it is possible to calculate the *speed of convergence*, λ , and the so-called *Half-Life* H , while the latter is the time taken to reduce the disparities by one half (Allington/McCombie 2007, Goecke/Huether 2016). There is *sigma convergence*, when the dispersion of the variable (σ), e.g. calculated as standard deviation or coefficient of variation, reduces from t to $t + T$. This can be measured using ANOVA for two years or trend regression with respect to several years (Furceri 2005, Goecke/Huether 2016).

This function calculates the *speed of convergence*, λ , and the *Half-Life*, H , based on a given β value and time interval.

Value

A matrix containing the following objects:

Lambda	Lambda value (convergence speed)
Half-Life	Half-life values

Author(s)

Thomas Wieland

References

- Allington, N. F. B./McCombie, J. S. L. (2007): "Economic growth and beta-convergence in the East European Transition Economies". In: Arestis, P./Baddely, M./McCombie, J. S. L. (eds.): *Economic Growth. New Directions in Theory and Policy*. Cheltenham: Elgar. p. 200-222.
- Capello, R./Nijkamp, P. (2009): "Introduction: regional growth and development theories in the twenty-first century - recent theoretical advances and future challenges". In: Capello, R./Nijkamp, P. (eds.): *Handbook of Regional Growth and Development Theories*. Cheltenham: Elgar. p. 1-16.
- Dapena, A. D./Vazquez, E. F./Morollon, F. R. (2016): "The role of spatial scale in regional convergence: the effect of MAUP in the estimation of beta-convergence equations". In: *The Annals of Regional Science*, **56**, 2, p. 473-489.
- Furceri, D. (2005): "Beta and sigma-convergence: A mathematical relation of causality". In: *Economics Letters*, **89**, 2, p. 212-215.
- Goecke, H./Huether, M. (2016): "Regional Convergence in Europe". In: *Intereconomics*, **51**, 3, p. 165-171.
- Young, A. T./Higgins, M. J./Levy, D. (2008): "Sigma Convergence versus Beta Convergence: Evidence from U.S. County-Level Data". In: *Journal of Money, Credit and Banking*, **40**, 5, p. 1083-1093.

See Also

[betaconv.nls](#), [betaconv.ols](#), [sigmaconv](#), [sigmaconv.t](#), [cv](#), [sd2](#), [var2](#)

Examples

```
speed <- betaconv.speed(-0.008070533, 1)
speed[1] # lambda
speed[2] # half-life
```

 conc

Measures of industry concentration

Description

Calculating three measures of industry concentration (Gini, Krugman, Hoover) for a set of I industries

Usage

```
conc(e_ij, industry.id, region.id, na.rm = TRUE)
```

Arguments

e_ij	a numeric vector with the employment of the industry i in region j
industry.id	a vector containing the IDs of the industries i
region.id	a vector containing the IDs of the regions j
na.rm	logical argument that indicates whether NA values should be excluded before computing results

Details

This function is a convenient wrapper for all functions calculating measures of spatial concentration of industries (Gini, Krugman, Hoover)

Value

A matrix with three columns (Gini coefficient, Krugman coefficient, Hoover coefficient) and I rows (one for each regarded industry).

Author(s)

Thomas Wieland

References

Farhauer, O./Kroell, A. (2014): "Standorttheorien: Regional- und Stadtoekonomik in Theorie und Praxis". Wiesbaden : Springer.

Schaetzl, L. (2000): "Wirtschaftsgeographie 2: Empirie". Paderborn : Schoeningh.

See Also

[gini.conc](#), [krugman.conc2](#), [hoover](#)

Examples

```
data(G.regions.industries)

conc_i <- conc (e_ij = G.regions.industries$emp_all,
              industry.id = G.regions.industries$ind_code,
              region.id = G.regions.industries$region_code)
```

 converse

Breaking point formula by Converse

Description

Calculating the breaking point between two cities or retail locations

Usage

```
converse(P_a, P_b, D_ab)
```

Arguments

P_a	a single numeric value of attractivity/population size of location/city <i>a</i>
P_b	a single numeric value of attractivity/population size of location/city <i>b</i>
D_ab	a single numeric value of the transport costs (e.g. distance) between <i>a</i> and <i>b</i>

Details

The *breaking point formula* by Converse (1949) is a modification of the *law of retail gravitation* by Reilly (1929, 1931) (see the functions `reilly` and `reilly.lambda`). The aim of the calculation is to determine the boundaries of the market areas between two locations/cities in consideration of their attractivity/population size and the transport costs (e.g. distance) between them. The models by Reilly and Converse are simple *spatial interaction models* and are considered as *deterministic market area models* due to their exact allocation of demand origins to locations. A probabilistic approach including a theoretical framework was developed by Huff (1962) (see the function `huff`).

Value

a list with two values (B_a: distance from location *a* to breaking point, B_b: distance from location *b* to breaking point)

Author(s)

Thomas Wieland

References

- Berman, B. R./Evans, J. R. (2012): “Retail Management: A Strategic Approach”. 12th edition. Boston : Pearson.
- Converse, P. D. (1949): “New Laws of Retail Gravitation”. In: *Journal of Marketing*, **14**, 3, p. 379-384.
- Huff, D. L. (1962): “Determination of Intra-Urban Retail Trade Areas”. Los Angeles : University of California.
- Levy, M./Weitz, B. A. (2012): “Retailing management”. 8th edition. New York : McGraw-Hill Irwin.
- Loeffler, G. (1998): “Market areas - a methodological reflection on their boundaries”. In: *GeoJournal*, **45**, 4, p. 265-272
- Reilly, W. J. (1929): “Methods for the Study of Retail Relationships”. *Studies in Marketing*, **4**. Austin : Bureau of Business Research, The University of Texas.
- Reilly, W. J. (1931): “The Law of Retail Gravitation”. New York.

See Also

[huff, reilly](#)

Examples

```
# Example from Huff (1962):
converse (400000, 200000, 80)
# two cities (population 400.000 and 200.000 with a distance separating them of 80 miles)
```

coulter

Coulter Coefficient

Description

Calculating the Coulter Coefficient e.g. with respect to regional income

Usage

```
coulter(x, weighting = NULL, na.rm = TRUE)
```

Arguments

x	A numeric vector (e.g. dataset of regional income)
weighting	a weighting vector, e.g. population
na.rm	logical argument that indicates whether NA values should be excluded before computing results

Details

The *Coulter Coefficient* (*CC*) varies between 0 (no inequality/concentration) and 1 (complete inequality/concentration). It can be used for economic inequality and/or regional disparities (Portnov/Felsenstein 2010).

Value

A single numeric value of the *Coulter Coefficient* ($0 < CC < 1$).

Author(s)

Thomas Wieland

References

Portnov, B.A./Felsenstein, D. (2010): "On the suitability of income inequality measures for regional analysis: Some evidence from simulation analysis and bootstrapping tests". In: *Socio-Economic Planning Sciences*, **44**, 4, p. 212-219.

See Also

[cv](#), [gini](#), [gini2](#), [herf](#), [theil](#), [hoover](#), [atkinson](#), [dalton](#), [disp](#)

Examples

```
bip <- c(400,400,400, 400, NA)
bev <- c(1,1,1,200, NA)
coulter(bip, bev)
```

curvefit

Curve fitting

Description

Curve fitting (similar to SPSS and Excel)

Usage

```
curvefit(x, y, y.max = NULL, extrapol = NULL,
plot.curves = TRUE, pcol = "black", ptype = 19, psize = 1,
lin.col = "blue", pow.col = "green", exp.col = "orange", logi.col = "red",
plot.title = "Curve fitting", plot.legend = TRUE,
xlab = "x", ylab = "y", y.min = NULL, ..., print.results = TRUE)
```

Arguments

<code>x</code>	a numeric vector containing the explanatory variable
<code>y</code>	a numeric vector containing the dependent variable
<code>y.max</code>	Optional: given maximum for the logistic regression function
<code>extrapol</code>	a single numeric value for how many x units the dependent variable y shall be extrapolated
<code>plot.curves</code>	Logical argument that indicates whether the curves shall be plotted or not
<code>pcol</code>	If <code>plot.curves = TRUE</code> : Point color
<code>ptype</code>	If <code>plot.curves = TRUE</code> : Point type (pch)
<code>psize</code>	If <code>plot.curves = TRUE</code> : Point size
<code>lin.col</code>	If <code>plot.curves = TRUE</code> : Color of linear regression line
<code>pow.col</code>	If <code>plot.curves = TRUE</code> : Color of power function regression line
<code>exp.col</code>	If <code>plot.curves = TRUE</code> : Color of exponential function regression line
<code>logi.col</code>	If <code>plot.curves = TRUE</code> : Color of logistic function regression line
<code>plot.title</code>	If <code>plot.curves = TRUE</code> : Plot title
<code>plot.legend</code>	If <code>plot.curves = TRUE</code> : Logical argument that indicates whether a legend is added to the plot or not
<code>xlab</code>	If <code>plot.curves = TRUE</code> : X axis label
<code>ylab</code>	If <code>plot.curves = TRUE</code> : Y axis label
<code>y.min</code>	Optional: Y axis minimum
<code>...</code>	Optional: other plot parameters
<code>print.results</code>	Logical argument that indicates whether the model results are shown or not

Details

Curve fitting for a given independent and dependent variable ($y = f(x)$). Similar to curve fitting in SPSS or Excel. Fitting of nonlinear regression models (power, exponential, logistic) via intrinsically linear models (Rawlings et al. 1998).

Value

A data frame containing the regression results (Parameters a and b, std. errors, t values, ...)

Author(s)

Thomas Wieland

References

Rawlings, J. O./Pantula, S. G./Dickey, D. A. (1998): "Applied Regression Analysis". Springer. 2nd edition.

Examples

```

x <- 1:20
y <- 3-2*x
curvefit(x, y, plot.curves = TRUE)
# fit with plot
curvefit(x, y, extrapol=10, plot.curves = TRUE)
# fit and extrapolation with plot

x <- runif(20, min = 0, max = 100)
# some random data

# linear
y_resid <- runif(20, min = 0, max = 10)
# random residuals
y <- 3+(-0.112*x)+y_resid
curvefit(x, y)

# power
y_resid <- runif(20, min = 0.1, max = 0.2)
# random residuals
y <- 3*(x^-0.112)*y_resid
curvefit(x, y)

# exponential
y_resid <- runif(20, min = 0.1, max = 0.2)
# random residuals
y <- 3*exp(-0.112*x)*y_resid
curvefit(x, y)

# logistic
y_resid <- runif(20, min = 0.1, max = 0.2)
# random residuals
y <- 100/(1+exp(3+(-0.112*x)))*y_resid
curvefit(x, y)

```

cv

Coefficient of variation

Description

Calculating the coefficient of variation (cv), standardized and non-standardized, weighted and non-weighted

Usage

```

cv (x, is.sample = TRUE, coefnorm = FALSE, weighting = NULL,
wmean = FALSE, na.rm = TRUE)

```

Arguments

<code>x</code>	a numeric vector
<code>is.sample</code>	logical argument that indicates if the dataset is a sample or the population (default: <code>is.sample = TRUE</code> , so the denominator of variance is $n - 1$)
<code>coefnorm</code>	logical argument that indicates if the function output is the standardized cv ($0 < v^* < 1$) or not ($0 < v < \infty$) (default: <code>coefnorm = FALSE</code>)
<code>weighting</code>	a numeric vector containing weighting data to compute the weighted coefficient of variation (instead of the non-weighted cv)
<code>wmean</code>	logical argument that indicates if the weighted mean is used when calculating the weighted coefficient of variation
<code>na.rm</code>	logical argument that whether NA values should be extracted or not

Details

The *coefficient of variation*, v , is a dimensionless measure of statistical dispersion ($0 < v < \infty$), based on variance and standard deviation, respectively. From a regional economic perspective, it is closely linked to the concept of *sigma convergence* (σ) which means a harmonization of regional economic output or income over time, while the other type of convergence, *beta convergence* (β), means a decline of dispersion because poor regions have a stronger growth than rich regions (Capello/Nijkamp 2009). The cv allows to summarize regional disparities (e.g. disparities in regional GDP per capita) in one indicator and is more frequently used for this purpose than the standard deviation, especially in analyzing of σ convergence over a long period (e.g. Lessmann 2005, Huang/Leung 2009, Siljak 2015). But the cv can also be used for any other types of disparities or dispersion, such as disparities in supply (e.g. density of physicians or grocery stores).

The cv (variance, standard deviation) can be weighted by using a second weighting vector. As there is more than one way to weight measures of statistical dispersion, this function uses the formula for the weighted cv (v_w) from Sheret (1984). The cv can be standardized, while this function uses the formula for the standardized cv (v^* , with $0 < v^* < 1$) from Kohn/Oetztuerk (2013). The vector `x` is automatically treated as a sample (such as in the base `sd` function), so the denominator of variance is $n - 1$, if it is not, set `is.sample = FALSE`.

Value

Single numeric value. If `coefnorm = FALSE` the function returns the non-standardized cv ($0 < v < \infty$). If `coefnorm = TRUE` the standardized cv ($0 < v^* < 1$) is returned.

Author(s)

Thomas Wieland

References

- Bahrenberg, G./Giese, E./Mevenkamp, N./Nipper, J. (2010): "Statistische Methoden in der Geographie. Band 1: Univariate und bivariate Statistik". Stuttgart: Bortraeger.
- Capello, R./Nijkamp, P. (2009): "Introduction: regional growth and development theories in the twenty-first century - recent theoretical advances and future challenges". In: Capello, R./Nijkamp, P. (eds.): *Handbook of Regional Growth and Development Theories*. Cheltenham: Elgar. p. 1-16.

Lessmann, C. (2005): “Regionale Disparitaeten in Deutschland und ausgesuchten OECD-Staaten im Vergleich”. *ifo Dresden berichtet*, **3/2005**. https://www.ifo.de/DocDL/ifodb_2005_3_25-33.pdf.

Huang, Y./Leung, Y. (2009): “Measuring Regional Inequality: A Comparison of Coefficient of Variation and Hoover Concentration Index”. In: *The Open Geography Journal*, **2**, p. 25-34.

Kohn, W./Oeztuerk, R. (2013): “Statistik fuer Oekonomen. Datenanalyse mit R und SPSS”. Berlin: Springer.

Sheret, M. (1984): “The Coefficient of Variation: Weighting Considerations”. In: *Social Indicators Research*, **15**, 3, p. 289-295.

Siljak, D. (2015): “Real Economic Convergence in Western Europe from 1995 to 2013”. In: *International Journal of Business and Economic Development*, **3**, 3, p. 56-67.

See Also

[gini](#), [herf](#), [hoover](#), [rca](#)

Examples

```
# Regional disparities / sigma convergence in Germany
data(G.counties.gdp)
# GDP per capita for German counties (Landkreise)
cvs <- apply (G.counties.gdp[54:68], MARGIN = 2, FUN = cv)
# Calculating cv for the years 2000-2014
years <- 2000:2014
plot(years, cvs, "l", ylim=c(0.3,0.6), xlab = "year",
      ylab = "CV of GDP per capita")
# Plot cv over time
```

dalton

Dalton Inequality Index

Description

Calculating the Dalton Inequality Index e.g. with respect to regional income

Usage

```
dalton(x, na.rm = TRUE)
```

Arguments

x	A numeric vector (e.g. dataset of regional income)
na.rm	logical argument that indicates whether NA values should be excluded before computing results

Details

The *Dalton Inequality Index* (δ) can be used for economic inequality and/or regional disparities (Portnov/Felsenstein 2010).

Value

A single numeric value of the *Dalton Inequality Index*.

Author(s)

Thomas Wieland

References

Portnov, B.A./Felsenstein, D. (2010): "On the suitability of income inequality measures for regional analysis: Some evidence from simulation analysis and bootstrapping tests". In: *Socio-Economic Planning Sciences*, **44**, 4, p. 212-219.

See Also

[cv](#), [gini](#), [gini2](#), [herf](#), [theil](#), [hoover](#), [coulter](#), [dalton](#), [disp](#)

Examples

```
dalton (c(10,10,10,10))
```

```
dalton (c(10,0,0,0))
```

```
dalton (c(10,1,1,1))
```

 disp

Concentration/inequality/dispersion measures

Description

Calculating a set of concentration/inequality/dispersion measures

Usage

```
disp(x, weighting = NULL, at.epsilon = 0.5, na.rm = TRUE)
```

Arguments

x	a numeric vector or matrix or columns from a data frame
weighting	a weighting vector, e.g. population
at.epsilon	Weighting parameter ϵ for the Atkinson index
na.rm	logical argument that indicates whether NA values should be excluded before computing results

Details

This function is a convenient wrapper for all functions calculating concentration/inequality measures.

Value

A matrix containing the concentration/inequality measures.

Author(s)

Thomas Wieland

References

Gluschenko, K. (2018): "Measuring regional inequality: to weight or not to weight?" In: *Spatial Economic Analysis*, **13**, 1, p. 36-59.

Portnov, B.A./Felsenstein, D. (2010): "On the suitability of income inequality measures for regional analysis: Some evidence from simulation analysis and bootstrapping tests". In: *Socio-Economic Planning Sciences*, **44**, 4, p. 212-219.

See Also

[atkinson](#), [coulter](#), [dalton](#), [cv](#), [gini2](#), [herf](#), [hoover](#), [sd2](#), [theil](#), [williamson](#)

Examples

```
data(Automotive)

disp(Automotive$Turnover2008)
disp(Automotive[4:8])
```

dist.buf

Counting points in a buffer

Description

Counting points within a buffer of a given distance with points with given coordinates

Usage

```
dist.buf(startpoints, sp_id, lat_start, lon_start, endpoints, ep_id, lat_end, lon_end,
ep_sum = NULL, bufdist = 500, extract_local = TRUE, unit = "m")
```

Arguments

startpoints	A data frame containing the start points
sp_id	Column containing the IDs of the startpoints in the data frame startpoints
lat_start	Column containing the latitudes of the start points in the data frame startpoints
lon_start	Column containing the longitudes of the start points in the data frame startpoints
endpoints	A data frame containing the points to count
ep_id	Column containing the IDs of the points to count in the data frame endpoints
lat_end	Column containing the latitudes of the points to count in the data frame endpoints
lon_end	Column containing the longitudes of the points to count in the data frame endpoints
ep_sum	Column of an additional variable in the data frame endpoints to sum
bufdist	The buffer distance
extract_local	Logical argument that indicates if the start points should be included or not (default: TRUE)
unit	Unit of the buffer distance: unit="m" for meters, unit="km" for kilometers or unit="miles" for miles

Details

The function is based on the idea of a buffer analysis in GIS (Geographic Information System), e.g. to count the points of interest within a given buffer distance.

Value

The function returns a list containing:

count_table	A data.frame containing two columns: The start point IDs (from) and the number of counted points in the given buffer distance (count_location)
distmat	A data.frame containing the corresponding distance matrix with $I \times J$ rows

Author(s)

Thomas Wieland

References

- de Lange, N. (2013): "Geoinformatik in Theorie und Praxis". 3rd edition. Berlin : Springer Spektrum.
- Krider, R. E./Putler, R. S. (2013): "Which Birds of a Feather Flock Together? Clustering and Avoidance Patterns of Similar Retail Outlets". In: *Geographical Analysis*, **45**, 2, p. 123-149

See Also

[dist](#), [dist.mat](#)

Examples

```
citynames <- c("Goettingen", "Karlsruhe", "Freiburg")
lat <- c(51.556307, 49.009603, 47.9874)
lon <- c(9.947375, 8.417004, 7.8945)
citynames <- c("Goettingen", "Karlsruhe", "Freiburg")
cities <- data.frame(citynames, lat, lon)
dist.mat (cities, "citynames", "lat", "lon", cities, "citynames", "lat", "lon")
# Euclidean distance matrix (3 x 3 cities = 9 distances)
dist.buf (cities, "citynames", "lat", "lon", cities, "citynames", "lat", "lon", bufdist = 300000)
# Cities within 300 km
```

dist.calc

Euclidean distance between coordinates

Description

Calculation of the euclidean distance between two points with stated coordinates (lat, lon)

Usage

```
dist.calc(lat1, lon1, lat2, lon2, unit = "km")
```

Arguments

lat1	Latitude of the regarded start point
lon1	Longitude of the regarded start point
lat2	Latitude of the regarded end point
lon2	Longitude of the regarded end point
unit	Unit of the resulting distance: unit="m" for meters, unit="km" for kilometers or unit="miles" for miles

Value

A single numeric value

Author(s)

Thomas Wieland

See Also

[dist.buf](#), [dist.mat](#)

Examples

```
dist.calc(51.556307, 9.947375, 49.009603, 8.417004)
# about 304 kilometers
```

 dist.mat

Euclidean distance matrix between points

Description

Calculation of an euclidean distance matrix between points with stated coordinates (lat, lon)

Usage

```
dist.mat(startpoints, sp_id, lat_start, lon_start, endpoints, ep_id,
lat_end, lon_end, unit = "km")
```

Arguments

startpoints	A data frame containing the start points
sp_id	Column containing the IDs of the startpoints in the data frame startpoints
lat_start	Column containing the latitudes of the start points in the data frame startpoints
lon_start	Column containing the longitudes of the start points in the data frame startpoints
endpoints	A data frame containing the end points
ep_id	Column containing the IDs of the endpoints in the data frame endpoints
lat_end	Column containing the latitudes of the end points in the data frame endpoints
lon_end	Column containing the longitudes of the end points in the data frame endpoints
unit	Unit of the resulting distance: unit="m" for meters, unit="km" for kilometers or unit="miles" for miles

Details

The function calculates an euclidean distance matrix between points with stated coordinates (lat and lon). While m start points and n end points are given, the output is a linear $m * n$ distance matrix.

Value

The function returns a data.frame containing 4 columns: The start point IDs (from), the end point IDs (to), the combination of both (from_to) and the calculated distance (distance).

Author(s)

Thomas Wieland

References

de Lange, N. (2013): "Geoinformatik in Theorie und Praxis". 3rd edition. Berlin : Springer Spektrum.

Krider, R. E./Putler, R. S. (2013): "Which Birds of a Feather Flock Together? Clustering and Avoidance Patterns of Similar Retail Outlets". In: *Geographical Analysis*, **45**, 2, p. 123-149

See Also[dist](#), [dist.buf](#)**Examples**

```
citynames <- c("Goettingen", "Karlsruhe", "Freiburg")
lat <- c(51.556307, 49.009603, 47.9874)
lon <- c(9.947375, 8.417004, 7.8945)
citynames <- c("Goettingen", "Karlsruhe", "Freiburg")
cities <- data.frame(citynames, lat, lon)
dist.mat (cities, "citynames", "lat", "lon", cities, "citynames", "lat", "lon")
# Euclidean distance matrix (3 x 3 cities = 9 distances)
dist.buf (cities, "citynames", "lat", "lon", cities, "citynames", "lat", "lon", bufdist = 300000)
# Cities within 300 km
```

durpug

Relative diversity index by Duranton and Puga

Description

Calculating the relative diversity index (RDI) by Duranton and Puga based on regional industry data (normally employment data)

Usage

```
durpug(e_ij, e_i)
```

Arguments

`e_ij` a numeric vector with the employment of the industries i in region j
`e_i` a numeric vector with the all-over employment in the industries i

Value

A single numeric value of *RDI*

Author(s)

Thomas Wieland

References

Duranton, G./Puga, D. (2000): "Diversity and Specialisation in Cities: Why, Where and When Does it Matter?". In: *Urban Studies*, **37**, 3, p. 533-555.
Farhauer, O./Kroell, A. (2013): "Standorttheorien: Regional- und Stadtoekonomik in Theorie und Praxis". Wiesbaden : Springer.

See Also

[gini.spec](#), [krugman.spec](#), [hoover](#)

Examples

```
# Example Goettingen:

data(Goettingen)
# Loads the data

durpug (Goettingen$Goettingen2008[2:13], Goettingen$BRD2008[2:13])
# Returns the Duranton-Puga RDI for Goettingen
```

ellison.a

Ellison-Glaeser Agglomeration Index

Description

Calculating the Agglomeration Index by Ellison and Glaeser for a single industry i

Usage

```
ellison.a(e_ik, e_j, regions, print.results = TRUE)
```

Arguments

e_ik	a numeric vector containing the no. of employees of firm k from industry i
e_j	a numeric vector containing the no. of employees in the regions j
regions	a vector containing the IDs/names of the regions j
print.results	logical argument that indicates whether the function prints the results or not (only for internal use)

Details

The Ellison-Glaeser Agglomeration Index is not standardized. A value of $\gamma_i = 0$ indicates a spatial distribution of firms equal to a dartboard approach. Values below zero indicate spatial dispersion, values greater than zero indicate clustering.

Value

A matrix with five columns (γ_i , G_i , z_{G_i} , K_i and HHI_i).

Author(s)

Thomas Wieland

References

- Ellison G./Glaeser, E. (1997): “Geographic concentration in u.s. manufacturing industries: A dartboard approach”. In: *Journal of Political Economy*, **105**, 5, p. 889-927.
- Farhauer, O./Kroell, A. (2014): “Standorttheorien: Regional- und Stadtoekonomik in Theorie und Praxis”. Wiesbaden : Springer.
- Nakamura R./Morrison Paul, C. (2009): “Measuring agglomeration”. In: Capello, R./Nijkamp, P. (eds): *Handbook of Regional Growth and Development Theories*, p. 305-328.

See Also

[gini.conc](#), [gini.spec](#), [locq](#), [locq2](#), [howard.cl](#), [howard.xcl](#), [howard.xcl2](#), [litzenberger](#), [litzenberger2](#)

Examples

```
# Example from Farhauer/Kroell (2014):
j <- c("Wien", "Wien", "Wien", "Wien", "Wien", "Linz",
      "Linz", "Linz", "Linz", "Graz")
E_ik <- c(200,650,12000,100,50,16000,13000,1500,1500,25000)
E_j <- c(500000,400000,100000)
ellison.a(E_ik, E_j, j)
# 0.05990628
```

ellison.a2

Ellison-Glaeser Agglomeration Index

Description

Calculating the Agglomeration Index by Ellison and Glaeser for a given number of I industries

Usage

```
ellison.a2(e_ik, industry, region, print.results = TRUE)
```

Arguments

<code>e_ik</code>	a numeric vector containing the no. of employees of firm k from industry i
<code>industry</code>	a vector containing the IDs/names of the industries i
<code>region</code>	a vector containing the IDs/names of the regions j
<code>print.results</code>	logical argument that indicates whether the function prints the results or not (only for internal use)

Details

The Ellison-Glaeser Agglomeration Index is not standardized. A value of $\gamma_i = 0$ indicates a spatial distribution of firms equal to a dartboard approach. Values below zero indicate spatial dispersion, values greater than zero indicate clustering.

Value

A matrix with five columns (γ_i , G_i , z_{G_i} , K_i and HHI_i) and I rows (one for each industry).

Author(s)

Thomas Wieland

References

Ellison G./Glaeser, E. (1997): “Geographic concentration in u.s. manufacturing industries: A dashboard approach”. In: *Journal of Political Economy*, **105**, 5, p. 889-927.

Farhauer, O./Kroell, A. (2014): “Standorttheorien: Regional- und Stadtoekonomik in Theorie und Praxis”. Wiesbaden : Springer.

Nakamura R./Morrison Paul, C. (2009): “Measuring agglomeration”. In: Capello, R./Nijkamp, P. (eds): *Handbook of Regional Growth and Development Theories*, p. 305-328.

See Also

[ellison.a](#), [gini.conc](#), [gini.spec](#), [locq](#), [locq2](#), [howard.cl](#), [howard.xcl](#), [howard.xcl2](#), [litzenberger](#), [litzenberger2](#)

Examples

```
# Example data from Farhauer/Kroell (2014):
data(FK2014_EGC)

ellison.a2 (FK2014_EGC$emp_firm, FK2014_EGC$industry,
FK2014_EGC$region)
```

ellison.c

Ellison-Glaeser Coagglomeration Index

Description

Calculating the Coagglomeration Index by Ellison and Glaeser for one set of U industries

Usage

```
ellison.c(e_ik, industry, region, e_j = NULL, c.industries = NULL)
```

Arguments

e_ik	a numeric vector containing the no. of employees of firm k from industry i
industry	a vector containing the IDs/names of the industries i
region	a vector containing the IDs/names of the regions j
e_j	a numeric vector containing the total employment of the regions j
c.industries	optional: a vector containing the regarded U industries (where $U \leq I$)

Details

The Ellison-Glaeser Coagglomeration Index is not standardized. A value of $\gamma_c = 0$ indicates a spatial distribution of firms equal to a dartboard approach. Values below zero indicate spatial dispersion, values greater than zero indicate clustering.

Value

A single value of γ_c

Author(s)

Thomas Wieland

References

Ellison G./Glaeser, E. (1997): “Geographic concentration in u.s. manufacturing industries: A dartboard approach”. In: *Journal of Political Economy*, **105**, 5, p. 889-927.

Farhauer, O./Kroell, A. (2014): “Standorttheorien: Regional- und Stadtoekonomik in Theorie und Praxis”. Wiesbaden : Springer.

Nakamura R./Morrison Paul, C. (2009): “Measuring agglomeration”. In: Capello, R./Nijkamp, P. (eds): *Handbook of Regional Growth and Development Theories*, p. 305-328.

See Also

[ellison.a](#), [ellison.a2](#), [ellison.c2](#), [gini.conc](#), [gini.spec](#), [locq](#), [locq2](#), [howard.cl](#), [howard.xcl](#), [howard.xcl2](#), [litzengerger](#), [litzengerger2](#)

Examples

```
# Example from Farhauer/Kroell (2014):
data(FK2014_EGC)

ellison.c(FK2014_EGC$emp_firm, FK2014_EGC$industry,
FK2014_EGC$region, FK2014_EGC$emp_region)
```

ellison.c2

Ellison-Glaeser Coagglomeration Index

Description

Calculating the Coagglomeration Index by Ellison and Glaeser for $I \times I$ sets of two industries

Usage

```
ellison.c2(e_ik, industry, region, e_j = NULL, print.results = TRUE)
```

Arguments

e_ik	a numeric vector containing the no. of employees of firm k from industry i
industry	a vector containing the IDs/names of the industries i
region	a vector containing the IDs/names of the regions j
e_j	a numeric vector containing the total employment of the regions j
print.results	logical argument that indicates whether the results are printed or not (for internal use)

Details

The Ellison-Glaeser Coagglomeration Index is not standardized. A value of $\gamma^c = 0$ indicates a spatial distribution of firms equal to a dartboard approach. Values below zero indicate spatial dispersion, values greater than zero indicate clustering.

Value

A single value of γ^c

Author(s)

Thomas Wieland

References

- Ellison G./Glaeser, E. (1997): “Geographic concentration in u.s. manufacturing industries: A dartboard approach”. In: *Journal of Political Economy*, **105**, 5, p. 889-927.
- Farhauer, O./Kroell, A. (2014): “Standorttheorien: Regional- und Stadtoekonomik in Theorie und Praxis”. Wiesbaden : Springer.
- Nakamura R./Morrison Paul, C. (2009): “Measuring agglomeration”. In: Capello, R./Nijkamp, P. (eds): *Handbook of Regional Growth and Development Theories*, p. 305-328.

See Also

[ellison.a](#), [ellison.a2](#), [ellison.c](#), [gini.conc](#), [gini.spec](#), [locq](#), [locq2](#), [howard.cl](#), [howard.xcl](#), [howard.xcl2](#), [litzenberger](#), [litzenberger2](#)

Examples

```
# Example from Farhauer/Kroell (2014):
data(FK2014_EGC)

ellison.c2(FK2014_EGC$emp_firm, FK2014_EGC$industry,
FK2014_EGC$region, FK2014_EGC$emp_region)
# this may take a while
```

`EU28.emp`*Eurostat national employment data 2004-2016*

Description

Employment data for EU countries 2004-2016 (Source: Eurostat)

Usage

```
data("EU28.emp")
```

Format

A data frame with 3000 observations on the following 7 variables.

`unit` measuring unit: thousand persons (THS_PER)

`nace_r2` NACE industry classification

`s_adj` Adjustment of data: Not seasonally adjusted data (NSA)

`na_item` a factor with levels SAL_DC

`geo` NUTS nation code

`time` year

`emp1000` Industry-specific employment in thousand persons

Source

Eurostat (2018): Breakdowns of GDP aggregates and employment data by main industries and asset classes, Tab. code `namq_10_a10_e`. http://appsso.eurostat.ec.europa.eu/nui/show.do?dataset=namq_10_a10_e. Own postprocessing.

Examples

```
data(EU28.emp)
EU28.emp[EU28.emp$time == 2016,]
# only data for 2016
```

FK2014_EGC

Fictional sample data of 42 firms

Description

Dataset with 42 firms from 4 industries in 3 regions (fictional sample data from Farhauer/Kroell 2014)

Usage

```
data("FK2014_EGC")
```

Format

A data frame with 42 observations on the following 5 variables.

region unique ID of the region

industry name of the industry (German language)

firm firm ID

emp_firm each firm's no. of employees

emp_region total employment of the region

Source

Farhauer, O./Kroell, A. (2014): "Standorttheorien: Regional- und Stadtoekonomik in Theorie und Praxis". Wiesbaden : Springer.

References

Farhauer, O./Kroell, A. (2014): "Standorttheorien: Regional- und Stadtoekonomik in Theorie und Praxis". Wiesbaden : Springer.

Examples

```
# Example from Farhauer/Kroell (2014):  
data(FK2014_EGC)  
  
ellison.c(FK2014_EGC$emp_firm, FK2014_EGC$industry,  
FK2014_EGC$region, FK2014_EGC$emp_region)
```

Freiburg

Employment data in Freiburg and Germany

Description

Dataset with industry-specific employment in Freiburg and Germany in the years 2008 and 2014

Usage

```
data("Freiburg")
```

Format

A data frame with 9 observations on the following 8 variables.

industry a factor with levels for the regarded industry based on the German official economic statistics (WZ2008)

e_Freiburg2008 a numeric vector with industry-specific employment in Freiburg 2008

e_Freiburg2014 a numeric vector with industry-specific employment in Freiburg 2014

e_g_Freiburg_0814 a numeric vector containing the growth of industry-specific employment in Freiburg 2008-2014, percentage

e_Germany2008 a numeric vector with industry-specific employment in Germany 2008

e_Germany2014 a numeric vector with industry-specific employment in Germany 2014

e_g_Germany_0814 a numeric vector containing the growth of industry-specific employment in Germany 2008-2014, percentage

color a factor containing colors (blue, brown, ...)

Source

Statistische Aemter des Bundes und der Laender: Regionaldatenbank Deutschland, Tab. 254-74-4, own calculations

Examples

```
data(Freiburg)
# Loads the data
growth(Freiburg$e_Freiburg2008, Freiburg$e_Freiburg2014, growth.type = "rate")
# Industry-specific growth rates for Freiburg 2008 to 2014
```

G.counties.gdp	<i>Gross Domestic Product (GDP) per capita for German counties 1992-2014</i>
----------------	--

Description

The dataset contains the Gross Domestic Product (GDP) absolute and per capita (in EUR, at current prices) for the 402 German counties (Landkreise) from 1992 to 2014.

Usage

```
data("G.counties.gdp")
```

Format

A data frame with 402 observations on the following 68 variables.

region_code_EU a factor containing der EU regional code

region_code a factor containing the German regional code

gdp1992 a numeric vector containing the GDP for German counties (Landkreise) for 1992

gdp1994 a numeric vector containing the GDP for German counties (Landkreise) for 1994

gdp1995 a numeric vector containing the GDP for German counties (Landkreise) for 1995

gdp1996 a numeric vector containing the GDP for German counties (Landkreise) for 1996

gdp1997 a numeric vector containing the GDP for German counties (Landkreise) for 1997

gdp1998 a numeric vector containing the GDP for German counties (Landkreise) for 1998

gdp1999 a numeric vector containing the GDP for German counties (Landkreise) for 1999

gdp2000 a numeric vector containing the GDP for German counties (Landkreise) for 2000

gdp2001 a numeric vector containing the GDP for German counties (Landkreise) for 2001

gdp2002 a numeric vector containing the GDP for German counties (Landkreise) for 2002

gdp2003 a numeric vector containing the GDP for German counties (Landkreise) for 2003

gdp2004 a numeric vector containing the GDP for German counties (Landkreise) for 2004

gdp2005 a numeric vector containing the GDP for German counties (Landkreise) for 2005

gdp2006 a numeric vector containing the GDP for German counties (Landkreise) for 2006

gdp2007 a numeric vector containing the GDP for German counties (Landkreise) for 2007

gdp2008 a numeric vector containing the GDP for German counties (Landkreise) for 2008

gdp2009 a numeric vector containing the GDP for German counties (Landkreise) for 2009

gdp2010 a numeric vector containing the GDP for German counties (Landkreise) for 2010

gdp2011 a numeric vector containing the GDP for German counties (Landkreise) for 2011

gdp2012 a numeric vector containing the GDP for German counties (Landkreise) for 2012

gdp2013 a numeric vector containing the GDP for German counties (Landkreise) for 2013

gdp2014 a numeric vector containing the GDP for German counties (Landkreise) for 2014

pop1992 a numeric vector containing the population for German counties (Landkreise) for 1992

pop1994 a numeric vector containing the population for German counties (Landkreise) for 1994

pop1995 a numeric vector containing the population for German counties (Landkreise) for 1995

pop1996 a numeric vector containing the population for German counties (Landkreise) for 1996

pop1997 a numeric vector containing the population for German counties (Landkreise) for 1997

pop1998 a numeric vector containing the population for German counties (Landkreise) for 1998

pop1999 a numeric vector containing the population for German counties (Landkreise) for 1999

pop2000 a numeric vector containing the population for German counties (Landkreise) for 2000

pop2001 a numeric vector containing the population for German counties (Landkreise) for 2001

pop2002 a numeric vector containing the population for German counties (Landkreise) for 2002

pop2003 a numeric vector containing the population for German counties (Landkreise) for 2003

pop2004 a numeric vector containing the population for German counties (Landkreise) for 2004

pop2005 a numeric vector containing the population for German counties (Landkreise) for 2005

pop2006 a numeric vector containing the population for German counties (Landkreise) for 2006

pop2007 a numeric vector containing the population for German counties (Landkreise) for 2007

pop2008 a numeric vector containing the population for German counties (Landkreise) for 2008

pop2009 a numeric vector containing the population for German counties (Landkreise) for 2009

pop2010 a numeric vector containing the population for German counties (Landkreise) for 2010

pop2011 a numeric vector containing the population for German counties (Landkreise) for 2011

pop2012 a numeric vector containing the population for German counties (Landkreise) for 2012

pop2013 a numeric vector containing the population for German counties (Landkreise) for 2013

pop2014 a numeric vector containing the population for German counties (Landkreise) for 2014

gdppc1992 a numeric vector containing the GDP per capita for German counties (Landkreise) for 1992

gdppc1994 a numeric vector containing the GDP per capita for German counties (Landkreise) for 1994

gdppc1995 a numeric vector containing the GDP per capita for German counties (Landkreise) for 1995

gdppc1996 a numeric vector containing the GDP per capita for German counties (Landkreise) for 1996

gdppc1997 a numeric vector containing the GDP per capita for German counties (Landkreise) for 1997

gdppc1998 a numeric vector containing the GDP per capita for German counties (Landkreise) for 1998

gdppc1999 a numeric vector containing the GDP per capita for German counties (Landkreise) for 1999

gdppc2000 a numeric vector containing the GDP per capita for German counties (Landkreise) for 2000

gdppc2001 a numeric vector containing the GDP per capita for German counties (Landkreise) for 2001

gdppc2002 a numeric vector containing the GDP per capita for German counties (Landkreise) for 2002

gdppc2003 a numeric vector containing the GDP per capita for German counties (Landkreise) for 2003

gdppc2004 a numeric vector containing the GDP per capita for German counties (Landkreise) for 2004

gdppc2005 a numeric vector containing the GDP per capita for German counties (Landkreise) for 2005

gdppc2006 a numeric vector containing the GDP per capita for German counties (Landkreise) for 2006

gdppc2007 a numeric vector containing the GDP per capita for German counties (Landkreise) for 2007

gdppc2008 a numeric vector containing the GDP per capita for German counties (Landkreise) for 2008

gdppc2009 a numeric vector containing the GDP per capita for German counties (Landkreise) for 2009

gdppc2010 a numeric vector containing the GDP per capita for German counties (Landkreise) for 2010

gdppc2011 a numeric vector containing the GDP per capita for German counties (Landkreise) for 2011

gdppc2012 a numeric vector containing the GDP per capita for German counties (Landkreise) for 2012

gdppc2013 a numeric vector containing the GDP per capita for German counties (Landkreise) for 2013

gdppc2014 a numeric vector containing the GDP per capita for German counties (Landkreise) for 2014

regional Region West or East

Details

For the years 1992 to 1999, the GDP data is incomplete.

Source

Arbeitskreis "Volkswirtschaftliche Gesamtrechnungen der Laender" im Auftrag der Statistischen Aemter der 16 Bundeslaender, des Statistischen Bundesamtes und des Buergeramtes, Statistik und Wahlen, Frankfurt a. M. (2016): "Bruttoinlandsprodukt, Bruttowertschoepfung in den kreisfreien Staedten und Landkreisen der Bundesrepublik Deutschland 1992 und 1994 bis 2014".

References

Arbeitskreis "Volkswirtschaftliche Gesamtrechnungen der Laender" im Auftrag der Statistischen Aemter der 16 Bundeslaender, des Statistischen Bundesamtes und des Buergeramtes, Statistik und Wahlen, Frankfurt a. M. (2016): "Bruttoinlandsprodukt, Bruttowertschoepfung in den kreisfreien Staedten und Landkreisen der Bundesrepublik Deutschland 1992 und 1994 bis 2014".

Examples

```
# Regional disparities / sigma convergence in Germany
data(G.counties.gdp)
# GDP per capita for German counties (Landkreise)
cvs <- apply (G.counties.gdp[54:68], MARGIN = 2, FUN = cv)
# Calculating cv for the years 2000-2014
years <- 2000:2014
plot(years, cvs, "l", ylim=c(0.3,0.6), xlab = "year",
ylab = "CV of GDP per capita")
# Plot cv over time
```

G.regions.emp

Employment data for German regions 2008-2014

Description

The dataset contains the industry-specific employment in the German region ("Bundeslaender") for the years 2008 to 2014.

Usage

```
data("G.regions.emp")
```

Format

A data frame with 1428 observations on the following 4 variables.

industry a factor containing the industry (in German language, e.g. "Baugewerbe" = construction, "Handel, Gastgewerbe, Verkehr (G-I)" = retail, hospitality industry and transport industry)

region a factor containing the names of the German regions (Bundeslaender)

year a numeric vector containing the related year

emp a numeric vector containing the related number of employees

Source

Statistische Aemter des Bundes und der Laender, Regionaldatenbank (2017): Sozialversicherungspflichtig Beschaeftigte: Beschaeftigte am Arbeitsort nach Geschlecht, Nationalitaet und Wirtschaftszweigen (Beschaeftigungsstatistik der Bundesagentur fuer Arbeit) - Stichtag 30.06. - regionale Ebenen(Tab. 254-74-4-B).

References

Statistische Aemter des Bundes und der Laender, Regionaldatenbank (2017): Sozialversicherungspflichtig Beschaeftigte: Beschaeftigte am Arbeitsort nach Geschlecht, Nationalitaet und Wirtschaftszweigen (Beschaeftigungsstatistik der Bundesagentur fuer Arbeit) - Stichtag 30.06. - regionale Ebenen(Tab. 254-74-4-B).

Examples

```

data(G.regions.emp)
# Concentration of construction industry in Germany
# based on 16 German regions (Bundeslaender) for the year 2008
construction2008 <- G.regions.emp[(G.regions.emp$industry == "Baugewerbe (F)" |
G.regions.emp$industry == "Insgesamt") & G.regions.emp$year == "2008",]
# only data for construction industry (Baugewerbe) and all-over (Insgesamt)
# for the 16 German regions in the year 2008
construction2008 <- construction2008[construction2008$region != "Insgesamt",]
# delete all-over data for all industries
gini.conc(construction2008[construction2008$industry=="Baugewerbe (F)",]$emp,
construction2008[construction2008$industry=="Insgesamt",]$emp)

# Concentration of financial industry in Germany 2008 vs. 2014
# based on 16 German regions (Bundeslaender) for 2008 and 2014
finance2008 <- G.regions.emp[(G.regions.emp$industry ==
"Erbringung von Finanz- und Vers.leistungen (K)" |
G.regions.emp$industry == "Insgesamt") & G.regions.emp$year == "2008",]
finance2008 <- finance2008[finance2008$region != "Insgesamt",]
# delete all-over data for all industries
gini.conc(finance2008[finance2008$industry ==
"Erbringung von Finanz- und Vers.leistungen (K)",]$emp,
finance2008[finance2008$industry=="Insgesamt",]$emp)
finance2014 <- G.regions.emp[(G.regions.emp$industry ==
"Erbringung von Finanz- und Vers.leistungen (K)" | G.regions.emp$industry ==
"Insgesamt") & G.regions.emp$year == "2014",]
finance2014 <- finance2014[finance2014$region != "Insgesamt",]
# delete all-over data for all industries
gini.conc(finance2014[finance2014$industry ==
"Erbringung von Finanz- und Vers.leistungen (K)",]$emp,
finance2014[finance2014$industry=="Insgesamt",]$emp)

```

G.regions.industries *Firms and employment data for German regions 2015*

Description

The dataset contains the industry-specific firm stock and employment in the German regions ("Bundeslaender") for 2015.

Usage

```
data("G.regions.industries")
```

Format

A data frame with 272 observations on the following 9 variables.

year a numeric vector containing the related year

region a factor containing the names of the German regions (Bundeslaender)
 region_code a factor containing the codes of the German regions (Bundeslaender)
 ind_code a factor containing the codes of the industries (WZ2008)
 ind_name a factor containing the names of the industries (WZ2008)
 firms a numeric vector containing the related number of firms
 emp_all a numeric vector containing the related number of employees (incl. self-employed)
 pop a numeric vector containing the related population
 area_sqkm a numeric vector containing the related region size (in sqkm)

Source

Compiled from:

Statistisches Bundesamt (2019): Tab. 11111-0001 - Gebietsflaeche: Bundeslaender, Stichtag.

Statistisches Bundesamt (2019): Tab. 12411-0010 - Bevoelkerung: Bundeslaender, Stichtag.

Statistisches Bundesamt (2019): Tab. 13311-0002 - Erwerbstaetige, Arbeitnehmer, Selbstaeendige und mithelfende Familienangehoerige (im Inland): Bundeslaender, Jahre, Wirtschaftszweige (Arbeitskreis "Erwerbstaetigenrechnung des Bundes und der Laender").

Statistisches Bundesamt (2019): Tab. 52111-0004 - Betriebe (Unternehmensregister-System): Bundeslaender, Jahre, Wirtschaftszweige (Abschnitte), Beschaeftigtengroessenklassen.

Examples

```

data (G.regions.industries)

lqs <- locq2(e_ij = G.regions.industries$emp_all,
G.regions.industries$ind_code, G.regions.industries$region_code,
LQ.output = "df")
# output as data frame

lqs_sort <- lqs[order(lqs$LQ, decreasing = TRUE),]
# Sort decreasing by size of LQ

lqs_sort[1:5,]

```

gifpro

Commercial area prognosis

Description

This function contains the basic GIFPRO model for commercial area prognosis (GIFPRO = Gewerbe- und Industrie- flaechenprognose)

Usage

```

gifpro(e_ij, a_i, sq_ij, rq_ij, ru_ij = NULL, ai_ij, time.base, tinterval = 1,
industry.names = NULL, output = "short")

```

Arguments

<code>e_ij</code>	a numeric vector with i values containing the current employment in i industries in region j
<code>a_i</code>	a numeric vector with i values containing the share of employees in industry i which is located in commercial areas
<code>sq_ij</code>	a numeric vector with i values containing the annual quote of resettled employees (<i>Neuansiedlungsquote</i> in German) in industry i , in percent
<code>rq_ij</code>	a numeric vector with i values containing the annual quote of relocated employees (<i>Verlagerungsquote</i> in German) in industry i , in percent
<code>ru_ij</code>	a numeric vector with i values containing the annual quote of employees in industry i which is located in reused commercial area (<i>Wiedernutzungsquote</i> in German), in percent (default: <code>ru_ij = NULL</code> , which represents a quote of 0 percent, meaning that no commercial area can be reused)
<code>ai_ij</code>	a numeric vector with i values containing the areal index (<i>Flaechenkennziffer</i> in German), representing the area requirement (e.g. in sqm) per employee in industry i
<code>time.base</code>	a single value representing the start time of the prognose (typically current year + 1)
<code>tinterval</code>	a single value representing the forecast horizon (length of time into the future for which the commercial area prognosis is done), in time units (e.g. <code>tinterval = 10 = 10 years</code>)
<code>industry.names</code>	a vector containing the industry names (e.g. from the relevant statistical classification of economic activities)
<code>output</code>	Type of output: <code>output = "short"</code> (default) shows the final number of relevant employment and commercial area requirement. If <code>output = "full"</code> , employment and commercial area are displayed for each time unit (year)

Details

In municipal land use planning (mostly in Germany), the future need of local commercial area (which is a type of land use, defined in official land-use plans) is mostly forecasted by models founded on the GIFPRO model (*Gewerbe- und Industrieflaechenbedarfsprognose*, prognosis of future demand of commercial area). GIFPRO is a demand-side model, which means predicting the demand of commercial area based on a prognosis of future employment in different industries (Bonny/Kahnert 2005). The key parameters of the model are the (assumed) shares of employees located in commercial areas (a_i), the (assumed) quotas of resettlement (sq_{ij}), relocation (rq_{ij}) and (sometimes) reuse (ru_{ij}) as well as the (assumed) area requirement per employee (ai_{ij}). Outgoing from current employment in i industries in region j , e_{ij} , the future employment is predicted based on the quotas mentioned above and, finally, multiplied by the industry-specific (and maybe region-specific) areal index. The GIFPRO model has been modified and extended several times, especially with respect to industry- and region-specific employment growth, quotas and areal indices (Deutsches Institut fuer Urbanistik 2010, Vallee et al. 2012).

Value

A list containing the following objects:

components	Matrices containing the single components (resettlement, relocation, reuse, relevant employment)
results	Matrices containing the final results per year and all over

Author(s)

Thomas Wieland

References

Bonny, H.-W./Kahnert, R. (2005): "Zur Ermittlung des Gewerbeflaechenbedarfs: Ein Vergleich zwischen einer Monitoring gestuetzten Prognose und einer analytischen Bestimmung". In: *Raumforschung und Raumordnung*, **63**, 3, p. 232-240.

Deutsches Institut fuer Urbanistik (ed.) (2010): "Stadtentwicklungskonzept Gewerbe fuer die Landeshauptstadt Potsdam". Berlin. https://www.potsdam.de/sites/default/files/documents/STEK_Gewerbe_Langfassung_2010.pdf (accessed October 13, 2017).

Vallee, D./Witte, A./Brandt, T./Bischof, T. (2012): "Bedarfsberechnung fuer die Darstellung von Allgemeinen Siedlungsbereichen (ASB) und Gewerbe- und Industrieansiedlungsbereichen (GIB) in Regionalplaenen". Im Auftrag der Staatskanzlei des Landes Nordrhein-Westfalen. Abschlussbericht Oktober 2012. Aachen.

See Also

[gifpro.tbs](#), [portfolio](#), [shift](#), [shiftd](#), [shifti](#)

Examples

```
# Data for the city Kempten (2012):
emp2012 <- c(7228, 12452, 11589)
sharesCA <- c(100, 40, 10)
rsquote <- c(0.3, 0.3, 0.3)
rlquote <- c(0.7, 0.7, 0.7)
arealindex <- c(148, 148, 148)
industries <- c("Manufacturing", "Wholesale and retail trade, Transportation
and storage, Information and communication", "Other services")

gifpro (e_ij = emp2012, a_i = sharesCA, sq_ij = rsquote,
rq_ij = rlquote, ai_ij = arealindex, time.base = 2012,
tinterval = 13, industry.names = industries, output = "short")
# short output

gifpro (e_ij = emp2012, a_i = sharesCA, sq_ij = rsquote,
rq_ij = rlquote, ai_ij = arealindex, time.base = 2012,
tinterval = 13, industry.names = industries, output = "full")
# full output

gifpro_results <- gifpro (e_ij = emp2012, a_i = sharesCA, sq_ij = rsquote,
rq_ij = rlquote, ai_ij = arealindex, time.base = 2012,
tinterval = 13, industry.names = industries, output = "short")
# saving results as gifpro object
```

```

gifpro_results$components
# single components

gifpro_results$results
# results (as shown in full output)

```

```
gifpro.tbs
```

Trend-based and location-specific commercial area prognosis

Description

This function contains the TBS-GIFPRO model for commercial area prognosis (TBS-GIFPRO = Trendbasierte und standortspezifische Gewerbe- und Industrieflaechenprognose; trend-based and location-specific commercial area prognosis)

Usage

```

gifpro.tbs(e_ij, a_i, sq_ij, rq_ij, ru_ij = NULL, ai_ij,
time.base, tinterval = 1, prog.func = rep("lin", nrow(e_ij)),
prog.plot = TRUE, plot.single = FALSE,
multiplot.col = NULL, multiplot.row = NULL,
industry.names = NULL, emp.only = FALSE, output = "short")

```

Arguments

<code>e_ij</code>	a numeric vector with i values containing the current employment in i industries in region j
<code>a_i</code>	a numeric vector with i values containing the share of employees in industry i which is located in commercial areas
<code>sq_ij</code>	a numeric vector with i values containing the annual quote of resettled employees (<i>Neuansiedlungsquote</i> in German) in industry i , in percent
<code>rq_ij</code>	a numeric vector with i values containing the annual quote of relocated employees (<i>Verlagerungsquote</i> in German) in industry i , in percent
<code>ru_ij</code>	a numeric vector with i values containing the annual quote of employees in industry i which is located in reused commercial area (<i>Wiedernutzungsquote</i> in German), in percent (default: <code>ru_ij = NULL</code> , which represents a quote of 0 percent, meaning that no commercial area can be reused)
<code>ai_ij</code>	a numeric vector with i values containing the areal index (<i>Flaechenkennziffer</i> in German), representing the area requirement (e.g. in sqm) per employee in industry i
<code>time.base</code>	a single value representing the start time of the prognose (typically current year + 1)
<code>tinterval</code>	a single value representing the forecast horizon (length of time into the future for which the commercial area prognosis is done), in time units (e.g. <code>tinterval = 10 = 10 years</code>)

prog.func	a vector containing the estimation function types for employment prognosis ("lin" for linear, "pow" for power, "exp" for exponential and "logi" for logistic function); must have the same length as e_ij and industry.names, respectively
prog.plot	Logical argument that indicates if the employment prognoses have to be plotted
plot.single	If prog.plot = TRUE: Logical argument that indicates if the plots are stored as single graphic devices or integrated in one plot
multiplot.col	No. of columns in plot
multiplot.row	No. of rows in plot
industry.names	a vector containing the industry names (e.g. from the relevant statistical classification of economic activities)
emp.only	Logical argument that indicates if the analysis only contains employment prognosis
output	Type of output: output = "short" (default) shows the final number of relevant employment and commercial area requirement. If output = "full", employment and commercial area are displayed for each time unit (year)

Details

In municipal land use planning (mostly in Germany), the future need of local commercial area (which is a type of land use, defined in official land-use plans) is mostly forecasted by models founded on the GIFPRO model (*Gewerbe- und Industrieflaechenbedarfsprognose*, prognosis of future demand of commercial area). GIFPRO is a demand-side model, which means predicting the demand of commercial area based on a prognosis of future employment in different industries (Bonny/Kahnert 2005). The key parameters of the model are the (assumed) shares of employees located in commercial areas (a_i), the (assumed) quotas of resettlement (sq_{ij}), relocation (rq_{ij}) and (sometimes) reuse (ru_{ij}) as well as the (assumed) area requirement per employee (ai_{ij}). Outgoing from current employment in i industries in region j , e_{ij} , the future employment is predicted based on the quotas mentioned above and, finally, multiplied by the industry-specific (and maybe region-specific) areal index. The GIFPRO model has been modified and extended several times, especially with respect to industry- and region-specific employment growth, quotas and areal indices (Deutsches Institut fuer Urbanistik 2010, Vallee et al. 2012).

This function contains the TBS-GIFPRO model for commercial area prognosis (TBS-GIFPRO = Trendbasierte und standortspezifische Gewerbe- und Industrieflaechenprognose; trend-based and location-specific commercial area prognosis) (Deutsches Institut fuer Urbanistik 2010).

Value

A list containing the following objects:

components	List with matrices containing the single components (resettlement, relocation, reuse, relevant employment)
results	List with matrices containing the final results per year and all over as well as the industry-specific forecast data

Author(s)

Thomas Wieland

References

Bonny, H.-W./Kahnert, R. (2005): “Zur Ermittlung des Gewerbeflaechenbedarfs: Ein Vergleich zwischen einer Monitoring gestuetzten Prognose und einer analytischen Bestimmung”. In: *Raumforschung und Raumordnung*, **63**, 3, p. 232-240.

Deutsches Institut fuer Urbanistik (ed.) (2010): “Stadtentwicklungskonzept Gewerbe fuer die Landeshauptstadt Potsdam”. Berlin. https://www.potsdam.de/sites/default/files/documents/STEK_Gewerbe_Langfassung_2010.pdf (accessed October 13, 2017).

Vallee, D./Witte, A./Brandt, T./Bischof, T. (2012): “Bedarfsberechnung fuer die Darstellung von Allgemeinen Siedlungsbereichen (ASB) und Gewerbe- und Industrieansiedlungsbereichen (GIB) in Regionalplaenen”. Im Auftrag der Staatskanzlei des Landes Nordrhein-Westfalen. Abschlussbericht Oktober 2012.

See Also

[gifpro](#), [portfolio](#), [shift](#), [shiftd](#), [shifti](#)

Examples

```
# Data for Goettingen:
data(Goettingen)

anteileGOE <- rep(100,15)
nvquote <- rep (0.3, 15)
vlquote <- rep (0.7, 15)

gifpro.tbs (e_ij = Goettingen[2:16,3:12],
a_i = anteileGOE, sq_ij = nvquote,
rq_ij = vlquote, tinterval = 12, prog.func =
rep("lin", nrow(Goettingen[2:16,3:12])),
ai_ij = 150, time.base = 2008, output = "full",
industry.names = Goettingen$WZ2008_Code[2:16],
prog.plot = TRUE, plot.single = FALSE)
```

gini

Gini coefficient

Description

Calculating the Gini coefficient of inequality (or concentration), standardized and non-standardized, and optionally plotting the Lorenz curve

Usage

```
gini(x, coefnorm = FALSE, weighting = NULL, na.rm = TRUE, lc = FALSE,
lcx = "% of objects", lcy = "% of regarded variable",
lctitle = "Lorenz curve", le.col = "blue", lc.col = "black",
lsize = 1, ltype = "solid",
```

```

bg.col = "gray95", bgrid = TRUE, bgrid.col = "white",
bgrid.size = 2, bgrid.type = "solid",
lcn = FALSE, lcn = FALSE, lcn.caption = NULL,
lcn.lab.x = 0, lcn.lab.y = 1, add.lcn = FALSE)

```

Arguments

x	A numeric vector (e.g. dataset of household income, sales turnover or supply)
coefnorm	logical argument that indicates if the function output is the non-standardized or the standardized Gini coefficient (default: coefnorm = FALSE, that means the non-standardized Gini coefficient is returned)
weighting	A numeric vector containing the weighting data (e.g. size of income classes when calculating a Gini coefficient for aggregated income data)
na.rm	logical argument that indicates whether NA values should be excluded before computing results
lc	logical argument that indicates if the Lorenz curve is plotted additionally (default: lc = FALSE, so no Lorenz curve is displayed)
lcx	if lc = TRUE (plot of Lorenz curve), lcx defines the x axis label
lcy	if lc = TRUE (plot of Lorenz curve), lcy defines the y axis label
lctitle	if lc = TRUE (plot of Lorenz curve), lctitle defines the overall title of the Lorenz curve plot
le.col	if lc = TRUE (plot of Lorenz curve), le.col defines the color of the diagonale (line of equality)
lc.col	if lc = TRUE (plot of Lorenz curve), lc.col defines the color of the Lorenz curve
lsize	if lc = TRUE (plot of Lorenz curve), lsize defines the size of the lines (default: 1)
ltype	if lc = TRUE (plot of Lorenz curve), ltype defines the type of the lines (default: "solid")
bg.col	if lc = TRUE (plot of Lorenz curve), bg.col defines the background color of the plot (default: "gray95")
bgrid	if lc = TRUE (plot of Lorenz curve), the logical argument bgrid defines if a grid is shown in the plot
bgrid.col	if lc = TRUE (plot of Lorenz curve) and bgrid = TRUE (background grid), bgrid.col defines the color of the background grid (default: "white")
bgrid.size	if lc = TRUE (plot of Lorenz curve) and bgrid = TRUE (background grid), bgrid.size defines the size of the background grid (default: 2)
bgrid.type	if lc = TRUE (plot of Lorenz curve) and bgrid = TRUE (background grid), bgrid.type defines the type of lines of the background grid (default: "solid")
lcn	if lc = TRUE (plot of Lorenz curve), the logical argument lcn defines if the non-standardized Gini coefficient is displayed in the Lorenz curve plot
lcn	if lc = TRUE (plot of Lorenz curve), the logical argument lcn defines if the standardized Gini coefficient is displayed in the Lorenz curve plot

l _{cg} .caption	if l _{cg} = TRUE (displaying the Gini coefficient in the plot), l _{cg} .caption specifies the caption above the coefficients
l _{cg} .lab.x	if l _{cg} = TRUE (displaying the Gini coefficient in the plot), l _{cg} .lab.x specifies the x coordinate of the label
l _{cg} .lab.y	if l _{cg} = TRUE (displaying the Gini coefficient in the plot), l _{cg} .lab.y specifies the y coordinate of the label
add.lc	if l _c = TRUE (plot of Lorenz curve), add.lc specifies if a new Lorenz curve is plotted (add.lc = "FALSE") or the plot is added to an existing Lorenz curve plot (add.lc = "TRUE")

Details

The *Gini coefficient* (Gini 1912) is a popular measure of statistical dispersion, especially used for analyzing inequality or concentration. The *Lorenz curve* (Lorenz 1905), though developed independently, can be regarded as a graphical representation of the degree of inequality/concentration calculated by the *Gini coefficient* (G) and can also be used for additional interpretations of it. In an economic-geographical context, these methods are frequently used to analyse the concentration/inequality of income or wealth within countries (Aoyama et al. 2011). Other areas of application are analyzing regional disparities (Lessmann 2005, Nakamura 2008) and concentration in markets (sales turnover of competing firms) which makes Gini and Lorenz part of economic statistics in general (Doersam 2004, Roberts 2014).

The *Gini coefficient* (G) varies between 0 (no inequality/concentration) and 1 (complete inequality/concentration). The *Lorenz curve* displays the deviations of the empirical distribution from a perfectly equal distribution as the difference between two graphs (the distribution curve and a diagonal line of perfect equality). This function calculates G and plots the *Lorenz curve* optionally. As there are several ways to calculate the *Gini coefficient*, this function uses the formula given in Doersam (2004). Because the maximum of G is not equal to 1, also a standardized coefficient (G^*) with a maximum equal to 1 can be calculated alternatively. If a Gini coefficient for aggregated data (e.g. income classes with averaged incomes) or the Gini coefficient has to be weighted, use a weighting vector (e.g. size of the income classes).

Value

A single numeric value of the *Gini coefficient* ($0 < G < 1$) or the *standardized Gini coefficient* ($0 < G^* < 1$) and, optionally, a plot of the *Lorenz curve*.

Author(s)

Thomas Wieland

References

- Aoyama, Y./Murphy, J. T./Hanson, S. (2011): "Key Concepts in Economic Geography". London : SAGE.
- Bahrenberg, G./Giese, E./Mevenkamp, N./Nipper, J. (2010): "Statistische Methoden in der Geographie. Band 1: Univariate und bivariate Statistik". Stuttgart: Borntraeger.
- Cerlani, L./Verme, P. (2012): "The origins of the Gini index: extracts from Variabilita e Mutabilita (1912) by Corrado Gini". In: *The Journal of Economic Inequality*, **10**, 3, p. 421-443.

- Doersam, P. (2004): "Wirtschaftsstatistik anschaulich dargestellt". Heidenau : PD-Verlag.
- Gini, C. (1912): "Variabilita e Mutabilita". Contributo allo Studio delle Distribuzioni e delle Relazioni Statistiche. Bologna : Cuppini.
- Lessmann, C. (2005): "Regionale Disparitaeten in Deutschland und ausgesuchten OECD-Staaten im Vergleich". *ifo Dresden berichtet*, **3/2005**. https://www.ifo.de/DocDL/ifodb_2005_3_25-33.pdf.
- Lorenz, M. O. (1905): "Methods of Measuring the Concentration of Wealth". In: *Publications of the American Statistical Association*, **9**, 70, p. 209-219.
- Nakamura, R. (2008): "Agglomeration Effects on Regional Economic Disparities: A Comparison between the UK and Japan". In: *Urban Studies*, **45**, 9, p. 1947-1971.
- Roberts, T. (2014): "When Bigger Is Better: A Critique of the Herfindahl-Hirschman Index's Use to Evaluate Mergers in Network Industries". In: *Pace Law Review*, **34**, 2, p. 894-946.

See Also

[cv](#), [gini.conc](#), [gini.spec](#), [herf](#), [hoover](#)

Examples

```
# Market concentration (example from Doersam 2004):
sales <- c(20,50,20,10)
# sales turnover of four car manufacturing companies
gini(sales, lc = TRUE, lcx = "percentage of companies", lcy = "percentage of sales",
lctitle = "Lorenz curve of sales", lcg = TRUE, lcgcn = TRUE)
# returns the non-standardized Gini coefficient (0.3) and
# plots the Lorenz curve with user-defined title and labels
gini(sales, coefnorm = TRUE)
# returns the standardized Gini coefficient (0.4)

# Income classes (example from Doersam 2004):
income <- c(500, 1500, 2500, 4000, 7500, 15000)
# average income of 6 income classes
sizeofclass <- c(1000, 1200, 1600, 400, 200, 600)
# size of income classes
gini(income, weighting = sizeofclass)
# returns the non-standardized Gini coefficient (0.5278)

# Market concentration in automotive industry
data(Automotive)
gini(Automotive$Turnover2008, lsize=1, lc=TRUE, le.col = "black",
lc.col = "orange", lcx = "Shares of companies", lcy = "Shares of turnover / cars",
lctitle = "Automotive industry: market concentration",
lcg = TRUE, lcgcn = TRUE, lcg.caption = "Turnover 2008:", lcg.lab.x = 0, lcg.lab.y = 1)
# Gini coefficient and Lorenz curve for turnover 2008
gini(Automotive$Turnover2013, lsize=1, lc = TRUE, add.lc = TRUE, lc.col = "red",
lcg = TRUE, lcgcn = TRUE, lcg.caption = "Turnover 2013:", lcg.lab.x = 0, lcg.lab.y = 0.85)
# Adding Gini coefficient and Lorenz curve for turnover 2013
gini(Automotive$Quantity2014_car, lsize=1, lc = TRUE, add.lc = TRUE, lc.col = "blue",
lcg = TRUE, lcgcn = TRUE, lcg.caption = "Cars 2014:", lcg.lab.x = 0, lcg.lab.y = 0.7)
# Adding Gini coefficient and Lorenz curve for cars 2014
```

```
# Regional disparities in Germany:
gdp <- c(460.69, 549.19, 124.16, 65.29, 31.59, 109.27, 263.44, 39.87, 258.53,
645.59, 131.95, 35.03, 112.66, 56.22, 85.61, 56.81)
# GDP of german regions (Bundeslaender) 2015 (in billion EUR)
gini(gdp)
# returns the non-standardized Gini coefficient (0.5009)
```

gini.conc

Gini coefficient of spatial industry concentration

Description

Calculating the Gini coefficient of spatial industry concentration based on regional industry data (normally employment data)

Usage

```
gini.conc(e_ij, e_j, lc = FALSE, lcx = "% of objects",
lcy = "% of regarded variable", lctitle = "Lorenz curve",
le.col = "blue", lc.col = "black", lsize = 1, ltype = "solid",
bg.col = "gray95", bgrid = TRUE, bgrid.col = "white",
bgrid.size = 2, bgrid.type = "solid", lcg = FALSE, lcg.n = FALSE,
lcg.caption = NULL, lcg.lab.x = 0, lcg.lab.y = 1,
add.lc = FALSE, plot.lc = TRUE)
```

Arguments

e_ij	a numeric vector with the employment of the industry i in region j
e_j	a numeric vector with the employment in region j
lc	logical argument that indicates if the Lorenz curve is plotted additionally (default: lc = FALSE, so no Lorenz curve is displayed)
lcx	if lc = TRUE (plot of Lorenz curve), lcx defines the x axis label
lcy	if lc = TRUE (plot of Lorenz curve), lcy defines the y axis label
lctitle	if lc = TRUE (plot of Lorenz curve), lctitle defines the overall title of the Lorenz curve plot
le.col	if lc = TRUE (plot of Lorenz curve), le.col defines the color of the diagonale (line of equality)
lc.col	if lc = TRUE (plot of Lorenz curve), lc.col defines the color of the Lorenz curve
lsize	if lc = TRUE (plot of Lorenz curve), lsize defines the size of the lines (default: 1)
ltype	if lc = TRUE (plot of Lorenz curve), ltype defines the type of the lines (default: "solid")
bg.col	if lc = TRUE (plot of Lorenz curve), bg.col defines the background color of the plot (default: "gray95")

<code>bgrid</code>	if <code>lc = TRUE</code> (plot of Lorenz curve), the logical argument <code>bgrid</code> defines if a grid is shown in the plot
<code>bgrid.col</code>	if <code>lc = TRUE</code> (plot of Lorenz curve) and <code>bgrid = TRUE</code> (background grid), <code>bgrid.col</code> defines the color of the background grid (default: "white")
<code>bgrid.size</code>	if <code>lc = TRUE</code> (plot of Lorenz curve) and <code>bgrid = TRUE</code> (background grid), <code>bgrid.size</code> defines the size of the background grid (default: 2)
<code>bgrid.type</code>	if <code>lc = TRUE</code> (plot of Lorenz curve) and <code>bgrid = TRUE</code> (background grid), <code>bgrid.type</code> defines the type of lines of the background grid (default: "solid")
<code>lcg</code>	if <code>lc = TRUE</code> (plot of Lorenz curve), the logical argument <code>lcg</code> defines if the non-standardized Gini coefficient is displayed in the Lorenz curve plot
<code>lcn</code>	if <code>lc = TRUE</code> (plot of Lorenz curve), the logical argument <code>lcn</code> defines if the standardized Gini coefficient is displayed in the Lorenz curve plot
<code>lcg.caption</code>	if <code>lcg = TRUE</code> (displaying the Gini coefficient in the plot), <code>lcg.caption</code> specifies the caption above the coefficients
<code>lcg.lab.x</code>	if <code>lcg = TRUE</code> (displaying the Gini coefficient in the plot), <code>lcg.lab.x</code> specifies the x coordinate of the label
<code>lcg.lab.y</code>	if <code>lcg = TRUE</code> (displaying the Gini coefficient in the plot), <code>lcg.lab.y</code> specifies the y coordinate of the label
<code>add.lc</code>	if <code>lc = TRUE</code> (plot of Lorenz curve), <code>add.lc</code> specifies if a new Lorenz curve is plotted (<code>add.lc = "FALSE"</code>) or the plot is added to an existing Lorenz curve plot (<code>add.lc = "TRUE"</code>)
<code>plot.lc</code>	logical argument that indicates if the Lorenz curve itself is plotted (if <code>plot.lc = FALSE</code> , only the line of equality is plotted)

Details

The *Gini coefficient of spatial industry concentration* (G_i) is a special spatial modification of the Gini coefficient of inequality (see the function `gini()`). It represents the rate of spatial concentration of the industry i referring to j regions (e.g. cities, counties, states). The coefficient G_i varies between 0 (perfect distribution, respectively no concentration) and 1 (complete concentration in one region). Optionally a Lorenz curve is plotted (if `lc = TRUE`).

Value

A single numeric value ($0 < G_i < 1$)

Author(s)

Thomas Wieland

References

Farhauer, O./Kroell, A. (2013): "Standorttheorien: Regional- und Stadtoekonomik in Theorie und Praxis". Wiesbaden : Springer.

Nakamura, R./Morrison Paul, C. J. (2009): "Measuring agglomeration". In: Capello, R./Nijkamp, P. (eds.): *Handbook of Regional Growth and Development Theories*. Cheltenham: Elgar. p. 305-328.

See Also

[gini](#), [gini.spec](#)

Examples

```
# Example from Farhauer/Kroell (2013):
E_ij <- c(500,500,1000,7000,1000)
# employment of the industry in five regions
E_j <- c(20000,15000,20000,40000,5000)
# employment in the five regions
gini.conc (E_ij, E_j)
# Returns the Gini coefficient of industry concentration (0.4068966)

data(G.regions.emp)
# Concentration of construction industry in Germany
# based on 16 German regions (Bundeslaender) for the year 2008
construction2008 <- G.regions.emp[(G.regions.emp$industry == "Baugewerbe (F)" |
G.regions.emp$industry == "Insgesamt") & G.regions.emp$year == "2008",]
# only data for construction industry (Baugewerbe) and all-over (Insgesamt)
# for the 16 German regions in the year 2008
construction2008 <- construction2008[construction2008$region != "Insgesamt",]
# delete all-over data for all industries
gini.conc(construction2008[construction2008$industry=="Baugewerbe (F)",]$emp,
construction2008[construction2008$industry=="Insgesamt",]$emp)

# Concentration of financial industry in Germany 2008 vs. 2014
# based on 16 German regions (Bundeslaender) for 2008 and 2014
finance2008 <- G.regions.emp[(G.regions.emp$industry ==
"Erbringung von Finanz- und Vers.leistungen (K)" |
G.regions.emp$industry == "Insgesamt") & G.regions.emp$year == "2008",]
finance2008 <- finance2008[finance2008$region != "Insgesamt",]
# delete all-over data for all industries
gini.conc(finance2008[finance2008$industry ==
"Erbringung von Finanz- und Vers.leistungen (K)",]$emp,
finance2008[finance2008$industry=="Insgesamt",]$emp)
finance2014 <- G.regions.emp[(G.regions.emp$industry ==
"Erbringung von Finanz- und Vers.leistungen (K)" | G.regions.emp$industry ==
"Insgesamt") & G.regions.emp$year == "2014",]
finance2014 <- finance2014[finance2014$region != "Insgesamt",]
# delete all-over data for all industries
gini.conc(finance2014[finance2014$industry ==
"Erbringung von Finanz- und Vers.leistungen (K)",]$emp,
finance2014[finance2014$industry=="Insgesamt",]$emp)
```

gini.spec

Gini coefficient of regional specialization

Description

Calculating the Gini coefficient of regional specialization based on regional industry data (normally employment data)

Usage

```
gini.spec(e_ij, e_i, lc = FALSE, lcx = "% of objects",
  lcy = "% of regarded variable", lctitle = "Lorenz curve",
  le.col = "blue", lc.col = "black", lsize = 1, ltype = "solid",
  bg.col = "gray95", bgrid = TRUE, bgrid.col = "white",
  bgrid.size = 2, bgrid.type = "solid", lcg = FALSE, lcg.n = FALSE,
  lcg.caption = NULL, lcg.lab.x = 0, lcg.lab.y = 1,
  add.lc = FALSE, plot.lc = TRUE)
```

Arguments

<code>e_ij</code>	a numeric vector with the employment of the industries i in region j
<code>e_i</code>	a numeric vector with the employment in the industries i
<code>lc</code>	logical argument that indicates if the Lorenz curve is plotted additionally (default: <code>lc = FALSE</code> , so no Lorenz curve is displayed)
<code>lcx</code>	if <code>lc = TRUE</code> (plot of Lorenz curve), <code>lcx</code> defines the x axis label
<code>lcy</code>	if <code>lc = TRUE</code> (plot of Lorenz curve), <code>lcy</code> defines the y axis label
<code>lctitle</code>	if <code>lc = TRUE</code> (plot of Lorenz curve), <code>lctitle</code> defines the overall title of the Lorenz curve plot
<code>le.col</code>	if <code>lc = TRUE</code> (plot of Lorenz curve), <code>le.col</code> defines the color of the diagonale (line of equality)
<code>lc.col</code>	if <code>lc = TRUE</code> (plot of Lorenz curve), <code>lc.col</code> defines the color of the Lorenz curve
<code>lsize</code>	if <code>lc = TRUE</code> (plot of Lorenz curve), <code>lsize</code> defines the size of the lines (default: 1)
<code>ltype</code>	if <code>lc = TRUE</code> (plot of Lorenz curve), <code>ltype</code> defines the type of the lines (default: "solid")
<code>bg.col</code>	if <code>lc = TRUE</code> (plot of Lorenz curve), <code>bg.col</code> defines the background color of the plot (default: "gray95")
<code>bgrid</code>	if <code>lc = TRUE</code> (plot of Lorenz curve), the logical argument <code>bgrid</code> defines if a grid is shown in the plot
<code>bgrid.col</code>	if <code>lc = TRUE</code> (plot of Lorenz curve) and <code>bgrid = TRUE</code> (background grid), <code>bgrid.col</code> defines the color of the background grid (default: "white")
<code>bgrid.size</code>	if <code>lc = TRUE</code> (plot of Lorenz curve) and <code>bgrid = TRUE</code> (background grid), <code>bgrid.size</code> defines the size of the background grid (default: 2)
<code>bgrid.type</code>	if <code>lc = TRUE</code> (plot of Lorenz curve) and <code>bgrid = TRUE</code> (background grid), <code>bgrid.type</code> defines the type of lines of the background grid (default: "solid")
<code>lcg</code>	if <code>lc = TRUE</code> (plot of Lorenz curve), the logical argument <code>lcg</code> defines if the non-standardized Gini coefficient is displayed in the Lorenz curve plot
<code>lcg.n</code>	if <code>lc = TRUE</code> (plot of Lorenz curve), the logical argument <code>lcg.n</code> defines if the standardized Gini coefficient is displayed in the Lorenz curve plot
<code>lcg.caption</code>	if <code>lcg = TRUE</code> (displaying the Gini coefficient in the plot), <code>lcg.caption</code> specifies the caption above the coefficients

lcg.lab.x	if lcg = TRUE (displaying the Gini coefficient in the plot), lcg.lab.x specifies the x coordinate of the label
lcg.lab.y	if lcg = TRUE (displaying the Gini coefficient in the plot), lcg.lab.y specifies the y coordinate of the label
add.lc	if lc = TRUE (plot of Lorenz curve), add.lc specifies if a new Lorenz curve is plotted (add.lc = "FALSE") or the plot is added to an existing Lorenz curve plot (add.lc = "TRUE")
plot.lc	logical argument that indicates if the Lorenz curve itself is plotted (if plot.lc = FALSE, only the line of equality is plotted)

Details

The *Gini coefficient of regional specialization* (G_j) is a special spatial modification of the *Gini coefficient* of inequality (see the function `gini()`). It represents the degree of regional specialization of the region j referring to i industries. The coefficient G_j varies between 0 (no specialization) and 1 (complete specialization). Optionally a Lorenz curve is plotted (if `lc = TRUE`).

Value

A single numeric value ($0 < G_j < 1$)

Author(s)

Thomas Wieland

References

Farhauer, O./Kroell, A. (2013): "Standorttheorien: Regional- und Stadtoekonomik in Theorie und Praxis". Wiesbaden : Springer.

Nakamura, R./Morrison Paul, C. J. (2009): "Measuring agglomeration". In: Capello, R./Nijkamp, P. (eds.): *Handbook of Regional Growth and Development Theories*. Cheltenham: Elgar. p. 305-328.

See Also

[gini](#), [gini.conc](#)

Examples

```
# Example from Farhauer/Kroell (2013):
E_ij <- c(700,600,500,10000,40000)
# employment of five industries in the region
E_i <- c(30000,15000,10000,60000,50000)
# over-all employment in the five industries
gini.spec(E_ij, E_i)
# Returns the Gini coefficient of regional specialization (0.6222222)

# Example Freiburg
data(Freiburg)
```

```

# Loads the data
E_ij <- Freiburg$e_Freiburg2014
# industry-specific employment in Freiburg 2014
E_i <- Freiburg$e_Germany2014
# industry-specific employment in Germany 2014
gini.spec (E_ij, E_i)
# Returns the Gini coefficient of regional specialization (0.2089009)

# Example Goettingen
data(Goettingen)
# Loads the data
gini.spec(Goettingen$Goettingen2017[2:16], Goettingen$BRD2017[2:16])
# Returns the Gini coefficient of regional specialization 2017 (0.359852)

```

gini2	<i>Gini coefficient</i>
-------	-------------------------

Description

Calculating the Gini coefficient of inequality (or concentration), standardized and non-standardized, and optionally plotting the Lorenz curve

Usage

```
gini2(x, weighting = NULL, coefnorm = FALSE, na.rm = TRUE)
```

Arguments

x	A numeric vector (e.g. dataset of regional incomes)
weighting	A numeric vector containing the weighting data (e.g. regional population)
coefnorm	logical argument that indicates if the function output is the non-standardized or the standardized Gini coefficient (default: coefnorm = FALSE, which means the non-standardized Gini coefficient is returned)
na.rm	logical argument that indicates whether NA values should be excluded before computing results

Details

The *Gini coefficient* (Gini 1912) is a popular measure of statistical dispersion, especially used for analyzing inequality or concentration. In an economic-geographical context, the Gini coefficient is frequently used to analyse the concentration/inequality of income or wealth within countries (Aoyama et al. 2011). Other areas of application are analyzing regional disparities (Lessmann 2005, Nakamura 2008) and concentration in markets (sales turnover of competing firms).

The *Gini coefficient* (G) varies between 0 (no inequality/concentration) and 1 (complete inequality/concentration). This function calculates G . As there are several ways to calculate the *Gini coefficient*, this function uses the formula given in Doersam (2004). Because the maximum of G is not equal to 1, also a standardized coefficient (G^*) with a maximum equal to 1 can be calculated alternatively. If a Gini coefficient for aggregated data (e.g. income classes with averaged incomes) or the Gini coefficient has to be weighted, use a weighting vector (e.g. size of the income classes).

Value

A single numeric value of the *Gini coefficient* ($0 < G < 1$) or the *standardized Gini coefficient* ($0 < G^* < 1$) and, optionally, a plot of the *Lorenz curve*.

Author(s)

Thomas Wieland

References

- Aoyama, Y./Murphy, J. T./Hanson, S. (2011): “Key Concepts in Economic Geography”. London : SAGE.
- Bahrenberg, G./Giese, E./Mevenkamp, N./Nipper, J. (2010): “Statistische Methoden in der Geographie. Band 1: Univariate und bivariate Statistik”. Stuttgart: Borntraeger.
- Cerlani, L./Verme, P. (2012): “The origins of the Gini index: extracts from Variabilita e Mutabilita (1912) by Corrado Gini”. In: *The Journal of Economic Inequality*, **10**, 3, p. 421-443.
- Doersam, P. (2004): “Wirtschaftsstatistik anschaulich dargestellt”. Heidenau : PD-Verlag.
- Gini, C. (1912): “Variabilita e Mutabilita”. Contributo allo Studio delle Distribuzioni e delle Relazioni Statistiche. Bologna : Cuppini.
- Lessmann, C. (2005): “Regionale Disparitaeten in Deutschland und ausgesuchten OECD-Staaten im Vergleich”. *ifo Dresden berichtet*, **3/2005**. https://www.ifo.de/DocDL/ifodb_2005_3_25-33.pdf.
- Lorenz, M. O. (1905): “Methods of Measuring the Concentration of Wealth”. In: *Publications of the American Statistical Association*, **9**, 70, p. 209-219.
- Nakamura, R. (2008): “Agglomeration Effects on Regional Economic Disparities: A Comparison between the UK and Japan”. In: *Urban Studies*, **45**, 9, p. 1947-1971.
- Roberts, T. (2014): “When Bigger Is Better: A Critique of the Herfindahl-Hirschman Index’s Use to Evaluate Mergers in Network Industries”. In: *Pace Law Review*, **34**, 2, p. 894-946.

See Also

[cv](#), [gini.conc](#), [gini.spec](#), [herf](#), [hoover](#)

Examples

```
# Market concentration (example from Doersam 2004):
sales <- c(20,50,20,10)
# sales turnover of four car manufacturing companies
gini (sales, lc = TRUE, lcx = "percentage of companies", lcy = "percentage of sales",
lctitle = "Lorenz curve of sales", lcg = TRUE, lcg = TRUE)
# returns the non-standardized Gini coefficient (0.3) and
# plots the Lorenz curve with user-defined title and labels
gini (sales, coefnorm = TRUE)
# returns the standardized Gini coefficient (0.4)

# Income classes (example from Doersam 2004):
income <- c(500, 1500, 2500, 4000, 7500, 15000)
```

```

# average income of 6 income classes
sizeofclass <- c(1000, 1200, 1600, 400, 200, 600)
# size of income classes
gini (income, weighting = sizeofclass)
# returns the non-standardized Gini coefficient (0.5278)

# Market concentration in automotive industry
data(Automotive)
gini(Automotive$Turnover2008, lsize=1, lc=TRUE, le.col = "black",
lc.col = "orange", lcx = "Shares of companies", lcy = "Shares of turnover / cars",
lctitle = "Automotive industry: market concentration",
lcg = TRUE, lcg = TRUE, lcg.caption = "Turnover 2008:", lcg.lab.x = 0, lcg.lab.y = 1)
# Gini coefficient and Lorenz curve for turnover 2008
gini(Automotive$Turnover2013, lsize=1, lc = TRUE, add.lc = TRUE, lc.col = "red",
lcg = TRUE, lcg = TRUE, lcg.caption = "Turnover 2013:", lcg.lab.x = 0, lcg.lab.y = 0.85)
# Adding Gini coefficient and Lorenz curve for turnover 2013
gini(Automotive$Quantity2014_car, lsize=1, lc = TRUE, add.lc = TRUE, lc.col = "blue",
lcg = TRUE, lcg = TRUE, lcg.caption = "Cars 2014:", lcg.lab.x = 0, lcg.lab.y = 0.7)
# Adding Gini coefficient and Lorenz curve for cars 2014

# Regional disparities in Germany:
gdp <- c(460.69, 549.19, 124.16, 65.29, 31.59, 109.27, 263.44, 39.87, 258.53,
645.59, 131.95, 35.03, 112.66, 56.22, 85.61, 56.81)
# GDP of german regions (Bundeslaender) 2015 (in billion EUR)
gini(gdp)
# returns the non-standardized Gini coefficient (0.5009)

```

Goettingen

Employment data for Goettingen and Germany 2008-2017

Description

This dataset contains the employees in 15 economic sections (German Classification of Economic Activities WZ2008) for the city Goettingen and Germany regarding the years 2008-2017 (date: 30 June each year).

Usage

```
data("Goettingen")
```

Format

A data frame with 16 observations on the following 22 variables.

WZ2008_Code a factor containing the code of the industry (15 economic sections from the German Classification of Economic Activities WZ2008 + total employees), in German language

WZ2008_Name a factor containing the name of the industry (15 economic sections from the German Classification of Economic Activities WZ2008 + total employees), in German language

Goettingen2008 industry employees in the city of Goettingen 2008

Goettingen2009 industry employees in the city of Goettingen 2009
 Goettingen2010 industry employees in the city of Goettingen 2010
 Goettingen2011 industry employees in the city of Goettingen 2011
 Goettingen2012 industry employees in the city of Goettingen 2012
 Goettingen2013 industry employees in the city of Goettingen 2013
 Goettingen2014 industry employees in the city of Goettingen 2014
 Goettingen2015 industry employees in the city of Goettingen 2015
 Goettingen2016 industry employees in the city of Goettingen 2016
 Goettingen2017 industry employees in the city of Goettingen 2017
 BRD2008 industry employees in Germany 2008
 BRD2009 industry employees in Germany 2009
 BRD2010 industry employees in Germany 2010
 BRD2011 industry employees in Germany 2011
 BRD2012 industry employees in Germany 2012
 BRD2013 industry employees in Germany 2013
 BRD2014 industry employees in Germany 2014
 BRD2015 industry employees in Germany 2015
 BRD2016 industry employees in Germany 2016
 BRD2017 industry employees in Germany 2017

Source

Bundesagentur fuer Arbeit (2018): "Beschaeftigungsstatistik, Beschaeftigte nach Wirtschaftszweigen (WZ 2008) (Zeitreihe Quartalszahlen) in Deutschland". <https://statistik.arbeitsagentur.de/DE/Navigation/Statistiken/Fachstatistiken/Beschaeftigung/Beschaeftigung-Nav.html> (accessed October 10, 2018). Own postprocessing (filtering and aggregation).

Stadt Goettingen - Referat Statistik und Wahlen (2008): "Stadt Goettingen: Beschaeftigte nach Wirtschaftsbereichen und Wirtschaftsabschnitten 1980 bis 2018. Table: IS 071.20". <https://duva-stg-extern.kdgoe.de/Informationsportal/Dateien/071.20-2018.pdf> (accessed November 21, 2019).

References

Bundesagentur fuer Arbeit (2018): "Beschaeftigungsstatistik, Beschaeftigte nach Wirtschaftszweigen (WZ 2008) (Zeitreihe Quartalszahlen) in Deutschland". <https://statistik.arbeitsagentur.de/DE/Navigation/Statistiken/Fachstatistiken/Beschaeftigung/Beschaeftigung-Nav.html> (accessed October 10, 2018).

Federal Statistical Office Germany (2008): "Classification of Economic Activities, Edition 2008 (WZ 2008)". <https://www.klassifikationsserver.de/klassService/jsp/common/url.jsf?variant=wz2008&lang=EN> (accessed June 07, 2019).

Stadt Goettingen - Referat Statistik und Wahlen (2008): "Stadt Goettingen: Beschaeftigte nach Wirtschaftsbereichen und Wirtschaftsabschnitten 1980 bis 2018. Table: IS 071.20". <https://duva-stg-extern.kdgoe.de/Informationsportal/Dateien/071.20-2018.pdf> (accessed November 21, 2019).

Examples

```

data(Goettingen)

# Location quotients for Goettingen 2017:
locq (Goettingen$Goettingen2017[2:16], Goettingen$Goettingen2017[1],
      Goettingen$BRD2017[2:16], Goettingen$BRD2017[1])

# Gini coefficient of regional specialization 2017:
gini.spec(Goettingen$Goettingen2017[2:16], Goettingen$BRD2017[2:16])

# Krugman coefficient of regional specialization 2017:
krugman.spec(Goettingen$Goettingen2017[2:16], Goettingen$BRD2017[2:16])

```

GoettingenHealth1 *Healthcare providers in South Lower Saxony*

Description

Dataset with healthcare providers (general practitioners, psychotherapists, pharmacies) in two German counties (Goettingen and Northeim)

Usage

```
data("GoettingenHealth1")
```

Format

A data frame with 617 observations on the following 5 variables.

location a numeric vector with unique IDs of the healthcare providers

lat Latitude

lon Longitude

type Type of healthcare provider: general practitioners (phyh_gen), psychotherapists (psych) or pharmacies (pharm)

district a numeric vector containing the IDs of the district the specific provider is located in

Source

Wieland T./Dittrich, C. (2016): "Bestands- und Erreichbarkeitsanalyse regionaler Gesundheitseinrichtungen in der Gesundheitsregion Goettingen". Research report, Georg-August-Universitaet Goettingen, Geographisches Institut, Abteilung Humangeographie. <http://webdoc.sub.gwdg.de/pub/mon/2016/3-wieland.pdf>.

References

Wieland T./Dittrich, C. (2016): "Bestands- und Erreichbarkeitsanalyse regionaler Gesundheitseinrichtungen in der Gesundheitsregion Goettingen". Research report, Georg-August-Universitaet Goettingen, Geographisches Institut, Abteilung Humangeographie. <http://webdoc.sub.gwdg.de/pub/mon/2016/3-wieland.pdf>.

Examples

```
## Not run:
data(GoettingenHealth1)
# general practitioners, psychotherapists and pharmacies

area_goe <- 1753000000
# area of Landkreis Goettingen (sqm)
area_nom <- 1267000000
# area of Landkreis Northeim (sqm)
area_gn <- area_goe+area_nom
sqrt(area_gn/pi)

# this takes some seconds
ripley(GoettingenHealth1[GoettingenHealth1$type == "phys_gen",],
"location", "lat", "lon", area = area_gn, t.max = 30000, t.sep = 300)

ripley(GoettingenHealth1[GoettingenHealth1$type == "pharm",],
"location", "lat", "lon", area = area_gn, t.max = 30000, t.sep = 300)

ripley(GoettingenHealth1[GoettingenHealth1$type == "psych",],
"location", "lat", "lon", area = area_gn, t.max = 30000, t.sep = 300)

## End(Not run)
```

GoettingenHealth2 *Healthcare provision in South Lower Saxony*

Description

Dataset with districts in two German counties (Goettingen and Northeim) and the corresponding healthcare providers (general practitioners, psychotherapists, pharmacies) and population size

Usage

```
data("GoettingenHealth2")
```

Format

A data frame with 420 observations on the following 7 variables.

district a numeric vector containing the IDs of the district

pop no. of inhabitants

lat Latitude

lon Longitude

phys_gen no. of general practitioners

psych no. of psychotherapists

pharm no. of pharmacies

Source

Wieland T./Dittrich, C. (2016): “Bestands- und Erreichbarkeitsanalyse regionaler Gesundheitseinrichtungen in der Gesundheitsregion Goettingen”. Research report, Georg-August-Universitaet Goettingen, Geographisches Institut, Abteilung Humangeographie. <http://webdoc.sub.gwdg.de/pub/mon/2016/3-wieland.pdf>.

References

Wieland T./Dittrich, C. (2016): “Bestands- und Erreichbarkeitsanalyse regionaler Gesundheitseinrichtungen in der Gesundheitsregion Goettingen”. Research report, Georg-August-Universitaet Goettingen, Geographisches Institut, Abteilung Humangeographie. <http://webdoc.sub.gwdg.de/pub/mon/2016/3-wieland.pdf>.

Examples

```
data(GoettingenHealth2)
# districts with healthcare providers and population size

williamson((GoettingenHealth2$phys_gen/GoettingenHealth2$pop),
GoettingenHealth2$pop)
```

growth	<i>Growth rates</i>
--------	---------------------

Description

This function calculates the growth from two input numeric vectors

Usage

```
growth(val1, val2, growth.type = "growth",
output = "rate", rate.perc = FALSE, log.rate = FALSE,
factor.mean = "mean", time.periods = NULL)
```

Arguments

<code>val1</code>	First numeric vector (e.g. employment at time t)
<code>val2</code>	Second numeric vector (e.g. employment at time t) or data frame for times $t+1$, $t+2$, $t+3$, ..., $t+n$
<code>growth.type</code>	Type of growth value that has to be calculated (absolute values or growth rate)
<code>output</code>	Type of output in the case of several years: growth rate (default: <code>output = "rate"</code>) or annual values (<code>output = "annual"</code>)
<code>rate.perc</code>	Logical argument that indicates whether growth rates are expressed in percent or not
<code>log.rate</code>	Logical argument that indicates whether growth rates are logged or not
<code>factor.mean</code>	If growth factors are returned: arithmetic mean (<code>factor.mean = "mean"</code>) or geometric mean (<code>factor.mean = "geom"</code>)
<code>time.periods</code>	No. of regarded time periods (for average growth rates)

Value

A numeric vector containing the growth rates in the same order as stated

Author(s)

Thomas Wieland

Examples

```
# Example from Farhauer/Kroell (2013):
region_A_t <- c(90,20,10,60)
region_A_t1 <- c(100,40,10,55)
# data for region A (time t and t+1)
nation_X_t <- c(400,150,150,400)
nation_X_t1 <- c(440,210,135,480)
# data for the national economy (time t and t+1)
growth(region_A_t, region_A_t1)

data(Freiburg)
# Loads the data
growth(Freiburg$e_Freiburg2008, Freiburg$e_Freiburg2014, growth.type = "rate")
# Industry-specific growth rates for Freiburg 2008 to 2014
```

hansen

Hansen accessibility

Description

Calculating the Hansen accessibility for given origins and destinations

Usage

```
hansen(od_dataset, origins, destinations, attrac, dist, gamma = 1, lambda = -2,
atp = "pow", dtp = "pow", gamma2 = NULL, lambda2 = NULL, dist_const = 0,
dist_max = NULL, extract_local = FALSE, accnorm = FALSE, check_df = TRUE,
print.results = TRUE)
```

Arguments

od_dataset	an interaction matrix which is a data.frame containing the origins, destinations, the distances between them and a size variable for the opportunities of the destinations
origins	the column in the interaction matrix od_dataset containing the origins
destinations	the column in the interaction matrix od_dataset containing the destinations
attrac	the column in the interaction matrix od_dataset containing the "attractivity" variable of the destinations (e.g. no. of opportunities)

dist	the column in the interaction matrix <code>od_dataset</code> containing the transport costs (e.g. travelling time, distance)
gamma	a single numeric value for the exponential weighting (γ) of size (default: 1)
lambda	a single numeric value for the exponential weighting (λ) of distance (transport costs, default: -2)
atype	Type of attractivity weighting function: <code>atype = "pow"</code> (power function), <code>atype = "exp"</code> (exponential function) or <code>atype = "logistic"</code> (default: <code>atype = "pow"</code>)
dtype	Type of distance weighting function: <code>dtype = "pow"</code> (power function), <code>dtype = "exp"</code> (exponential function) or <code>dtype = "logistic"</code> (default: <code>dtype = "pow"</code>)
gamma2	if <code>atype = "logistic"</code> a second γ parameter is needed
lambda2	if <code>dtype = "logistic"</code> a second λ parameter is needed
dist_const	a numeric value of a constant to be added to the transport costs (e.g. 1)
dist_max	a numeric value of a maximal value of transport costs for the opportunities to be recognized
extract_local	logical argument that indicates if the start points should be included in the analysis or not (if $i = j$). Default value: <code>extract_local = FALSE</code>
accnorm	logical argument that indicates if the Hansen accessibility should be standardized
check_df	logical argument that indicates if the given dataset is checked for correct input, only for internal use, should not be deselected (default: TRUE)
print.results	logical argument that indicates if the results are shown (default: TRUE)

Details

Accessibility and the inhibiting effect of transport costs on spatial interactions belong to the key concepts of economic geography (Aoyama et al. 2011). The *Hansen accessibility* (Hansen 1959) can be regarded as a *potential model of spatial interaction* that describes accessibility as the sum of all opportunities O in the regions j , O_j , weighted by distance or other types of transport costs from the origins, i , to them, d_{ij} : $A_i = \sum_j O_j f(d_{ij})$. The distance/travel time is weighted by a distance decay function ($f(d_{ij})$) to reflect the disutility (opportunity costs) of distance. From a microeconomic perspective, the accessibility of a region or zone can be seen as the sum of all utilities of every opportunity outgoing from given starting points, given an utility function containing the opportunities (utility) and transport costs (disutility) (Orpana/Lampinen 2003). As the accessibility model originally comes from urban land use theory, it can also be used to model spatial concentration/agglomeration, e.g. to quantify the rate of agglomeration of retail locations (Orpana/Lampinen 2003, Wieland 2015).

Originally the weighting function of distance is not explicitly stated and the "attractivities" (e.g. size of the activity at the destinations) is not weighted. These specifications are relaxed in this function, so both variables can be weighted by a power, exponential or logistic function. If `accnorm = TRUE`, the *Hansen accessibility* is standardized by weighting the non-standardized values by the sum of all opportunities without regarding transport costs; the standardized *Hansen accessibility* has a range between 0 and 1.

Value

A list containing the following objects:

origins	A data frame containing the origins
accessibility	A data frame containing the calculated accessibility values (optional: standardized accessibilities)

Author(s)

Thomas Wieland

References

- Aoyama, Y./Murphy, J. T./Hanson, S. (2011): “Key Concepts in Economic Geography”. London : SAGE.
- Hansen, W. G. (1959): “How Accessibility Shapes Land Use”. In: *Journal of the American Institute of Planners*, **25**, 2, p. 73-76.
- Orpana, T./Lampinen, J. (2003): “Building spatial choice models from aggregate data”. In: *Journal of Regional Science*, **43**, 2, p. 319-347.
- Wieland, T. (2015): “Räumliches Einkaufsverhalten und Standortpolitik im Einzelhandel unter Berücksichtigung von Agglomerationseffekten. Theoretische Erklärungsansätze, modellanalytische Zugänge und eine empirisch-ökonomische Marktgebietsanalyse anhand eines Fallbeispiels aus dem ländlichen Raum Ostwestfalens/Süd-niedersachsens”. *Geographische Handlungsforschung*, **23**. 289 pages. Mannheim : MetaGIS.

See Also

[converse](#), [dist.calc](#), [dist.mat](#), [dist.buf](#), [huff](#), [reilly](#)

Examples

```
# Example from Levy/Weitz (2009):
# Data for the existing and the new location
locations <- c("Existing Store", "New Store")
S_j <- c(5000, 10000)
location_data <- data.frame(locations, S_j)
# Data for the two communities (Rock Creek and Oak Hammock)
communities <- c("Rock Creek", "Oak Hammock")
C_i <- c(5000000, 3000000)
community_data <- data.frame(communities, C_i)
# Combining location and submarket data in the interaction matrix
interactionmatrix <- merge (community_data, location_data)
# Adding driving time:
interactionmatrix[1,5] <- 10
interactionmatrix[2,5] <- 5
interactionmatrix[3,5] <- 5
interactionmatrix[4,5] <- 15
colnames(interactionmatrix) <- c("communities", "C_i", "locations", "S_j", "d_ij")
shoppingcenters1 <- interactionmatrix
```

```

huff_shares <- huff(shoppingcenters1, "communities", "locations", "S_j", "d_ij")
# Market shares of the new location:
huff_shares$ijmatrix[huff_shares$ijmatrix$locations == "New Store",]
# Hansen accessibility for Oak Hammock and Rock Creek:
# hansen (huff_shares$ijmatrix, "communities", "locations", "S_j", "d_ij")

```

herf	<i>Herfindahl-Hirschman coefficient</i>
------	---

Description

Calculating the Herfindahl-Hirschman coefficient of concentration, standardized and non-standardized

Usage

```
herf(x, coefnorm = FALSE, output = "HHI", na.rm = TRUE)
```

Arguments

x	A numeric vector (e.g. dataset of sales turnover or size of firms)
coefnorm	logical argument that indicates if the function output is the non-standardized or the standardized Herfindahl-Hirschman coefficient (default: <code>coefnorm = FALSE</code> , that means the non-standardized Herfindahl-Hirschman coefficient is returned)
output	argument to state the output. If <code>output = "HHI"</code> (default), the Herfindahl-Hirschman coefficient is returned (standardized or non-standardized). If <code>output = "eq"</code> , the Herfindahl-Hirschman coefficient equivalent number is returned
na.rm	logical argument that indicates whether NA values should be excluded before computing results

Details

The *Herfindahl-Hirschman coefficient* is a popular measure of statistical dispersion, especially used for analyzing concentration in markets, regarding sales turnovers or sizes of n competing firms in an industry. This indicator is especially used as a measure of market power and distortions of competition in the governmental competition policy (Roberts 2014). But the coefficient is also utilized as a measure of geographic concentration of industries (Lessmann 2005, Nakamura/Morrison Paul 2009).

The coefficient (HHI) varies between $\frac{1}{n}$ (parity resp. no concentration) and 1 (complete concentration). Because the minimum of HHI is not equal to 0, also a standardized coefficient (HHI^*) with a minimum equal to 0 can be calculated alternatively. The *equivalent number* (which is the inverse of the *Herfindahl-Hirschman coefficient*) reflects the theoretical number of economic objects (normally firms) where a calculated coefficient is $\frac{1}{n}$, which means parity (Doersam 2004). In a regional context, the inverse of HHI is also used as a measure of diversity (Duranton/Puga 2000).

Value

A single numeric value of the *Herfindahl-Hirschman coefficient* ($\frac{1}{n} < HHI < 1$) or the *standardized Herfindahl-Hirschman coefficient* ($0 < HHI^* < 1$) or the *Herfindahl-Hirschman coefficient equivalent number* ($H_{eq} \geq 1$).

Author(s)

Thomas Wieland

References

- Doersam, P. (2004): “Wirtschaftsstatistik anschaulich dargestellt”. Heidenau : PD-Verlag.
- Duranton, G./Puga, D. (2000): “Diversity and Specialisation in Cities: Why, Where and When Does it Matter?”. In: *Urban Studies*, **37**, 3, p. 533-555.
- Lessmann, C. (2005): “Regionale Disparitäten in Deutschland und ausgesuchten OECD-Staaten im Vergleich”. *ifo Dresden berichtet*, **3/2005**. https://www.ifo.de/DocDL/ifodb_2005_3_25-33.pdf.
- Nakamura, R./Morrison Paul, C. J. (2009): “Measuring agglomeration”. In: Capello, R./Nijkamp, P. (eds.): *Handbook of Regional Growth and Development Theories*. Cheltenham: Elgar. p. 305-328.
- Roberts, T. (2014): “When Bigger Is Better: A Critique of the Herfindahl-Hirschman Index’s Use to Evaluate Mergers in Network Industries”. In: *Pace Law Review*, **34**, 2, p. 894-946.

See Also

[cv](#), [gini](#)

Examples

```
# Example from Doersam (2004):
sales <- c(20,50,20,10)
# sales turnover of four car manufacturing companies
herf(sales)
# returns the non-standardized HHI (0.34)
herf(sales, coefnorm=TRUE)
# returns the standardized HHI (0.12)
herf(sales, output = "eq")
# returns the HHI equivalent number (2.94)

# Regional disparities in Germany:
gdp <- c(460.69, 549.19, 124.16, 65.29, 31.59, 109.27, 263.44, 39.87, 258.53,
645.59, 131.95, 35.03, 112.66, 56.22, 85.61, 56.81)
# GDP of german regions 2015 (in billion EUR)
herf(gdp)
# returns the HHI (0.125)
```

hoover	<i>Hoover Concentration Index</i>
--------	-----------------------------------

Description

Calculating the Hoover Concentration Index with respect to regional income (e.g. GDP) and population

Usage

```
hoover(x, ref = NULL, weighting = NULL, output = "HC", na.rm = TRUE)
```

Arguments

x	A numeric vector (dataset of regional income, e.g. GDP)
ref	A numeric vector containing the reference distribution for the Hoover Index, e.g. population. If reg = NULL, the reference distribution is set to $1/n$
weighting	A numeric containing the weightings for the Hoover Index, e.g. population
output	Default option is the output of the Hoover Index. If output = "data", the corresponding data table is returned instead
na.rm	logical argument that indicates whether NA values should be excluded before computing results

Details

The *Hoover Concentration Index (CI)* measures the economic concentration of income across space by comparing the share of income (e.g. GDP - Gross Domestic Product) with the share of population. The index varies between 0 (no inequality/concentration) and 1 (complete inequality/concentration). It can be used for economic inequality and/or regional disparities (Huang/Leung 2009).

Value

A single numeric value of the *Hoover Concentration Index* ($0 < CI < 1$).

Author(s)

Thomas Wieland

References

- Bahrenberg, G./Giese, E./Mevenkamp, N./Nipper, J. (2010): "Statistische Methoden in der Geographie. Band 1: Univariate und bivariate Statistik". Stuttgart: Borntraeger.
- Huang, Y./Leung, Y. (2009): "Measuring Regional Inequality: A Comparison of Coefficient of Variation and Hoover Concentration Index". In: *The Open Geography Journal*, **2**, p. 25-34.
- Portnov, B.A./Felsenstein, D. (2010): "On the suitability of income inequality measures for regional analysis: Some evidence from simulation analysis and bootstrapping tests". In: *Socio-Economic Planning Sciences*, **44**, 4, p. 212-219.

See Also

[cv](#), [gini](#), [herf](#), [theil](#), [atkinson](#), [coulter](#), [disp](#)

Examples

```
# Regional disparities in Germany:
gdp <- c(460.69, 549.19, 124.16, 65.29, 31.59, 109.27, 263.44, 39.87, 258.53,
645.59, 131.95, 35.03, 112.66, 56.22, 85.61, 56.81)
# GDP of german regions 2015 (in billion EUR)
pop <- pop <- c(10879618, 12843514, 3520031, 2484826, 671489, 1787408, 6176172,
1612362, 7926599, 17865516, 4052803, 995597, 4084851, 2245470, 2858714, 2170714)
# population of german regions 2015
hoover(gdp, pop)
```

howard.cl

Howard-Newman-Tarp colocation index

Description

Calculating the colocation index (CL) by Howard, Newman and Tarp for two industries

Usage

```
howard.cl(k, industry, region, industry1, industry2, e_k = NULL)
```

Arguments

<code>k</code>	a vector containing the IDs/names of firms k
<code>industry</code>	a vector containing the IDs/names of the industries i
<code>region</code>	a vector containing the IDs/names of the regions j
<code>industry1</code>	Regarded industry 1 (out of the industry vector)
<code>industry2</code>	Regarded industry 2 (out of the industry vector)
<code>e_k</code>	Employment of firm k

Details

The Howard-Newman-Tarp colocation index (CL) is standardized ($0 \leq CL \leq 1$). Processing time depends on the number of firms.

Value

A single value of CL

Author(s)

Thomas Wieland

References

Howard, E./Newman, C./Tarp, F. (2016): “Measuring industry coagglomeration and identifying the driving forces”. In: *Journal of Economic Geography*, **16**, 5, p. 1055-1078.

See Also

[howard.xcl](#), [howard.xcl2](#), [ellison.c](#), [ellison.c2](#)

Examples

```
# example from Howard et al. (2016):
firms <- 1:6
industries <- c("A", "B", "A", "B", "A", "B")
locations <- c("X", "X", "X", "Y", "Y", "X")

howard.cl(firms, industries, locations, industry1 = "A",
industry2 = "B")
```

howard.xcl

Howard-Newman-Tarp excess colocation (XCL) index

Description

Calculating the excess colocation (XCL) index by Howard, Newman and Tarp for two industries

Usage

```
howard.xcl(k, industry, region, industry1, industry2, no.samples = 50, e_k = NULL)
```

Arguments

<code>k</code>	a vector containing the IDs/names of firms k
<code>industry</code>	a vector containing the IDs/names of the industries i
<code>region</code>	a vector containing the IDs/names of the regions j
<code>industry1</code>	Regarded industry 1 (out of the industry vector)
<code>industry2</code>	Regarded industry 2 (out of the industry vector)
<code>no.samples</code>	Number of samples for the counterfactual firm allocation via bootstrapping
<code>e_k</code>	Employment of firm k

Details

The Howard-Newman-Tarp excess colocation index (*XCL*) is standardized ($-1 \leq CL \leq 1$). The rationale behind is that the *CL* index (see `howard.cl`) is compared to a counterfactual (random) location pattern which is constructed via bootstrapping. Processing time depends on the number of firms and the number of samples.

Value

A single value of *XCL*

Author(s)

Thomas Wieland

References

Howard, E./Newman, C./Tarp, F. (2016): “Measuring industry coagglomeration and identifying the driving forces”. In: *Journal of Economic Geography*, **16**, 5, p. 1055-1078.

See Also

[howard.cl](#), [howard.xcl2](#), [ellison.c](#), [ellison.c2](#)

Examples

```
# example from Howard et al. (2016):
firms <- 1:6
industries <- c("A", "B", "A", "B", "A", "B")
locations <- c("X", "X", "X", "Y", "Y", "X")

howard.xcl(firms, industries, locations, industry1 = "A",
industry2 = "B")
```

howard.xcl2

Howard-Newman-Tarp excess colocation (XCL) index

Description

Calculating the excess colocation (XCL) index by Howard, Newman and Tarp for a given number of industries

Usage

```
howard.xcl2(k, industry, region, print.results = TRUE)
```

Arguments

<code>k</code>	a vector containing the IDs/names of firms k
<code>industry</code>	a vector containing the IDs/names of the industries i
<code>region</code>	a vector containing the IDs/names of the regions j
<code>print.results</code>	logical argument that indicates whether the calculated values are printed or not

Details

The Howard-Newman-Tarp excess colocation index (XCL) is standardized ($-1 \leq CL \leq 1$). The rationale behind is that the CL index (see `howard.cl`) is compared to a counterfactual (random) location pattern which is constructed via bootstrapping. Processing time depends on the number of firms and the number of samples. This function takes a while even for a relatively small number of industries!

Value

A matrix with I rows (one for each industry-industry combination) containing the XCL values

Author(s)

Thomas Wieland

References

Howard, E./Newman, C./Tarp, F. (2016): "Measuring industry coagglomeration and identifying the driving forces". In: *Journal of Economic Geography*, **16**, 5, p. 1055-1078.

See Also

[howard.cl](#), [howard.xcl2](#), [ellison.c](#), [ellison.c2](#)

Examples

```
## Not run:
# example data from Farhauer/Kroell (2014):
data (FK2014_EGC)

howard.xcl2 (FK2014_EGC$firm, FK2014_EGC$industry,
            FK2014_EGC$region)
# this may take a while!

## End(Not run)
```

huff

Huff model

Description

Calculating market areas using the probabilistic market area model by Huff

Usage

```
huff(huffdataset, origins, locations, attrac, dist, gamma = 1, lambda = -2,
     atype = "pow", dtype = "pow", gamma2 = NULL, lambda2 = NULL,
     localmarket_dataset = NULL, origin_id = NULL, localmarket = NULL,
     check_df = TRUE)
```

Arguments

<code>huffdataset</code>	an interaction matrix which is a <code>data.frame</code> containing the origins, locations and the explanatory variables
<code>origins</code>	the column in the interaction matrix <code>huffdataset</code> containing the origins (e.g. ZIP codes)
<code>locations</code>	the column in the interaction matrix <code>huffdataset</code> containing the locations (e.g. store codes)
<code>attrac</code>	the column in the interaction matrix <code>huffdataset</code> containing the attractivity variable (e.g. sales area)
<code>dist</code>	the column in the interaction matrix <code>huffdataset</code> containing the transport costs (e.g. travelling time)
<code>gamma</code>	a single numeric value for the exponential weighting of size (default: 1)
<code>lambda</code>	a single numeric value for the exponential weighting of distance (transport costs, default: -2)
<code>atype</code>	Type of attractivity weighting function: <code>atype = "pow"</code> (power function), <code>atype = "exp"</code> (exponential function) or <code>atype = "logistic"</code> (default: <code>atype = "pow"</code>)
<code>dtype</code>	Type of distance weighting function: <code>dtype = "pow"</code> (power function), <code>dtype = "exp"</code> (exponential function) or <code>dtype = "logistic"</code> (default: <code>dtype = "pow"</code>)
<code>gamma2</code>	if <code>atype = "logistic"</code> a second γ parameter is needed
<code>lambda2</code>	if <code>dtype = "logistic"</code> a second λ parameter is needed
<code>localmarket_dataset</code>	if <code>output = "total"</code> , a <code>data.frame</code> is needed which contains data about the origins
<code>origin_id</code>	the ID variable of the origins in <code>localmarket_dataset</code>
<code>localmarket</code>	the customer/purchasing power potential of the origins in <code>localmarket_dataset</code>
<code>check_df</code>	logical argument that indicates if the given dataset is checked for correct input, only for internal use, should not be deselected (default: TRUE)

Details

The *Huff Model* (Huff 1962, 1963, 1964) is the most popular *spatial interaction model* for retailing and services and belongs to the family of *probabilistic market area models*. The basic idea of the model is that consumer decisions are not deterministic but probabilistic, so the decision of customers for a shopping location in a competitive environment cannot be predicted exactly. The results of the model are probabilities for these decisions, which can be interpreted as market shares of the regarded locations (j) in the customer origins (i), p_{ij} , which can be regarded as an equilibrium solution with logically consistent market shares ($0 < p_{ij} < 1$, $\sum_{j=1}^n p_{ij} = 1$). From a theoretical perspective, the model is based on an utility function with two explanatory variables ("attractivity" of the locations, transport costs between origins and locations), which are weighted by an exponent: $U_{ij} = A_j^\gamma d_{ij}^{-\lambda}$. This specification is relaxed in this case, so both variables can be weighted by a power, exponential or logistic function.

This function computes the market shares from a given interaction matrix and given weighting parameters. The function returns an estimated interaction matrix. If local market information about the origins (e.g. purchasing power, population size etc.) is stated, the location total turnovers are filed in another `data.frame`. Note that each attractivity or distance value must be greater than zero.

Value

A list containing the following objects:

huffmat	A data frame containing the Huff interaction matrix
totals	If total turnovers are estimated: a data frame containing the total values (turnovers) of each location

Note

This function contains code from the authors' package MCI.

Author(s)

Thomas Wieland

References

- Berman, B. R./Evans, J. R. (2012): "Retail Management: A Strategic Approach". 12th edition. Boston : Pearson.
- Huff, D. L. (1962): "Determination of Intra-Urban Retail Trade Areas". Los Angeles : University of California.
- Huff, D. L. (1963): "A Probabilistic Analysis of Shopping Center Trade Areas". In: *Land Economics*, **39**, 1, p. 81-90.
- Huff, D. L. (1964): "Defining and Estimating a Trading Area". In: *Journal of Marketing*, **28**, 4, p. 34-38.
- Levy, M./Weitz, B. A. (2012): "Retailing management". 8th edition. New York : McGraw-Hill Irwin.
- Loeffler, G. (1998): "Market areas - a methodological reflection on their boundaries". In: *GeoJournal*, **45**, 4, p. 265-272.
- Wieland, T. (2015): "Nahversorgung im Kontext raumoeconomischer Entwicklungen im Lebensmitteleinzelhandel - Konzeption und Durchfuehrung einer GIS-gestuetzten Analyse der Strukturen des Lebensmitteleinzelhandels und der Nahversorgung in Freiburg im Breisgau". Projektbericht. Goettingen : GOEDOC, Dokumenten- und Publikationsserver der Georg-August-Universitaet Goettingen. <http://webdoc.sub.gwdg.de/pub/mon/2015/5-wieland.pdf>

See Also

[converse](#), [reilly](#), [hansen](#)

Examples

```
# Example from Levy/Weitz (2009):  
  
# Data for the existing and the new location  
locations <- c("Existing Store", "New Store")  
S_j <- c(5000, 10000)  
location_data <- data.frame(locations, S_j)
```

```

# Data for the two communities (Rock Creek and Oak Hammock)
communities <- c("Rock Creek", "Oak Hammock")
C_i <- c(5000000, 3000000)
community_data <- data.frame(communities, C_i)

# Combining location and submarket data in the interaction matrix
interactionmatrix <- merge (communities, location_data)
# Adding driving time:
interactionmatrix[1,4] <- 10
interactionmatrix[2,4] <- 5
interactionmatrix[3,4] <- 5
interactionmatrix[4,4] <- 15
colnames(interactionmatrix) <- c("communities", "locations", "S_j", "d_ij")

huff_shares <- huff(interactionmatrix, "communities", "locations", "S_j", "d_ij")
huff_shares
# Market shares of the new location:
huff_shares$ijmatrix[huff_shares$ijmatrix$locations == "New Store",]

huff_all <- huff(interactionmatrix, "communities", "locations", "S_j", "d_ij",
localmarket_dataset = community_data, origin_id = "communities", localmarket = "C_i")

huff_all

huff_all$totals

```

krugman.conc

Krugman coefficient of spatial industry concentration for two industries

Description

Calculating the Krugman coefficient for the spatial concentration of two industries based on regional industry data (normally employment data)

Usage

```
krugman.conc(e_ij, e_uj)
```

Arguments

e_ij a numeric vector with the employment of the industry *i* in regions *j*
e_uj a numeric vector with the employment of the industry *u* in region *j*

Details

The *Krugman coefficient of industry concentration* (K_{iu}) is a measure for the dissimilarity of the spatial structure of two industries (i and u) regarding the employment in the j regions. The coefficient K_{iu} varies between 0 (no concentration/same structure) and 2 (maximum difference, that means a complete other spatial structure of the industry compared to the others). The calculation is based on the formulae in Farhauer/Kroell (2013).

Value

A single numeric value ($0 < K_{iu} < 2$)

Author(s)

Thomas Wieland

References

Farhauer, O./Kroell, A. (2013): "Standorttheorien: Regional- und Stadtoekonomik in Theorie und Praxis". Wiesbaden : Springer.

Nakamura, R./Morrison Paul, C. J. (2009): "Measuring agglomeration". In: Capello, R./Nijkamp, P. (eds.): *Handbook of Regional Growth and Development Theories*. Cheltenham: Elgar. p. 305-328.

See Also

[gini.conc](#), [gini.spec](#), [krugman.conc2](#), [krugman.spec](#), [krugman.spec2](#), [locq](#)

Examples

```
E_ij <- c(4388, 37489, 129423, 60941)
E_uj <- E_ij/2
krugman.conc(E_ij, E_uj)
# exactly the same structure (= no concentration)
```

krugman.conc2

Krugman coefficient of spatial industry concentration for more than two industries

Description

Calculating the Krugman coefficient for the spatial concentration of an industry based on regional industry data (normally employment data) compared with a vector of other industries

Usage

```
krugman.conc2(e_ij, e_uj)
```

Arguments

e_ij a numeric vector with the employment of the industry i in regions j
 e_uj a data frame with the employment of the industry u in j regions

Details

The *Krugman coefficient of industry concentration* (K_i) is a measure for the dissimilarity of the spatial structure of one industry (i) compared to several others (u) regarding the employment in the j regions. The coefficient K_{iu} varies between 0 (no concentration/same structure) and 2 (maximum difference, that means a complete other spatial structure of the industry compared to the others). The calculation is based on the formulae in Farhauer/Kroell (2013).

Value

A single numeric value ($0 < K_i < 2$)

Author(s)

Thomas Wieland

References

Farhauer, O./Kroell, A. (2013): “Standorttheorien: Regional- und Stadtoekonomik in Theorie und Praxis”. Wiesbaden : Springer.

Nakamura, R./Morrison Paul, C. J. (2009): “Measuring agglomeration”. In: Capello, R./Nijkamp, P. (eds.): *Handbook of Regional Growth and Development Theories*. Cheltenham: Elgar. p. 305-328.

See Also

[gini.conc](#), [gini.spec](#), [krugman.conc](#), [krugman.spec](#), [krugman.spec2](#), [locq](#)

Examples

```
# Example from Farhauer/Kroell (2013):
Chemie <- c(20000,11000,31000,8000,20000)
Sozialwesen <- c(40000,10000,25000,9000,16000)
Elektronik <- c(10000,11000,14000,14000,13000)
Holz <- c(7000,7500,11000,1500,36000)
Bergbau <- c(4320, 7811, 3900, 2300, 47560)
# five industries
industries <- data.frame(Chemie, Sozialwesen, Elektronik, Holz)
# data frame with all comparison industries
krugman.conc2(Bergbau, industries)
# returns the Krugman coefficient for the concentration
# of the mining industry (Bergbau) compared to
# chemistry (Chemie), social services (Sozialwesen),
# electronics (Elektronik) and wood industry (Holz)
# 0.8619
```

`krugman.spec`*Krugman coefficient of regional specialization for two regions*

Description

Calculating the Krugman coefficient for the specialization of two regions based on regional industry data (normally employment data)

Usage`krugman.spec(e_ij, e_il)`**Arguments**

`e_ij` a numeric vector with the employment of the industries i in region j
`e_il` a numeric vector with the employment of the industries i in region l

Details

The *Krugman coefficient of regional specialization* (K_{jl}) is a measure for the dissimilarity of the industrial structure of two regions (j and l) regarding the employment in the i industries in these regions. The coefficient K_{jl} varies between 0 (no specialization/same structure) and 2 (maximum difference, that means there is no single industry localized in both regions). The calculation is based on the formulae in Farhauer/Kroell (2013).

Value

A single numeric value ($0 < K_{jl} < 2$)

Author(s)

Thomas Wieland

References

Farhauer, O./Kroell, A. (2013): "Standorttheorien: Regional- und Stadtoekonomik in Theorie und Praxis". Wiesbaden : Springer.

Nakamura, R./Morrison Paul, C. J. (2009): "Measuring agglomeration". In: Capello, R./Nijkamp, P. (eds.): *Handbook of Regional Growth and Development Theories*. Cheltenham: Elgar. p. 305-328.

See Also

[gini.conc](#), [gini.spec](#), [krugman.conc](#), [krugman.conc2](#), [krugman.spec2](#), [locq](#)

Examples

```
# Example from Farhauer/Kroell (2013), modified:
E_ij <- c(20,10,70,0,0)
# employment of five industries in region j
E_il <- c(0,0,0,60,40)
# employment of five industries in region l
krugman.spec(E_ij, E_il)
# results the specialization coefficient (2)

# Example Goettingen:
data(Goettingen)
krugman.spec(Goettingen$Goettingen2017[2:16], Goettingen$BRD2017[2:16])
# Returns the Krugman coefficient of regional specialization 2017 (0.4508469)
```

krugman.spec2	<i>Krugman coefficient of regional specialization for more than two regions</i>
---------------	---

Description

Calculating the Krugman coefficient for the specialization of one region based on regional industry data (normally employment data) compared with a vector of other regions

Usage

```
krugman.spec2(e_ij, e_il)
```

Arguments

e_ij	a numeric vector with the employment of the industries i in region j
e_il	a data frame with the employment of the industries i in l regions

Details

The *Krugman coefficient of regional specialization* (K_{jl}) is a measure for the dissimilarity of the industrial structure of regions (j and other regions, l) regarding the employment in the i industries in these regions. The coefficient K_{jl} varies between 0 (no specialization/same structure) and 2 (maximum difference, that means there is no single industry localized in both regions).

Value

A single numeric value ($0 < K_{jl} < 2$)

Author(s)

Thomas Wieland

References

- Farhauer, O./Kroell, A. (2013): “Standorttheorien: Regional- und Stadtoekonomik in Theorie und Praxis”. Wiesbaden : Springer.
- Nakamura, R./Morrison Paul, C. J. (2009): “Measuring agglomeration”. In: Capello, R./Nijkamp, P. (eds.): *Handbook of Regional Growth and Development Theories*. Cheltenham: Elgar. p. 305-328.

See Also

[gini.conc](#), [gini.spec](#), [krugman.spec](#), [krugman.conc](#), [krugman.conc2](#), [locq](#)

Examples

```
# Example from Farhauer/Kroell (2013):
Sweden <- c(45000, 15000, 32000, 10000, 30000)
Norway <- c(35000, 12000, 30000, 8000, 22000)
Denmark <- c(40000, 10000, 25000, 9000, 18000)
Finland <- c(30000, 11000, 18000, 3000, 13000)
Island <- c(40000, 6000, 11000, 2000, 12000)
# industry jobs in five industries for five countries
countries <- data.frame(Norway, Denmark, Finland, Island)
# data frame with all comparison countries
krugman.spec2(Sweden, countries)
# returns the Krugman coefficient for the specialization
# of sweden compared to Norway, Denmark, Finland and Island
# 0.1595
```

litzenberger

Litzenberger-Sternberg Cluster Index

Description

Calculating the *Cluster Index* by Litzenberger and Sternberg

Usage

```
litzenberger(e_ij, e_i, a_j, a, p_j, p, b_ij, b_i)
```

Arguments

e_ij	a single numeric value with the employment of industry <i>i</i> in region <i>j</i>
e_i	a single numeric value with the over-all employment in industry <i>i</i>
a_j	a single numeric value of the area of region <i>j</i>
a	a single numeric value of the total area
p_j	a single numeric value of the population of region <i>j</i>
p	a single numeric value of the total population
b_ij	a single numeric value of the number of firms of industry <i>i</i> in region <i>j</i>
b_i	a single numeric value of the total number of firms of industry <i>i</i>

Details

The Litzenberger-Sternberg Cluster Index is not standardized and depends on the number of regarded industries and regions.

Value

A single numeric value of (*CI*).

Author(s)

Thomas Wieland

References

Farhauer, O./Kroell, A. (2014): "Standorttheorien: Regional- und Stadtoekonomik in Theorie und Praxis". Wiesbaden : Springer.

Hoffmann J./Hirsch, S./Simons, J. (2017): "Identification of spatial agglomerations in the German food processing industry". In: *Papers in Regional Science*, **96**, 1, p. 139-162.

Litzenberger, T./Sternberg, R. (2006): "Der Clusterindex - eine Methodik zur Identifizierung regionaler Cluster am Beispiel deutscher Industriebranchen". In: *Geographische Zeitschrift*, **94**, 2, p. 209-224.

See Also

[litzenberger2](#), [gini.conc](#), [gini.spec](#), [locq](#), [locq2](#), [ellison.a](#), [ellison.a2](#), [ellison.c](#), [ellison.c2](#)

Examples

```
# Example from Farhauer/Kroell (2014):
litzenberger(e_ij = 1743, e_i = 5740, a_j = 50,
a = 576, p_j = 488, p = 4621, b_ij = 35, b_i = 53)
# 21.87491
```

litzenberger2

Litzenberger-Sternberg Cluster Index

Description

Calculating the *Cluster Index* by Litzenberger and Sternberg for a given number of *I* industries and *J* regions

Usage

```
litzenberger2(e_ij, industry.id, region.id, a_j, p_j, b_ij,
CI.output = "mat", na.rm = TRUE)
```

Arguments

<code>e_ij</code>	a vector with the employment of industry i in region j
<code>industry.id</code>	a vector containing the IDs of the industries i
<code>region.id</code>	a vector containing the IDs of the regions j
<code>a_j</code>	a vector containing the areas of the regions j
<code>p_j</code>	a vector containing the populations of the regions j
<code>b_ij</code>	a vector containing the numbers of firms of industry i in region j
<code>CI.output</code>	Type of output: matrix (default: <code>CI.output = "mat"</code>) or data frame (<code>CI.output = "df"</code>)
<code>na.rm</code>	logical argument that indicates whether NA values should be excluded before computing results

Details

The Litzenberger-Sternberg Cluster Index is not standardized and depends on the number of regarded industries and regions.

Value

A matrix or data frame containing $I \times J$ values of CI

Author(s)

Thomas Wieland

References

- Farhauer, O./Kroell, A. (2014): "Standorttheorien: Regional- und Stadtoekonomie in Theorie und Praxis". Wiesbaden : Springer.
- Hoffmann J./Hirsch, S./Simons, J. (2017): "Identification of spatial agglomerations in the German food processing industry". In: *Papers in Regional Science*, **96**, 1, p. 139-162.
- Litzenberger, T./Sternberg, R. (2006): "Der Clusterindex - eine Methodik zur Identifizierung regionaler Cluster am Beispiel deutscher Industriebranchen". In: *Geographische Zeitschrift*, **94**, 2, p. 209-224.

See Also

[litzenberger](#), [gini.conc](#), [gini.spec](#), [locq](#), [locq2](#), [ellison.a](#), [ellison.a2](#), [ellison.c](#), [ellison.c2](#)

Examples

```
data (G.regions.industries)

lss <- litzenberger2(G.regions.industries$emp_all,
  G.regions.industries$ind_code, G.regions.industries$region_code,
  G.regions.industries$area_sqkm, G.regions.industries$pop,
  G.regions.industries$firms, CI.output = "df")
```

```
# output as data frame

lss_sort <- lss[order(lss$CI, decreasing = TRUE),]
# Sort decreasing by size of CI

lss_sort[1:5,]
```

lm.beta

Beta regression coefficients

Description

Calculating the standardized (beta) regression coefficients of linear models

Usage

```
lm.beta(linmod, dummy.na = TRUE)
```

Arguments

linmod	A lm object (linear regression model) with more than one independent variable
dummy.na	logical argument that indicates if dummy variables should be ignored when calculating the beta weights (default: TRUE). Note that beta weights of dummy variables do not make any sense

Details

Standardized coefficients (beta coefficients) show how many standard deviations a dependent variable will change when the regarded independent variable is increased by a standard deviation. The β values are used in multiple linear regression models to compare the real effect (power) of the independent variables when they are measured in different units. Note that β values do not make any sense for dummy variables since they cannot change by a standard deviation.

Value

A list containing all independent variables and the corresponding standardized coefficients.

Author(s)

Thomas Wieland

References

Backhaus, K./Erichson, B./Plinke, W./Weiber, R. (2016): "Multivariate Analysemethoden: Eine anwendungsorientierte Einfuehrung". Berlin: Springer.

Examples

```
x1 <- runif(100)
x2 <- runif(100)
# random values for two independent variables (x1, x2)
y <- runif(100)
# random values for the dependent variable (y)
testmodel <- lm(y~x1+x2)
# OLS regression
summary(testmodel)
# summary
lm.beta(testmodel)
# beta coefficients
```

locq

*Location quotient***Description**

Calculating the *location quotient* (a.k.a. *Hoover-Balassa quotient*)

Usage

```
locq(e_ij, e_j, e_i, e, industry.names = NULL, plot.results = FALSE,
LQ.method = "m", plot.title = "Localization quotients",
bar.col = "lightblue", line.col = "red", arg.size = 1)
```

Arguments

e_ij	a single numeric value or vector with the employment of industry/industries <i>i</i> in region <i>j</i>
e_j	a single numeric value with the over-all employment in region <i>j</i>
e_i	a single numeric value or vector with the over-all employment in industry/industries <i>i</i>
e	a single numeric value with the over-all employment in all regions
industry.names	Industry names (e.g. from the relevant statistical classification of economic activities)
plot.results	Logical argument that indicates if the results have to be plotted (only available if <i>i</i> > 1)
LQ.method	Indicates whether the multiplicative (default: LQ.method = "m") or the additive LQ (LQ.method = "a") is computed
plot.title	If plot.results = TRUE: Plot title
bar.col	If plot.results = TRUE: Bar colour
line.col	If plot.results = TRUE: LQ1-line colour
arg.size	If plot.results = TRUE: Size of industry names in bar plot

Details

The *location quotient* is a simple measure for the concentration of an industry (i) in a region (j) and is also the mathematical basis for other related indicators in regional economics (e.g. `gini.conc()`). The function returns the value LQ which is equal to 1 if the concentration of the regarded industry is exactly the same as the over-all concentration (that means, it is proportionally represented in region j). If the value of LQ is smaller (bigger) than 1, the industry is underrepresented (overrepresented). The function checks the input values for errors (i.e. if employment in a region is bigger than over-all employment).

Value

A single numeric value of (LQ) or a matrix with respect to all i industries. Optional: plot.

Author(s)

Thomas Wieland

References

- Farhauer, O./Kroell, A. (2013): “Standorttheorien: Regional- und Stadtoekonomik in Theorie und Praxis”. Wiesbaden : Springer.
- Hoen A.R./Oosterhaven, J. (2006): “On the measure of comparative advantage”. In: *The Annals of Regional Science*, **40**, 3, p. 677-691.
- Nakamura, R./Morrison Paul, C. J. (2009): “Measuring agglomeration”. In: Capello, R./Nijkamp, P. (eds.): *Handbook of Regional Growth and Development Theories*. Cheltenham: Elgar. p. 305-328.
- O’Donoghue, D./Gleave, B. (2004): “A Note on Methods for Measuring Industrial Agglomeration”. In: *Regional Studies*, **38**, 4, p. 419-427.
- Tian, Z. (2013): “Measuring agglomeration using the standardized location quotient with a bootstrap method”. In: *Journal of Regional Analysis and Policy*, **43**, 2, p. 186-197.

See Also

[gini.conc](#), [gini.spec](#), [locq2](#)

Examples

```
# Example from Farhauer/Kroell (2013):
locq (1714, 79006, 879213, 15593224)
# returns the location quotient (0.3847623)

# Location quotients for Goettingen 2017:
data(Goettingen)
locq (Goettingen$Goettingen2017[2:16], Goettingen$Goettingen2017[1],
Goettingen$BRD2017[2:16], Goettingen$BRD2017[1])
```

locq.growth	<i>Portfolio matrix for specialization and growth</i>
-------------	---

Description

Portfolio matrix plot comparing two numeric vectors (here: specialization and growth)

Usage

```
locq.growth(e_ij1, e_ij2, e_i1, e_i2, industry.names = NULL,
  y.axis = "r",
  psize, psize.factor = 10, time.periods = NULL,
  pmx = "Regional specialization", pmy = "Regional growth",
  pmtitle = "Portfolio matrix", pcol = NULL, pcol.border = NULL,
  leg = FALSE, leg.fsize = 1, leg.col = NULL,
  leg.x = 0, leg.y = y_min*1.5,
  bg.col = "gray95", bgrid = TRUE, bgrid.col = "white",
  bgrid.size = 2, bgrid.type = "solid",
  seg.x = 1, seg.y = 0)
```

Arguments

e_ij1	a numeric vector with i values containing the employment in i industries in region j at time 1
e_ij2	a numeric vector with i values containing the employment in i industries in region j at time 2
e_i1	a numeric vector with i values containing the total employment in i industries at time 1
e_i2	a numeric vector with i values containing the total employment in i industries at time 2
industry.names	Industry names (e.g. from the relevant statistical classification of economic activities)
y.axis	Declares which values shall be plotted on the Y axis: If <code>y.axis = "r"</code> , the Y axis shows the <i>regional</i> growth. If <code>y.axis = "n"</code> , the Y axis shows the <i>national</i> growth. To set both growths in ratio, choose <code>y.axis = "rn"</code> (regional vs. national growth)
psize	Point size in the portfolio matrix plot (mostly the absolute values of employment in i industries in region j at time 2)
psize.factor	Enlargement factor for the points in the plot
time.periods	No. of regarded time periods (for average growth rates)
pmx	Name of the X axis in the plot
pmy	Name of the Y axis in the plot
pmtitle	Plot title

<code>pcol</code>	Industry-specific point colors
<code>pcol.border</code>	Color of point border
<code>leg</code>	Logical argument that indicates if a legend has to be added to the plot
<code>leg.fsize</code>	If <code>leg = TRUE</code> : Font size in the plot legend
<code>leg.col</code>	No. of columns in the plot legend
<code>leg.x</code>	If <code>leg = TRUE</code> : X coordinate of the legend
<code>leg.y</code>	If <code>leg = TRUE</code> : Y coordinate of the legend
<code>bg.col</code>	Background color
<code>bgrid</code>	Logical argument that indicates if a grid has to be added to the plot
<code>bgrid.col</code>	If <code>bgrid = TRUE</code> : Color of the grid
<code>bgrid.size</code>	If <code>bgrid = TRUE</code> : Size of the grid
<code>bgrid.type</code>	If <code>bgrid = TRUE</code> : Type of the grid
<code>seg.x</code>	X coordinate of segmentation of the plot
<code>seg.y</code>	Y coordinate of segmentation of the plot

Details

The *portfolio matrix* is a graphic tool displaying the development of one variable compared to another variable. The plot shows the regarded variable on the x axis and a variable with which it is confronted on the y axis while the graph is divided in four quadrants. Originally, the *portfolio matrix* was developed by the *Boston Consulting Group* to analyze the performance of product lines in marketing, also known as the *growth-share matrix*. The quadrants show the performance of the regarded objects (stars, cash cows, question marks, dogs) (Henderson 1973). But the *portfolio matrix* can also be used to analyze/illustrate the world market integration of a region or a national economy by confronting e.g. the increase in world market share (x axis) and the world trade growth (y axis) (Baker et al. 2002). Another option is to analyze/illustrate the economic performance of a region (Howard 2007). E.g. it is possible to confront the growth of industries in a region with the all-over growth of these industries in the national economy.

This function is a special case of portfolio matrix, showing the regional specialization on the X axis instead of the regional growth (which can be plotted on the Y axis).

Value

A portfolio matrix plot.

Invisible: a list containing the following items:

<code>portfolio.data</code>	The data related to the plot
<code>locq</code>	The localization quotients for each year
<code>growth</code>	The growth values for each industry

Author(s)

Thomas Wieland

References

Baker, P./von Kirchbach, F./Mimouni, M./Pasteels, J.-M. (2002): “Analytical tools for enhancing the participation of developing countries in the Multilateral Trading System in the context of the Doha Development Agenda”. In: *Aussenwirtschaft*, **57**, 3, p. 343-372.

Howard, D. (2007): “A regional economic performance matrix - an aid to regional economic policy development”. In: *Journal of Economic and Social Policy*, **11**, 2, Art. 4.

Henderson, B. D. (1973): “The Experience Curve - Reviewed, IV. The Growth Share Matrix or The Product Portfolio”. The Boston Consulting Group (BCG).

See Also

[locq](#), [portfolio](#), [shift](#), [shiftd](#), [shifti](#)

Examples

```
data(Goettingen)
# Loads employment data for Goettingen and Germany (2008-2017)

locq.growth(Goettingen$Goettingen2008[2:16], Goettingen$Goettingen2017[2:16],
Goettingen$BRD2008[2:16], Goettingen$BRD2017[2:16],
psize = Goettingen$Goettingen2017[2:16],
industry.names = Goettingen$WA_WZ2008[2:16], pcol.border = "grey",
leg = TRUE, leg.fsize = 0.4, leg.x = -0.2)
```

locq2

Location quotient

Description

Calculating the *location quotient* (a.k.a. *Hoover-Balassa quotient*) for a given number of I industries and J regions

Usage

```
locq2(e_ij, industry.id, region.id, LQ.norm = "none",
LQ.output = "mat", na.rm = TRUE)
```

Arguments

<code>e_ij</code>	a vector with the employment of industry i in region j
<code>industry.id</code>	a vector containing the IDs of the industries i
<code>region.id</code>	a vector containing the IDs of the regions j
<code>LQ.norm</code>	Type of normalization of the location quotients: no normalization (default: <code>LQ.norm = "none"</code>), z values (<code>LQ.norm = "OG"</code>) or z values of logged location quotients (<code>LQ.norm = "T"</code>)

LQ.output	Type of output: matrix (default: CI.output = "mat") or data frame (CI.output = "df")
na.rm	logical argument that indicates whether NA values should be excluded before computing results

Details

The *location quotient* is a simple measure for the concentration of an industry (i) in a region (j) and is also the mathematical basis for other related indicators in regional economics (e.g. `gini.conc()`). The function returns the value LQ which is equal to 1 if the concentration of the regarded industry is exactly the same as the over-all concentration (that means, it is proportionally represented in region j). If the value of LQ is smaller (bigger) than 1, the industry is underrepresented (overrepresented). The function checks the input values for errors (i.e. if employment in a region is bigger than over-all employment).

Two types of normalization are available: z values of the location quotients (O'Donoghue/Gleave 2004) or z values of logged location quotients (Tian 2013).

Value

A matrix or data frame containing $I \times J$ values of LQ

Author(s)

Thomas Wieland

References

- Farhauer, O./Kroell, A. (2014): "Standorttheorien: Regional- und Stadtoekonomik in Theorie und Praxis". Wiesbaden : Springer.
- Hoen A.R./Oosterhaven, J. (2006): "On the measure of comparative advantage". In: *The Annals of Regional Science*, **40**, 3, p. 677-691.
- Nakamura, R./Morrison Paul, C. J. (2009): "Measuring agglomeration". In: Capello, R./Nijkamp, P. (eds.): *Handbook of Regional Growth and Development Theories*. Cheltenham: Elgar. p. 305-328.
- O'Donoghue, D./Gleave, B. (2004): "A Note on Methods for Measuring Industrial Agglomeration". In: *Regional Studies*, **38**, 4, p. 419-427.
- Tian, Z. (2013): "Measuring agglomeration using the standardized location quotient with a bootstrap method". In: *Journal of Regional Analysis and Policy*, **43**, 2, p. 186-197.

See Also

[litzengerger](#), [gini.conc](#), [gini.spec](#), [locq](#), [hoover](#), [ellison.a](#), [ellison.a2](#), [ellison.c](#), [ellison.c2](#)

Examples

```
data (G.regions.industries)

lqs <- locq2(e_ij = G.regions.industries$emp_all,
```

```
G.regions.industries$ind_code, G.regions.industries$region_code,
LQ.output = "df")
# output as data frame

lqs_sort <- lqs[order(lqs$LQ, decreasing = TRUE),]
# Sort decreasing by size of LQ

lqs_sort[1:5,]
```

lorenz

Lorenz curve

Description

Calculating and plotting the Lorenz curve

Usage

```
lorenz(x, weighting = NULL, z = NULL, na.rm = TRUE,
lcx = "% of objects", lcy = "% of regarded variable",
lctitle = "Lorenz curve", le.col = "blue", lc.col = "black",
lsize = 1.5, ltype = "solid", bg.col = "gray95", bgrid = TRUE,
bgrid.col = "white", bgrid.size = 2, bgrid.type = "solid",
lcg = FALSE, lcg.n = FALSE, lcg.caption = NULL, lcg.lab.x = 0,
lcg.lab.y = 1, add.lc = FALSE, plot.lc = TRUE)
```

Arguments

x	A numeric vector (e.g. dataset of household income, sales turnover or supply)
weighting	A numeric vector containing the weighting data (e.g. size of income classes when calculating a Lorenz curve for aggregated income data)
z	A numeric vector for (optionally) comparing the cumulative distribution
na.rm	logical argument that indicates whether NA values should be excluded before computing results
lcx	defines the x axis label
lcy	defines the y axis label
lctitle	defines the overall title of the Lorenz curve plot
le.col	defines the color of the diagonale (line of equality)
lc.col	defines the color of the Lorenz curve
lsize	defines the size of the lines (default: 1)
ltype	defines the type of the lines (default: "solid")
bg.col	defines the background color of the plot (default: "gray95")
bgrid	logical argument that indicates if a grid is shown in the plot

<code>bgrid.col</code>	if <code>bgrid = TRUE</code> (background grid), <code>bgrid.col</code> defines the color of the background grid (default: "white")
<code>bgrid.size</code>	if <code>bgrid = TRUE</code> (background grid), <code>bgrid.size</code> defines the size of the background grid (default: 2)
<code>bgrid.type</code>	if <code>bgrid = TRUE</code> (background grid), <code>bgrid.type</code> defines the type of lines of the background grid (default: "solid")
<code>lcg</code>	logical argument that indicates if the non-standardized Gini coefficient is displayed in the Lorenz curve plot
<code>lcgn</code>	logical argument that indicates if the standardized Gini coefficient is displayed in the Lorenz curve plot
<code>lcg.caption</code>	specifies the caption above the coefficients
<code>lcg.lab.x</code>	specifies the x coordinate of the label
<code>lcg.lab.y</code>	specifies the y coordinate of the label
<code>add.lc</code>	specifies if a new Lorenz curve is plotted (<code>add.lc = "FALSE"</code>) or the plot is added to an existing Lorenz curve plot (<code>add.lc = "TRUE"</code>)
<code>plot.lc</code>	logical argument that indicates if the Lorenz curve itself is plotted (if <code>plot.lc = FALSE</code> , only the line of equality is plotted)

Details

The *Gini coefficient* (Gini 1912) is a popular measure of statistical dispersion, especially used for analyzing inequality or concentration. The *Lorenz curve* (Lorenz 1905), though developed independently, can be regarded as a graphical representation of the degree of inequality/concentration calculated by the *Gini coefficient* (G) and can also be used for additional interpretations of it. In an economic-geographical context, these methods are frequently used to analyse the concentration/inequality of income or wealth within countries (Aoyama et al. 2011). Other areas of application are analyzing regional disparities (Lessmann 2005, Nakamura 2008) and concentration in markets (sales turnover of competing firms) which makes Gini and Lorenz part of economic statistics in general (Doersam 2004, Roberts 2014).

The *Gini coefficient* (G) varies between 0 (no inequality/concentration) and 1 (complete inequality/concentration). The *Lorenz curve* displays the deviations of the empirical distribution from a perfectly equal distribution as the difference between two graphs (the distribution curve and a diagonal line of perfect equality). This function calculates G and plots the *Lorenz curve* optionally. As there are several ways to calculate the *Gini coefficient*, this function uses the formula given in Doersam (2004). Because the maximum of G is not equal to 1, also a standardized coefficient (G^*) with a maximum equal to 1 can be calculated alternatively. If a Lorenz curve for aggregated data (e.g. income classes with averaged incomes) or the Lorenz curve has to be weighted, use a weighting vector (e.g. size of the income classes).

Value

A plot of the *Lorenz curve*.

Author(s)

Thomas Wieland

References

- Aoyama, Y./Murphy, J. T./Hanson, S. (2011): “Key Concepts in Economic Geography”. London : SAGE.
- Bahrenberg, G./Giese, E./Mevenkamp, N./Nipper, J. (2010): “Statistische Methoden in der Geographie. Band 1: Univariate und bivariate Statistik”. Stuttgart: Borotraeger.
- Cerlani, L./Verme, P. (2012): “The origins of the Gini index: extracts from Variabilita e Mutabilita (1912) by Corrado Gini”. In: *The Journal of Economic Inequality*, **10**, 3, p. 421-443.
- Doersam, P. (2004): “Wirtschaftsstatistik anschaulich dargestellt”. Heidenau : PD-Verlag.
- Gini, C. (1912): “Variabilita e Mutabilita”. Contributo allo Studio delle Distribuzioni e delle Relazioni Statistiche. Bologna : Cuppini.
- Lessmann, C. (2005): “Regionale Disparitaeten in Deutschland und ausgesuchten OECD-Staaten im Vergleich”. *ifo Dresden berichtet*, **3/2005**. https://www.ifo.de/DocDL/ifoedb_2005_3_25-33.pdf.
- Lorenz, M. O. (1905): “Methods of Measuring the Concentration of Wealth”. In: *Publications of the American Statistical Association*, **9**, 70, p. 209-219.
- Nakamura, R. (2008): “Agglomeration Effects on Regional Economic Disparities: A Comparison between the UK and Japan”. In: *Urban Studies*, **45**, 9, p. 1947-1971.
- Roberts, T. (2014): “When Bigger Is Better: A Critique of the Herfindahl-Hirschman Index’s Use to Evaluate Mergers in Network Industries”. In: *Pace Law Review*, **34**, 2, p. 894-946.

See Also

[cv](#), [gini.conc](#), [gini.spec](#), [herf](#), [hoover](#)

Examples

```
# Market concentration (example from Doersam 2004):
sales <- c(20,50,20,10)
# sales turnover of four car manufacturing companies
lorenz (sales, lcx = "percentage of companies", lcy = "percentage of sales",
lctitle = "Lorenz curve of sales", lcg = TRUE, lcg = TRUE)
# plots the Lorenz curve with user-defined title and labels
# including Gini coefficient

# Income classes (example from Doersam 2004):
income <- c(500, 1500, 2500, 4000, 7500, 15000)
# average income of 6 income classes
sizeofclass <- c(1000, 1200, 1600, 400, 200, 600)
# size of income classes
lorenz (income, weighting = sizeofclass, lcg = TRUE, lcg = TRUE)
# plots the Lorenz curve with user-defined title and labels
# including Gini coefficient

# Regional disparities in Germany:
gdp <- c(460.69, 549.19, 124.16, 65.29, 31.59, 109.27, 263.44, 39.87, 258.53,
645.59, 131.95, 35.03, 112.66, 56.22, 85.61, 56.81)
# GDP of german regions 2015 (in billion EUR)
lorenz (gdp, lcg = TRUE, lcg = TRUE)
```

```
# plots the Lorenz curve with user-defined title and labels
# including Gini coefficient
```

mean2	<i>Calculation of mean (extended)</i>
-------	---------------------------------------

Description

Calculating the arithmetic mean, weighted or non-weighted, or the geometric mean

Usage

```
mean2(x, weighting = NULL, output = "mean", na.rm = TRUE)
```

Arguments

x	a numeric vector
weighting	a numeric vector containing weighting data to compute the weighted arithmetic mean (instead of the non-weighted)
output	argument to specify the output (output = "mean" returns the arithmetic mean, output = "geom" returns the geometric mean)
na.rm	logical argument that whether NA values should be extracted or not

Details

This function uses the formula for the weighted arithmetic mean from Sheret (1984).

Value

Single numeric value. If output = "mean" and weighting is specified, the function returns a weighted arithmetic mean. If output = "geom", the geometric mean is returned.

Author(s)

Thomas Wieland

References

Bahrenberg, G./Giese, E./Mevenkamp, N./Nipper, J. (2010): "Statistische Methoden in der Geographie. Band 1: Univariate und bivariate Statistik". Stuttgart: Borntraeger.

Sheret, M. (1984): "The Coefficient of Variation: Weighting Considerations". In: *Social Indicators Research*, **15**, 3, p. 289-295.

See Also

[sd2](#)

Examples

```
avector <- c(5, 17, 84, 55, 39)
mean(avector)
mean2(avector)
wvector <- c(9, 757, 44, 18, 682)
mean2 (avector, weighting = wvector)
mean2 (avector, output = "geom")
```

mssd

Mean square successive difference

Description

Calculating the mean square successive difference

Usage

```
mssd (x)
```

Arguments

x a numeric vector arranged in chronological order

Details

The *mean square successive difference*, δ^2 , is a dimensionless measure of variability over time (von Neumann et al. 1941). It can be used for assessing the volatility of a variable with respect to different subjects/groups.

Value

Single numeric value (the *mean square successive difference*, δ^2).

Author(s)

Thomas Wieland

References

Von Neumann, J./Kent, R. H./Bellinson, H. R./Hart, B. I. (1941): "The mean square successive difference". In: *The Annals of Mathematical Statistics*, **12**, 2, p. 153-162.

See Also

[var2](#), [sd2](#), [cv](#)

Examples

```

data1 <- c(10,10,10,20,20,20,30,30,30)
# stable growth
data2 <- c(20,10,30,10,30,20,30,20,10)
# high variability

# Means:
mean2(data1)
mean2(data2)
# Same means

# Standard deviation:
sd2(data1)
sd2(data2)
# Coefficient of variation:
cv(data1)
cv(data2)
# Measures of statistical dispersion are equal

mssd(data1)
mssd(data2)
# high differences in variability

```

portfolio

Portfolio matrix

Description

Portfolio matrix plot comparing two numeric vectors

Usage

```

portfolio(e_ij1, e_ij2, e_i1, e_i2, industry.names = NULL,
psize, psize.factor = 10, time.periods = NULL,
pmx = "Regional growth", pmy = "National growth",
pmtitle = "Portfolio matrix", pcol = NULL, pcol.border = NULL,
leg = FALSE, leg.fsize = 1, leg.col = NULL,
leg.x = -max_val, leg.y = -max_val*1.5,
bg.col = "gray95", bgrid = TRUE, bgrid.col = "white",
bgrid.size = 2, bgrid.type = "solid",
seg.x = 0, seg.y = 0)

```

Arguments

e_ij1	a numeric vector with i values containing the employment in i industries in region j at time 1
e_ij2	a numeric vector with i values containing the employment in i industries in region j at time 2

e_i1	a numeric vector with i values containing the total employment in i industries at time 1
e_i2	a numeric vector with i values containing the total employment in i industries at time 2
industry.names	Industry names (e.g. from the relevant statistical classification of economic activities)
psize	Point size in the portfolio matrix plot (mostly the absolute values of employment in i industries in region j at time 2)
psize.factor	Enlargement factor for the points in the plot
time.periods	No. of regarded time periods (for average growth rates)
pmx	Name of the X axis in the plot
pmy	Name of the Y axis in the plot
pmtitle	Plot title
pcol	Industry-specific point colors
pcol.border	Color of point border
leg	Logical argument that indicates if a legend has to be added to the plot
leg.fsize	If leg = TRUE: Font size in the plot legend
leg.col	No. of columns in the legend
leg.x	If leg = TRUE: X coordinate of the legend
leg.y	If leg = TRUE: Y coordinate of the legend
bg.col	Background color
bgrid	Logical argument that indicates if a grid has to be added to the plot
bgrid.col	If bgrid = TRUE: Color of the grid
bgrid.size	If bgrid = TRUE: Size of the grid
bgrid.type	If bgrid = TRUE: Type of the grid
seg.x	X coordinate of segmentation of the plot
seg.y	Y coordinate of segmentation of the plot

Details

The *portfolio matrix* is a graphic tool displaying the development of one variable compared to another variable. The plot shows the regarded variable on the x axis and a variable with which it is confronted on the y axis while the graph is divided in four quadrants. Originally, the *portfolio matrix* was developed by the *Boston Consulting Group* to analyze the performance of product lines in marketing, also known as the *growth-share matrix*. The quadrants show the performance of the regarded objects (stars, cash cows, question marks, dogs) (Henderson 1973). But the *portfolio matrix* can also be used to analyze/illustrate the world market integration of a region or a national economy by confronting e.g. the increase in world market share (x axis) and the world trade growth (y axis) (Baker et al. 2002). Another option is to analyze/illustrate the economic performance of a region (Howard 2007). E.g. it is possible to confront the growth of industries in a region with the all-over growth of these industries in the national economy.

Value

A portfolio matrix plot and a data frame containing the related data (invisible).

Author(s)

Thomas Wieland

References

Baker, P./von Kirchbach, F./Mimouni, M./Pasteels, J.-M. (2002): “Analytical tools for enhancing the participation of developing countries in the Multilateral Trading System in the context of the Doha Development Agenda”. In: *Aussenwirtschaft*, **57**, 3, p. 343-372.

Howard, D. (2007): “A regional economic performance matrix - an aid to regional economic policy development”. In: *Journal of Economic and Social Policy*, **11**, 2, Art. 4.

Henderson, B. D. (1973): “The Experience Curve - Reviewed, IV. The Growth Share Matrix or The Product Portfolio”. The Boston Consulting Group (BCG).

See Also

[shift](#), [shiftd](#), [shifti](#)

Examples

```
data(Freiburg)
# Loads employment data for Freiburg and Germany (2008 and 2014)

portfolio(Freiburg$e_Freiburg2008, Freiburg$e_Freiburg2014,
Freiburg$e_Germany2008, Freiburg$e_Germany2014,
industry.names = Freiburg$industry, Freiburg$e_Freiburg2014, psize.factor = 12,
pmx = "Freiburg", pmy = "Deutschland", pmtitle = "Freiburg und BRD",
pcol = Freiburg$color, leg = TRUE, leg.fsize = 0.6, bgrid = TRUE, leg.y = -0.17)
```

Description

This function provides the analysis of absolute and conditional regional economic beta convergence and sigma convergence for cross-sectional data. Beta convergence can be estimated using an OLS or NLS technique. Sigma convergence can be analyzed using ANOVA or trend regression.

Usage

```
rca(gdp1, time1, gdp2, time2,
    conditions = NULL, conditions.formula = NULL, conditions.startval = NULL,
    beta.estimate = "ols", beta.plot = FALSE, beta.plotPSize = 1, beta.plotPCol = "black",
    beta.plotLine = FALSE, beta.plotLineCol = "red", beta.plotX = "Ln (initial)",
    beta.plotY = "Ln (growth)", beta.plotTitle = "Beta convergence", beta.bgCol = "gray95",
    beta.bgrid = TRUE, beta.bgridCol = "white", beta.bgridSize = 2, beta.bgridType = "solid",
    sigma.type = "anova", sigma.measure = "sd", sigma.log = TRUE, sigma.weighting = NULL,
    sigma.issample = FALSE, sigma.plot = FALSE, sigma.plotLSize = 1,
    sigma.plotLineCol = "black", sigma.plotRLine = FALSE, sigma.plotRLineCol = "blue",
    sigma.Ymin = 0, sigma.plotX = "Time", sigma.plotY = "Variation",
    sigma.plotTitle = "Sigma convergence", sigma.bgCol = "gray95", sigma.bgrid = TRUE,
    sigma.bgridCol = "white", sigma.bgridSize = 2, sigma.bgridType = "solid")
```

Arguments

<code>gdp1</code>	A numeric vector containing the GDP per capita (or another economic variable) at time t
<code>time1</code>	A single value of time t (= the initial year)
<code>gdp2</code>	A numeric vector containing the GDP per capita (or another economic variable) at time $t+1$ or a data frame containing the GDPs per capita (or another economic variable) at time $t+1, t+2, t+3, \dots, t+n$
<code>time2</code>	A single value of time $t+1$ or t_n , respectively
<code>conditions</code>	A data frame containing the conditions for conditional beta convergence
<code>conditions.formula</code>	If <code>beta.estimate = "nls"</code> : A formula for the functional linkage of the conditions in the case of conditional beta convergence
<code>conditions.startval</code>	If <code>beta.estimate = "nls"</code> : Starting values for the parameters of the conditions in the case of conditional beta convergence
<code>beta.estimate</code>	Beta estimate via ordinary least squares (OLS) or nonlinear least squares (NLS). Default: <code>beta.estimate = "ols"</code>
<code>beta.plot</code>	Boolean argument that indicates if a plot of beta convergence has to be created
<code>beta.plotPSize</code>	If <code>beta.plot = TRUE</code> : Point size in the beta convergence plot
<code>beta.plotPCol</code>	If <code>beta.plot = TRUE</code> : Point color in the beta convergence plot
<code>beta.plotLine</code>	If <code>beta.plot = TRUE</code> : Logical argument that indicates if a regression line has to be added to the plot
<code>beta.plotLineCol</code>	If <code>beta.plot = TRUE</code> and <code>beta.plotLine = TRUE</code> : Line color of regression line
<code>beta.plotX</code>	If <code>beta.plot = TRUE</code> : Name of the X axis
<code>beta.plotY</code>	If <code>beta.plot = TRUE</code> : Name of the Y axis
<code>beta.plotTitle</code>	If <code>beta.plot = TRUE</code> : Plot title
<code>beta.bgCol</code>	If <code>beta.plot = TRUE</code> : Plot background color

<code>beta.bgrid</code>	If <code>beta.plot = TRUE</code> : Logical argument that indicates if the plot contains a grid
<code>beta.bgridCol</code>	If <code>beta.plot = TRUE</code> and <code>beta.bgrid = TRUE</code> : Color of the grid
<code>beta.bgridSize</code>	If <code>beta.plot = TRUE</code> and <code>beta.bgrid = TRUE</code> : Size of the grid
<code>beta.bgridType</code>	If <code>beta.plot = TRUE</code> and <code>beta.bgrid = TRUE</code> : Type of the grid
<code>sigma.type</code>	Estimating sigma convergence via ANOVA (two years) or trend regression (more than two years). Default: <code>sigma.type = "anova"</code>
<code>sigma.measure</code>	argument that indicates how the sigma convergence should be measured. The default is <code>output = "sd"</code> , which means that the standard deviation is used. If <code>output = "var"</code> or <code>output = "cv"</code> , the variance or the coefficient of variation is used, respectively.
<code>sigma.log</code>	Logical argument. Per default (<code>sigma.log = TRUE</code>), also in the sigma convergence analysis, the economic variables are transformed by natural logarithm. If the original values should be used, state <code>sigma.log = FALSE</code>
<code>sigma.weighting</code>	If the measure of statistical dispersion in the sigma convergence analysis (coefficient of variation or standard deviation) should be weighted, a weighting vector has to be stated
<code>sigma.issample</code>	Logical argument that indicates if the dataset is a sample or the population (default: <code>is.sample = FALSE</code> , so the denominator of variance is n)
<code>sigma.plot</code>	Logical argument that indicates if a plot of sigma convergence has to be created
<code>sigma.plotLSize</code>	If <code>sigma.plot = TRUE</code> : Line size of the sigma convergence plot
<code>sigma.plotLineCol</code>	If <code>sigma.plot = TRUE</code> : Line color of the sigma convergence plot
<code>sigma.plotRLLine</code>	If <code>sigma.plot = TRUE</code> : Logical argument that indicates if a regression line has to be added to the plot
<code>sigma.plotRLLineCol</code>	If <code>sigma.plot = TRUE</code> and <code>sigma.plotRLLine = TRUE</code> : Color of the regression line
<code>sigma.Ymin</code>	If <code>sigma.plot = TRUE</code> : start value of the Y axis in the plot
<code>sigma.plotX</code>	If <code>sigma.plot = TRUE</code> : Name of the X axis
<code>sigma.plotY</code>	If <code>sigma.plot = TRUE</code> : Name of the Y axis
<code>sigma.plotTitle</code>	If <code>sigma.plot = TRUE</code> : Title of the plot
<code>sigma.bgCol</code>	If <code>sigma.plot = TRUE</code> : Plot background color
<code>sigma.bgrid</code>	If <code>sigma.plot = TRUE</code> : Logical argument that indicates if the plot contains a grid
<code>sigma.bgridCol</code>	If <code>sigma.plot = TRUE</code> and <code>sigma.bgrid = TRUE</code> : Color of the grid
<code>sigma.bgridSize</code>	If <code>sigma.plot = TRUE</code> and <code>sigma.bgrid = TRUE</code> : Size of the grid
<code>sigma.bgridType</code>	If <code>sigma.plot = TRUE</code> and <code>sigma.bgrid = TRUE</code> : Type of the grid

Details

From the regional economic perspective (in particular the neoclassical growth theory), regional disparities are expected to decline. This *convergence* can have different meanings: *Sigma convergence* (σ) means a harmonization of regional economic output or income over time, while *beta convergence* (β) means a decline of dispersion because poor regions have a stronger economic growth than rich regions (Capello/Nijkamp 2009). Regardless of the theoretical assumptions of a harmonization in reality, the related analytical framework allows to analyze both types of convergence for cross-sectional data (GDP p.c. or another economic variable, y , for i regions and two points in time, t and $t + T$), or one starting point (t) and the average growth within the following n years ($t + 1, t + 2, \dots, t + n$), respectively. Beta convergence can be calculated either in a linearized OLS regression model or in a nonlinear regression model. When no other variables are integrated in this model, it is called *absolute* beta convergence. Implementing other region-related variables (conditions) into the model leads to *conditional* beta convergence. If there is beta convergence ($\beta < 0$), it is possible to calculate the *speed of convergence*, λ , and the so-called *Half-Life* H , while the latter is the time taken to reduce the disparities by one half (Allington/McCombie 2007, Goecke/Huether 2016). There is *sigma convergence*, when the dispersion of the variable (σ), e.g. calculated as standard deviation or coefficient of variation, reduces from t to $t + T$. This can be measured using ANOVA for two years or trend regression with respect to several years (Furceri 2005, Goecke/Huether 2016).

The `rca` function is a wrapper for the functions `betaconv.ols`, `betaconv.nls`, `sigmaconv` and `sigmaconv.t`. This function calculates (absolute and/or conditional) beta convergence and sigma convergence. Regional disparities are measured by the standard deviation (or variance, coefficient of variation) for all GDPs per capita (or another economic variable) for the given years. Beta convergence is estimated either using ordinary least squares (OLS) or nonlinear least squares (NLS). If the beta coefficient is negative (using OLS) or positive (using NLS), there is beta convergence. Sigma convergence is analyzed either using an analysis of variance (ANOVA) for these deviation measures (year 1 divided by year 2, F-statistic) or a trend regression (F-statistic, t-statistic). In the former case, if $\sigma_t1/\sigma_t2 > 0$, there is sigma convergence. In the latter case, if the slope of the trend regression is negative, there is sigma convergence.

Value

A list containing the following objects:

<code>betaconv</code>	A list containing the following objects:
<code>regdata</code>	A data frame containing the regression data, including the \ln -transformed economic variables
<code>tinterval</code>	The time interval
<code>abeta</code>	A list containing the estimates of the absolute beta convergence regression model, including lambda and half-life
<code>cbeta</code>	If conditions are stated: a list containing the estimates of the conditional beta convergence regression model, including lambda and half-life
<code>sigmaconv</code>	A list containing the following objects:
<code>sigmaconv</code>	A matrix containing either the standard deviations, their quotient and the results of the significance test (F-statistic) or the results of trend regression

Author(s)

Thomas Wieland

References

Allington, N. F. B./McCombie, J. S. L. (2007): “Economic growth and beta-convergence in the East European Transition Economies”. In: Arestis, P./Baddely, M./McCombie, J. S. L. (eds.): *Economic Growth. New Directions in Theory and Policy*. Cheltenham: Elgar. p. 200-222.

Capello, R./Nijkamp, P. (2009): “Introduction: regional growth and development theories in the twenty-first century - recent theoretical advances and future challenges”. In: Capello, R./Nijkamp, P. (eds.): *Handbook of Regional Growth and Development Theories*. Cheltenham: Elgar. p. 1-16.

Dapena, A. D./Vazquez, E. F./Morollon, F. R. (2016): “The role of spatial scale in regional convergence: the effect of MAUP in the estimation of beta-convergence equations”. In: *The Annals of Regional Science*, **56**, 2, p. 473-489.

Furceri, D. (2005): “Beta and sigma-convergence: A mathematical relation of causality”. In: *Economics Letters*, **89**, 2, p. 212-215.

Goecke, H./Huether, M. (2016): “Regional Convergence in Europe”. In: *Intereconomics*, **51**, 3, p. 165-171.

Young, A. T./Higgins, M. J./Levy, D. (2008): “Sigma Convergence versus Beta Convergence: Evidence from U.S. County-Level Data”. In: *Journal of Money, Credit and Banking*, **40**, 5, p. 1083-1093.

See Also

[betaconv.ols](#), [betaconv.nls](#), [betaconv.speed](#), [sigmaconv](#), [sigmaconv.t](#), [cv](#), [sd2](#), [var2](#)

Examples

```
data (G.counties.gdp)
# Loading GDP data for Germany (counties = Landkreise)

rca (G.counties.gdp$gdppc2010, 2010, G.counties.gdp$gdppc2011, 2011,
conditions = NULL, beta.plot = TRUE)
# Two years, no conditions (Absolute beta convergence)

regionaldummies <- to.dummy(G.counties.gdp$regional)
# Creating dummy variables for West/East
G.counties.gdp$West <- regionaldummies[,2]
G.counties.gdp$East <- regionaldummies[,1]
# Adding dummy variables to data

rca (G.counties.gdp$gdppc2010, 2010, G.counties.gdp$gdppc2011, 2011,
conditions = G.counties.gdp[c(70,71)])
# Two years, with conditions
# (Absolute and conditional beta convergence)

converg1 <- rca (G.counties.gdp$gdppc2010, 2010, G.counties.gdp$gdppc2011, 2011,
conditions = G.counties.gdp[c(70,71)])
```

```

# Store results in object
converg1$betaconv$abeta
# Addressing estimates for the conditional beta model

rca (G.counties.gdp$gdppc2010, 2010, G.counties.gdp[65:68], 2014, conditions = NULL,
sigma.type = "trend", beta.plot = TRUE, sigma.plot = TRUE)
# Five years, no conditions (Absolute beta convergence)
# with plots for both beta and sigma convergence

```

reilly

*Law of retail gravitation by Reilly***Description**

Calculating the proportion of sales from an intermediate town between two cities or retail locations

Usage

```
reilly(P_a, P_b, D_a, D_b, gamma = 1, lambda = 2, relation = NULL)
```

Arguments

P_a	a single numeric value of attractivity/population size of location/city <i>a</i>
P_b	a single numeric value of attractivity/population size of location/city <i>b</i>
D_a	a single numeric value of the distance from the intermediate town to location/city <i>a</i>
D_b	a single numeric value of the distance from the intermediate town to location/city <i>b</i>
gamma	a single numeric value for the exponential weighting of size (default: 1)
lambda	a single numeric value for the exponential weighting of distance (transport costs, default: -2)
relation	a single numeric value containing the relation of trade between cities/locations <i>a</i> and <i>b</i> (only needed if the distance decay parameters has to be estimated instead of the sales flows)

Details

The *law of retail gravitation* by Reilly (1929, 1931) was the first *spatial interaction model* for retailing and services. This "law" states that two cities/locations attract customers from an intermediate town proportionally to the attractivity/population size of the two cities/locations and in inverse proportion to the squares of the transport costs (e.g. distance, travelling time) from these two locations to the intermediate town. But both variables can be weighted by exponents. The distance exponent can also be derived from empirical data (if an empirical `relation` is stated). The *breaking point formula* by Converse (1949) is a separate transformation of Reilly's law (see the function `converse`). The models by Reilly and Converse are simple *spatial interaction models* and are considered as

deterministic market area models due to their exact allocation of demand origins to locations. A probabilistic approach including a theoretical framework was developed by Huff (1962) (see the function `huff`).

Value

If no relation is stated, a list with three values:

<code>relation_AB</code>	relation of trade between cities/locations <i>a</i> and <i>b</i>
<code>prop_A</code>	proportion of city/location <i>a</i>
<code>prop_B</code>	proportion of city/location <i>b</i>

If a relation is stated instead of weighting parameters, a single numeric value containing the estimated distance decay parameter.

Author(s)

Thomas Wieland

References

- Berman, B. R./Evans, J. R. (2012): “Retail Management: A Strategic Approach”. 12th edition. Boston : Pearson.
- Converse, P. D. (1949): “New Laws of Retail Gravitation”. In: *Journal of Marketing*, **14**, 3, p. 379-384.
- Huff, D. L. (1962): “Determination of Intra-Urban Retail Trade Areas”. Los Angeles : University of California.
- Levy, M./Weitz, B. A. (2012): “Retailing management”. 8th edition. New York : McGraw-Hill Irwin.
- Loeffler, G. (1998): “Market areas - a methodological reflection on their boundaries”. In: *GeoJournal*, **45**, 4, p. 265-272
- Reilly, W. J. (1929): “Methods for the Study of Retail Relationships”. *Studies in Marketing*, **4**. Austin : Bureau of Business Research, The University of Texas.
- Reilly, W. J. (1931): “The Law of Retail Gravitation”. New York.

See Also

[huff](#), [converse](#)

Examples

```
# Example from Converse (1949):
reilly (39851, 37366, 27, 25)
# two cities (pop. size 39.851 and 37.366)
# with distances of 27 and 25 miles to intermediate town
myresults <- reilly (39851, 37366, 27, 25)
myresults$prop_A
# proportion of location a
```

```
# Distance decay parameter for the given sales relation:
reilly (39851, 37366, 27, 25, gamma = 1, lambda = NULL, relation = 0.9143555)
# returns 2
```

ripley

Ripley's K

Description

Analyzing point clustering with Ripley's K function

Usage

```
ripley(loc_df, loc_id, loc_lat, loc_lon,
area, t.max, t.sep = 10, K.local = FALSE,
ci.boot = FALSE, ci.alpha = 0.05, ciboot.samples = 100,
progmsg = FALSE, K.plot = TRUE, Kplot.func = "K",
plot.title = "Ripley's K", plotX = "t",
plotY = paste(Kplot.func, "Observed vs. expected"),
lcol.exp = "blue", lcol.emp = "red", lsize.exp = 1,
ltype.exp = "solid", lsize.emp = 1, ltype.emp = "solid",
bg.col = "gray95", bgrid = TRUE, bgrid.col = "white",
bgrid.size = 2, bgrid.type = "solid")
```

Arguments

loc_df	A data frame containing the points
loc_id	Column containing the IDs of the points in the data frame loc_df
loc_lat	Column containing the latitudes of the points in the data frame loc_df
loc_lon	Column containing the longitudes of the points in the data frame loc_df
area	Total area of the regarded region
t.max	Maximum distance
t.sep	Number of distance intervals
K.local	Logical arguments that indicates whether local K values are computed or not
ci.boot	Logical arguments that indicates whether bootstrap confidence intervals are computed or not
ci.alpha	Significance level of the bootstrap confidence intervals
ciboot.samples	No. of bootstrap samples
progmsg	Logical argument: Printing progress messages or not
K.plot	Logical argument: Plot K function or not
Kplot.func	Which function has to be plotted? K function (Kplot.func = "K"), L function (Kplot.func = "L") or H function (Kplot.func = "H")
plot.title	If K.plot = TRUE: Plot title

<code>plotX</code>	If <code>K.plot = TRUE</code> : name of the X axis
<code>plotY</code>	If <code>K.plot = TRUE</code> : name of the Y axis
<code>lcol.exp</code>	If <code>K.plot = TRUE</code> : color of the line representing the expected values
<code>lcol.emp</code>	If <code>K.plot = TRUE</code> : color of the line representing the empirical values
<code>lsize.exp</code>	If <code>K.plot = TRUE</code> : size of the line representing the expected values
<code>lsize.emp</code>	If <code>K.plot = TRUE</code> : size of the line representing the empirical values
<code>ltype.exp</code>	If <code>K.plot = TRUE</code> : type of the line representing the expected values
<code>ltype.emp</code>	If <code>K.plot = TRUE</code> : type of the line representing the empirical values
<code>bg.col</code>	if <code>lc = TRUE</code> (plot of Lorenz curve), <code>bg.col</code> defines the background color of the plot (default: "gray95")
<code>bgrid</code>	if <code>lc = TRUE</code> (plot of Lorenz curve), the logical argument <code>bgrid</code> defines if a grid is shown in the plot
<code>bgrid.col</code>	if <code>lc = TRUE</code> (plot of Lorenz curve) and <code>bgrid = TRUE</code> (background grid), <code>bgrid.col</code> defines the color of the background grid (default: "white")
<code>bgrid.size</code>	if <code>lc = TRUE</code> (plot of Lorenz curve) and <code>bgrid = TRUE</code> (background grid), <code>bgrid.size</code> defines the size of the background grid (default: 2)
<code>bgrid.type</code>	if <code>lc = TRUE</code> (plot of Lorenz curve) and <code>bgrid = TRUE</code> (background grid), <code>bgrid.type</code> defines the type of lines of the background grid (default: "solid")

Details

Calculating and plotting of the K function and its derivations (L function, H function) and, optionally, bootstrap confidence intervals.

Value

The function returns a list containing:

<code>K</code>	A data.frame containing the K/L/H/t values
<code>K_local</code>	A data.frame containing the local K values (if stated)
<code>local_ci</code>	A data.frame containing the local confidence intervals (if stated)

Author(s)

Thomas Wieland

References

- Kiskowski, M.A./Hancock, J. F./Kenworthy, A. (2009): "On the Use of Ripley's K-function and its Derivatives to Analyze Domain Size". In: *Biophysical Journal*, **97**, 4, p. 1095-1103.
- Krider, R. E./Putler, R. S. (2013): "Which Birds of a Feather Flock Together? Clustering and Avoidance Patterns of Similar Retail Outlets". In: *Geographical Analysis*, **45**, 2, p. 123-149.

See Also

[dist](#), [dist.buf](#), [dist.mat](#)

Examples

```
## Not run:
data(GoettingenHealth1)
# general practitioners, psychotherapists and pharmacies

area_goe <- 1753000000
# area of Landkreis Goettingen (sqm)
area_nom <- 1267000000
# area of Landkreis Northeim (sqm)
area_gn <- area_goe+area_nom
sqrt(area_gn/pi)

# this takes some seconds
ripley(GoettingenHealth1[GoettingenHealth1$type == "phys_gen",],
"location", "lat", "lon", area = area_gn, t.max = 30000, t.sep = 300)

ripley(GoettingenHealth1[GoettingenHealth1$type == "pharm",],
"location", "lat", "lon", area = area_gn, t.max = 30000, t.sep = 300)

ripley(GoettingenHealth1[GoettingenHealth1$type == "psych",],
"location", "lat", "lon", area = area_gn, t.max = 30000, t.sep = 300)

## End(Not run)
```

sd2

Standard deviation (extended)

Description

Calculating the standard deviation (sd), weighted or non-weighted, for samples or populations

Usage

```
sd2 (x, is.sample = TRUE, weighting = NULL, wmean = FALSE, na.rm = TRUE)
```

Arguments

x	a numeric vector
is.sample	logical argument that indicates if the dataset is a sample or the population (default: is.sample = TRUE, so the denominator of variance is $n - 1$)
weighting	a numeric vector containing weighting data to compute the weighted standard deviation (instead of the non-weighted sd)
wmean	logical argument that indicates if the weighted mean is used when calculating the weighted standard deviation
na.rm	logical argument that whether NA values should be extracted or not

Details

The function calculates the *standard deviation*. Unlike the R base `sd` function, the `sd2` function allows to choose if the data is treated as sample (denominator of variance is $n - 1$) or not (denominator of variance is n)

From a regional economic perspective, the `sd` is closely linked to the concept of *sigma convergence* (σ) which means a harmonization of regional economic output or income over time, while the other type of convergence, *beta convergence* (β), means a decline of dispersion because poor regions have a stronger growth than rich regions (Capello/Nijkamp 2009). The `sd` allows to summarize regional disparities (e.g. disparities in regional GDP per capita) in one indicator. The coefficient of variation (see the function `cv`) is more frequently used for this purpose (e.g. Lessmann 2005, Huang/Leung 2009, Siljak 2015). But the `sd` can also be used for any other types of disparities or dispersion, such as disparities in supply (e.g. density of physicians or grocery stores).

The standard deviation can be weighted by using a second weighting vector. As there is more than one way to weight measures of statistical dispersion, this function uses the formula for the weighted `sd` (σ_w) from Sheret (1984). The vector `x` is automatically treated as a sample (such as in the base `sd` function), so the denominator of variance is $n - 1$, if it is not, set `is.sample = FALSE`.

Value

Single numeric value. If `weighting` is specified, the function returns a weighted standard deviation (optionally using a weighted arithmetic mean if `wmean = TRUE`).

Author(s)

Thomas Wieland

References

- Bahrenberg, G./Giese, E./Mevenkamp, N./Nipper, J. (2010): "Statistische Methoden in der Geographie. Band 1: Univariate und bivariate Statistik". Stuttgart: Borntraeger.
- Capello, R./Nijkamp, P. (2009): "Introduction: regional growth and development theories in the twenty-first century - recent theoretical advances and future challenges". In: Capello, R./Nijkamp, P. (eds.): *Handbook of Regional Growth and Development Theories*. Cheltenham: Elgar. p. 1-16.
- Lessmann, C. (2005): "Regionale Disparitaeten in Deutschland und ausgesuchten OECD-Staaten im Vergleich". *ifo Dresden berichtet*, **3/2005**. https://www.ifo.de/DocDL/ifoedb_2005_3_25-33.pdf.
- Huang, Y./Leung, Y. (2009): "Measuring Regional Inequality: A Comparison of Coefficient of Variation and Hoover Concentration Index". In: *The Open Geography Journal*, **2**, p. 25-34.
- Sheret, M. (1984): "The Coefficient of Variation: Weighting Considerations". In: *Social Indicators Research*, **15**, 3, p. 289-295.
- Siljak, D. (2015): "Real Economic Convergence in Western Europe from 1995 to 2013". In: *International Journal of Business and Economic Development*, **3**, 3, p. 56-67.

See Also

[gini](#), [herf](#), [hoover](#), [mean2](#), [rca](#)

Examples

```
# Regional disparities / sigma convergence in Germany
data(G.counties.gdp)
# GDP per capita for German counties (Landkreise)
sd_gdppc <- apply (G.counties.gdp[54:68], MARGIN = 2, FUN = sd2)
# Calculating standard deviation for the years 2000-2014
years <- 2000:2014
# vector of years (2000-2014)
plot(years, sd_gdppc, "l", ylim = c(0,15000), xlab = "Year",
ylab = "SD of GDP per capita")
# Plot sd over time
```

 shift

Shift-share analysis

Description

Analyzing regional growth with the shift-share analysis

Usage

```
shift(e_ij1, e_ij2, e_i1, e_i2, industry.names = NULL,
shift.method = "Dunn", print.results = TRUE, plot.results = FALSE,
plot.colours = NULL, plot.title = NULL, plot.portfolio = FALSE, ...)
```

Arguments

e_ij1	a numeric vector with i values containing the employment in i industries in region j at time 1
e_ij2	a numeric vector with i values containing the employment in i industries in region j at time 2
e_i1	a numeric vector with i values containing the total employment in i industries at time 1
e_i2	a numeric vector with i values containing the total employment in i industries at time 2
industry.names	Industry names (e.g. from the relevant statistical classification of economic activities)
shift.method	Method of shift-share-analysis to be used ("Dunn", "Esteban", "Gerfin") (default: shift.method = "Dunn")
print.results	Logical argument that indicates if the function shows the results or not
plot.results	Logical argument that indicates if the results have to be plotted
plot.colours	If plot.results = TRUE: Plot colours
plot.title	If plot.results = TRUE: Plot title
plot.portfolio	Logical argument that indicates if the results have to be plotted in a portfolio matrix additionally
...	Additional arguments for the portfolio plot (see the function portfolio)

Details

The *shift-share analysis* (Dunn 1960) addresses the regional growth (or decline) regarding the overall development in the national economy. The aim of this analysis model is to identify which parts of the regional economic development can be traced back to national trends, effects of the regional industry structure and (positive) regional factors. The growth (or decline) of regional employment consists of three factors: $l_{t+1} - l_t = nps + nds + nts$, where l is the employment in the region at time t and $t + 1$, respectively, and nps is the *net proportionality shift*, nds is the *net differential shift* and nts is the *net total shift*. Other variants are e.g. the shift-share method by Gerfin (Index method), the dynamic shift-share analysis (Barff/Knight 1988) or the extension by Esteban-Marquillas (1972).

As there is more than one way to calculate a Dunn-type *shift-share analysis* and the terms are not used consequently in the regional economic literature, this function and the documentation use the formulae and terms given in Farhauer/Kroell (2013). If `shift.method = "Dunn"`, this function calculates the *net proportionality shift* (nps), the *net differential shift* (nds) and the *net total shift* (nts) where the last one represents the residuum of (positive) regional factors.

This function calculates a shift-share analysis for two years.

Value

A list containing the following objects:

components	A matrix containing the shift-share components related to the chosen method
growth	A matrix containing the industry-specific growth values
method	The chosen method, e.g. "Dunn"

Author(s)

Thomas Wieland

References

- Arcelus, F. J. (1984): "An Extension of Shift-Share Analysis". In: *Growth and Change*, **15**, 1, p. 3-8.
- Barff, R. A./Knight, P. L. (1988): "Dynamic Shift-Share Analysis". In: *Growth and Change*, **19**, 2, p. 1-10.
- Casler, S. D. (1989): "A Theoretical Context for Shift and Share Analysis". In: *Regional Studies*, **23**, 1, p. 43-48.
- Dunn, E. S. Jr. (1960): "A statistical and analytical technique for regional analysis". In: *Papers and Proceedings of the Regional Science Association*, **6**, p. 97-112.
- Esteban-Marquillas, J. M. (1972): "Shift- and share analysis revisited". In: *Regional and Urban Economics*, **2**, 3, p. 249-261.
- Farhauer, O./Kroell, A. (2013): "Standorttheorien: Regional- und Stadtoekonomik in Theorie und Praxis". Wiesbaden : Springer.
- Gerfin, H. (1964): "Gesamtwirtschaftliches Wachstum und regionale Entwicklung". In: *Kyklos*, **17**, 4, p. 565-593.
- Schoenebeck, C. (1996): "Wirtschaftsstruktur und Regionalentwicklung: Theoretische und empirische Befunde fuer die Bundesrepublik Deutschland". *Dortmunder Beitrage zur Raumplanung*, **75**. Dortmund.

See Also

[portfolio](#), [shiftd](#), [shifti](#), [shift.growth](#)

Examples

```
# Example from Farhauer/Kroell (2013):
region_A_t <- c(90,20,10,60)
region_A_t1 <- c(100,40,10,55)
# data for region A (time t and t+1)
nation_X_t <- c(400,150,150,400)
nation_X_t1 <- c(440,210,135,480)
# data for the national economy (time t and t+1)
resultsA <- shift(region_A_t, region_A_t1, nation_X_t, nation_X_t1)
# results for region A
region_B_t <- c(60,30,30,40)
region_B_t1 <- c(85,55,40,35)
# data for region B (time t and t+1)
resultsB <- shift(region_B_t, region_B_t1, nation_X_t, nation_X_t1)
# results for region B
region_C_t <- c(250,100,110,300)
region_C_t1 <- c(255,115,85,390)
# data for region C (time t and t+1)
resultsC <- shift(region_C_t, region_C_t1, nation_X_t, nation_X_t1)
# results for region C

# Example Freiburg dataset
data(Freiburg)
# Loads the data
shift(Freiburg$e_Freiburg2008, Freiburg$e_Freiburg2014, Freiburg$e_Germany2008,
Freiburg$e_Germany2014)
# results for Freiburg and Germany (2008 vs. 2014)
```

shift.growth

Growth rates for shift-share analysis

Description

This function calculates industry-specific growth rates which are part of the shift-share analysis

Usage

```
shift.growth(e_ij1, e_ij2, e_i1, e_i2, time.periods = NULL,
industry.names = NULL)
```

Arguments

`e_ij1` a numeric vector with i values containing the employment in i industries in region j at time 1

e_ij2	a numeric vector with i values containing the employment in i industries in region j at time 2
e_i1	a numeric vector with i values containing the total employment in i industries at time 1
e_i2	a numeric vector with i values containing the total employment in i industries at time 2
time.periods	No. of regarded time periods (for average growth rates)
industry.names	Industry names (e.g. from the relevant statistical classification of economic activities)

Details

The *shift-share analysis* (Dunn 1960) addresses the regional growth (or decline) regarding the overall development in the national economy. The aim of this analysis model is to identify which parts of the regional economic development can be traced back to national trends, effects of the regional industry structure and (positive) regional factors. The growth (or decline) of regional employment consists of three factors: $l_{t+1} - l_t = nps + nds + nts$, where l is the employment in the region at time t and $t + 1$, respectively, and nps is the *net proportionality shift*, nds is the *net differential shift* and nts is the *net total shift*. Other variants are e.g. the shift-share method by Gerfin (Index method) and the dynamic shift-share analysis (Barff/Knight 1988).

As there is more than one way to calculate a Dunn-type *shift-share analysis* and the terms are not used consequently in the regional economic literature, this function and the documentation use the formulae and terms given in Farhauer/Kroell (2013). If `shift.method = "Dunn"`, this function calculates the *net proportionality shift* (nps), the *net differential shift* (nds) and the *net total shift* (nts) where the last one represents the residuum of (positive) regional factors.

This function calculates industry-specific growth rates which are part of a shift-share analysis.

Value

A matrix containing the industry-specific growth values

Author(s)

Thomas Wieland

References

- Arcelus, F. J. (1984): "An Extension of Shift-Share Analysis". In: *Growth and Change*, **15**, 1, p. 3-8.
- Barff, R. A./Knight, P. L. (1988): "Dynamic Shift-Share Analysis". In: *Growth and Change*, **19**, 2, p. 1-10.
- Casler, S. D. (1989): "A Theoretical Context for Shift and Share Analysis". In: *Regional Studies*, **23**, 1, p. 43-48.
- Dunn, E. S. Jr. (1960): "A statistical and analytical technique for regional analysis". In: *Papers and Proceedings of the Regional Science Association*, **6**, p. 97-112.
- Esteban-Marquillas, J. M. (1972): "Shift- and share analysis revisited". In: *Regional and Urban Economics*, **2**, 3, p. 249-261.

Farhauer, O./Kroell, A. (2013): “Standorttheorien: Regional- und Stadtoekonomik in Theorie und Praxis”. Wiesbaden : Springer.

Gerfin, H. (1964): “Gesamtwirtschaftliches Wachstum und regionale Entwicklung”. In: *Kyklos*, **17**, 4, p. 565-593.

Goschin, Z. (2014): “Regional growth in Romania after its accession to EU: a shift-share analysis approach”. In: *Procedia Economics and Finance*, **15**, p. 169-175.

Schoenebeck, C. (1996): “Wirtschaftsstruktur und Regionalentwicklung: Theoretische und empirische Befunde fuer die Bundesrepublik Deutschland”. *Dortmunder Beitrage zur Raumplanung*, **75**. Dortmund.

See Also

[portfolio](#), [shift](#), [shiftd](#), [shifti](#)

Examples

```
# Example from Farhauer/Kroell (2013):
region_A_t <- c(90,20,10,60)
region_A_t1 <- c(100,40,10,55)
# data for region A (time t and t+1)
nation_X_t <- c(400,150,150,400)
nation_X_t1 <- c(440,210,135,480)
# data for the national economy (time t and t+1)
shiftd.growth(region_A_t, region_A_t1, nation_X_t, nation_X_t1)
```

shiftd

Dynamic shift-share analysis

Description

Analyzing regional growth with the dynamic shift-share analysis

Usage

```
shiftd(e_ij1, e_ij2, e_i1, e_i2, time1, time2,
industry.names = NULL, shift.method = "Dunn",
gerfin.shifts = "mean", print.results = TRUE,
plot.results = FALSE, plot.colours = NULL, plot.title = NULL,
plot.portfolio = FALSE, ...)
```

Arguments

e_ij1 a numeric vector with i values containing the employment in i industries in region j at time 1

e_ij2 a numeric data frame or matrix with i rows containing the employment in i industries in region j and t columns, representing t ($t > 1$) years

e_i1	a numeric vector with i values containing the total employment in i industries at time 1
e_i2	a numeric data frame or matrix with i rows containing the total employment in i industries and t columns, representing t ($t > 1$) years
time1	Initial year
time2	Final year
industry.names	Industry names (e.g. from the relevant statistical classification of economic activities)
shift.method	Method of shift-share-analysis to be used ("Dunn", "Gerfin") (default: shift.method = "Dunn")
gerfin.shifts	If shift.method = "Gerfin": Logical argument that indicates if the shifts are calculated as sums or as means (default: gerfin = "mean")
print.results	Logical argument that indicates if the function shows the results or not
plot.results	Logical argument that indicates if the results have to be plotted
plot.colours	If plot.results = TRUE: Plot colours
plot.title	If plot.results = TRUE: Plot title
plot.portfolio	Logical argument that indicates if the results have to be plotted in a portfolio matrix additionally
...	Additional arguments for the portfolio plot (see the function portfolio)

Details

The *shift-share analysis* (Dunn 1960) addresses the regional growth (or decline) regarding the overall development in the national economy. The aim of this analysis model is to identify which parts of the regional economic development can be traced back to national trends, effects of the regional industry structure and (positive) regional factors. The growth (or decline) of regional employment consists of three factors: $l_{t+1} - l_t = nps + nds + nts$, where l is the employment in the region at time t and $t + 1$, respectively, and nps is the *net proportionality shift*, nds is the *net differential shift* and nts is the *net total shift*. Other variants are e.g. the shift-share method by Gerfin (Index method) and the dynamic shift-share analysis (Barff/Knight 1988).

As there is more than one way to calculate a Dunn-type *shift-share analysis* and the terms are not used consequently in the regional economic literature, this function and the documentation use the formulae and terms given in Farhauer/Kroell (2013). If shift.method = "Dunn", this function calculates the *net proportionality shift* (nps), the *net differential shift* (nds) and the *net total shift* (nts) where the last one represents the residuum of (positive) regional factors.

This function calculates a dynamic shift-share analysis for at least two years.

Value

A list containing the following objects:

components	A matrix containing the shift-share components related to the chosen method
components.year	A matrix containing the shift-share components for each year
growth	A matrix containing the industry-specific growth values
method	The chosen method, e.g. "Dunn"

Author(s)

Thomas Wieland

References

- Arcelus, F. J. (1984): "An Extension of Shift-Share Analysis". In: *Growth and Change*, **15**, 1, p. 3-8.
- Barff, R. A./Knight, P. L. (1988): "Dynamic Shift-Share Analysis". In: *Growth and Change*, **19**, 2, p. 1-10.
- Casler, S. D. (1989): "A Theoretical Context for Shift and Share Analysis". In: *Regional Studies*, **23**, 1, p. 43-48.
- Dunn, E. S. Jr. (1960): "A statistical and analytical technique for regional analysis". In: *Papers and Proceedings of the Regional Science Association*, **6**, p. 97-112.
- Esteban-Marquillas, J. M. (1972): "Shift- and share analysis revisited". In: *Regional and Urban Economics*, **2**, 3, p. 249-261.
- Farhauer, O./Kroell, A. (2013): "Standorttheorien: Regional- und Stadtoekonomie in Theorie und Praxis". Wiesbaden : Springer.
- Gerfin, H. (1964): "Gesamtwirtschaftliches Wachstum und regionale Entwicklung". In: *Kyklos*, **17**, 4, p. 565-593.
- Schoenebeck, C. (1996): "Wirtschaftsstruktur und Regionalentwicklung: Theoretische und empirische Befunde fuer die Bundesrepublik Deutschland". *Dortmunder Beitrage zur Raumplanung*, **75**. Dortmund.

See Also

[portfolio](#), [shift](#), [shifti](#), [shift.growth](#)

Examples

```
# Example from Farhauer/Kroell (2013), extended:
region_A_t <- c(90,20,10,60)
region_A_t1 <- c(100,40,10,55)
region_A_t2 <- c(105,45,15,60)
# data for region A (time t and t+1)
nation_X_t <- c(400,150,150,400)
nation_X_t1 <- c(440,210,135,480)
nation_X_t2 <- c(460,230,155,500)
# data for the national economy (time t and t+1)
shiftd(region_A_t, data.frame(region_A_t1, region_A_t2), nation_X_t,
data.frame(nation_X_t1, nation_X_t2), time1 = 2000, time2 = 2002,
plot.results = TRUE, plot.portfolio = TRUE, psize = region_A_t1)

data(Goettingen)
shiftd(Goettingen$Goettingen2008[2:16], Goettingen[2:16,3:11],
Goettingen$BRD2008[2:16], Goettingen[2:16,13:21],
time1 = 2008, time2 = 2017, industry.names = Goettingen$WA_WZ2008[2:16],
shift.method = "Dunn")
```

shifti *Shift-share analysis for industries*

Description

Analyzing industry-specific regional growth with the shift-share analysis

Usage

```
shifti(e_ij1, e_ij2, e_i1, e_i2, industry.names = NULL,
       shift.method = "Dunn", print.results = TRUE, plot.results = FALSE,
       plot.colours = NULL, plot.title = NULL, plot.portfolio = FALSE, ...)
```

Arguments

e_ij1	a numeric vector with i values containing the employment in i industries in region j at time 1
e_ij2	a numeric vector with i values containing the employment in i industries in region j at time 2
e_i1	a numeric vector with i values containing the total employment in i industries at time 1
e_i2	a numeric vector with i values containing the total employment in i industries at time 2
industry.names	Industry names (e.g. from the relevant statistical classification of economic activities)
shift.method	Method of shift-share-analysis to be used ("Dunn", "Gerfin") (default: shift.method = "Dunn")
print.results	Logical argument that indicates if the function shows the results or not
plot.results	Logical argument that indicates if the results have to be plotted
plot.colours	If plot.results = TRUE: Plot colours
plot.title	If plot.results = TRUE: Plot title
plot.portfolio	Logical argument that indicates if the results have to be plotted in a portfolio matrix additionally
...	Additional arguments for the portfolio plot (see the function portfolio)

Details

The *shift-share analysis* (Dunn 1960) addresses the regional growth (or decline) regarding the overall development in the national economy. The aim of this analysis model is to identify which parts of the regional economic development can be traced back to national trends, effects of the regional industry structure and (positive) regional factors. The growth (or decline) of regional employment consists of three factors: $l_{t+1} - l_t = nps + nds + nts$, where l is the employment in the region at time t and $t + 1$, respectively, and nps is the *net proportionality shift*, nds is the *net differential*

shift and *nts* is the *net total shift*. Other variants are e.g. the shift-share method by Gerfin (Index method) and the dynamic shift-share analysis (Barff/Knight 1988).

As there is more than one way to calculate a Dunn-type *shift-share analysis* and the terms are not used consequently in the regional economic literature, this function and the documentation use the formulae and terms given in Farhauer/Kroell (2013). If `shift.method = "Dunn"`, this function calculates the *net proportionality shift* (*nps*), the *net differential shift* (*nds*) and the *net total shift* (*nts*) where the last one represents the residuum of (positive) regional factors.

This function calculates a shift-share analysis for at least two years and results industry-specific shift-share components.

Value

A list containing the following objects:

<code>components</code>	A matrix containing the shift-share components related to the chosen method
<code>components.industry</code>	A matrix containing the shift-share components for each industry
<code>growth</code>	A matrix containing the industry-specific growth values
<code>method</code>	The chosen method, e.g. "Dunn"

Author(s)

Thomas Wieland

References

- Arcelus, F. J. (1984): "An Extension of Shift-Share Analysis". In: *Growth and Change*, **15**, 1, p. 3-8.
- Barff, R. A./Knight, P. L. (1988): "Dynamic Shift-Share Analysis". In: *Growth and Change*, **19**, 2, p. 1-10.
- Casler, S. D. (1989): "A Theoretical Context for Shift and Share Analysis". In: *Regional Studies*, **23**, 1, p. 43-48.
- Dunn, E. S. Jr. (1960): "A statistical and analytical technique for regional analysis". In: *Papers and Proceedings of the Regional Science Association*, **6**, p. 97-112.
- Esteban-Marquillas, J. M. (1972): "Shift- and share analysis revisited". In: *Regional and Urban Economics*, **2**, 3, p. 249-261.
- Farhauer, O./Kroell, A. (2013): "Standorttheorien: Regional- und Stadtoekonomik in Theorie und Praxis". Wiesbaden : Springer.
- Gerfin, H. (1964): "Gesamtwirtschaftliches Wachstum und regionale Entwicklung". In: *Kyklos*, **17**, 4, p. 565-593.
- Schoenebeck, C. (1996): "Wirtschaftsstruktur und Regionalentwicklung: Theoretische und empirische Befunde fuer die Bundesrepublik Deutschland". *Dortmunder Beitrage zur Raumplanung*, **75**. Dortmund.

See Also

[portfolio](#), [shift](#), [shifti](#), [shift.growth](#)

Examples

```
# Example from Farhauer/Kroell (2013):
region_A_t <- c(90,20,10,60)
region_A_t1 <- c(100,40,10,55)
# data for region A (time t and t+1)
nation_X_t <- c(400,150,150,400)
nation_X_t1 <- c(440,210,135,480)
# data for the national economy (time t and t+1)
shiftid(region_A_t, region_A_t1, nation_X_t, nation_X_t1,
plot.results = TRUE, plot.portfolio = TRUE, psize = region_A_t1)
```

 shiftid

Dynamic shift-share analysis for industries

Description

Analyzing industry-specific regional growth with the dynamic shift-share analysis

Usage

```
shiftid(e_ij1, e_ij2, e_i1, e_i2, time1, time2,
industry.names = NULL, shift.method = "Dunn",
gerfin.shifts = "mean", print.results = TRUE,
plot.results = FALSE, plot.colours = NULL, plot.title = NULL,
plot.portfolio = FALSE, ...)
```

Arguments

e_ij1	a numeric vector with i values containing the employment in i industries in region j at time 1
e_ij2	a numeric data frame or matrix with i rows containing the employment in i industries in region j and t columns, representing t ($t > 1$) years
e_i1	a numeric vector with i values containing the total employment in i industries at time 1
e_i2	a numeric data frame or matrix with i rows containing the total employment in i industries and t columns, representing t ($t > 1$) years
time1	Initial year
time2	Final year
industry.names	Industry names (e.g. from the relevant statistical classification of economic activities)
shift.method	Method of shift-share-analysis to be used ("Dunn", "Gerfin") (default: shift.method = "Dunn")
gerfin.shifts	If shift.method = "Gerfin": Logical argument that indicates if the shifts are calculated as sums or as means (default: gerfin = "mean")
print.results	Logical argument that indicates if the function shows the results or not

plot.results	Logical argument that indicates if the results have to be plotted
plot.colours	If plot.results = TRUE: Plot colours
plot.title	If plot.results = TRUE: Plot title
plot.portfolio	Logical argument that indicates if the results have to be plotted in a portfolio matrix additionally
...	Additional arguments for the portfolio plot (see the function portfolio)

Details

The *shift-share analysis* (Dunn 1960) addresses the regional growth (or decline) regarding the overall development in the national economy. The aim of this analysis model is to identify which parts of the regional economic development can be traced back to national trends, effects of the regional industry structure and (positive) regional factors. The growth (or decline) of regional employment consists of three factors: $l_{t+1} - l_t = nps + nds + nts$, where l is the employment in the region at time t and $t + 1$, respectively, and nps is the *net proportionality shift*, nds is the *net differential shift* and nts is the *net total shift*. Other variants are e.g. the shift-share method by Gerfin (Index method) and the dynamic shift-share analysis (Barff/Knight 1988).

As there is more than one way to calculate a Dunn-type *shift-share analysis* and the terms are not used consequently in the regional economic literature, this function and the documentation use the formulae and terms given in Farhauer/Kroell (2013). If `shift.method = "Dunn"`, this function calculates the *net proportionality shift* (nps), the *net differential shift* (nds) and the *net total shift* (nts) where the last one represents the residuum of (positive) regional factors.

This function calculates a dynamic shift-share analysis for at least two years.

Value

A list containing the following objects:

components	A matrix containing the shift-share components related to the chosen method
components.year	A matrix containing the shift-share components for each year
growth	A matrix containing the industry-specific growth values
method	The chosen method, e.g. "Dunn"

Author(s)

Thomas Wieland

References

- Arcelus, F. J. (1984): "An Extension of Shift-Share Analysis". In: *Growth and Change*, **15**, 1, p. 3-8.
- Barff, R. A./Knight, P. L. (1988): "Dynamic Shift-Share Analysis". In: *Growth and Change*, **19**, 2, p. 1-10.
- Casler, S. D. (1989): "A Theoretical Context for Shift and Share Analysis". In: *Regional Studies*, **23**, 1, p. 43-48.

Dunn, E. S. Jr. (1960): "A statistical and analytical technique for regional analysis". In: *Papers and Proceedings of the Regional Science Association*, **6**, p. 97-112.

Esteban-Marquillas, J. M. (1972): "Shift- and share analysis revisited". In: *Regional and Urban Economics*, **2**, 3, p. 249-261.

Farhauer, O./Kroell, A. (2013): "Standorttheorien: Regional- und Stadtoekonomie in Theorie und Praxis". Wiesbaden : Springer.

Gerfin, H. (1964): "Gesamtwirtschaftliches Wachstum und regionale Entwicklung". In: *Kyklos*, **17**, 4, p. 565-593.

Schoenebeck, C. (1996): "Wirtschaftsstruktur und Regionalentwicklung: Theoretische und empirische Befunde fuer die Bundesrepublik Deutschland". *Dortmunder Beitrage zur Raumplanung*, **75**. Dortmund.

See Also

[portfolio](#), [shift](#), [shifti](#), [shift.growth](#)

Examples

```
# Example from Farhauer/Kroell (2013), extended:
region_A_t <- c(90,20,10,60)
region_A_t1 <- c(100,40,10,55)
region_A_t2 <- c(105,45,15,60)
# data for region A (time t and t+1)
nation_X_t <- c(400,150,150,400)
nation_X_t1 <- c(440,210,135,480)
nation_X_t2 <- c(460,230,155,500)
# data for the national economy (time t and t+1)
shiftd(region_A_t, data.frame(region_A_t1, region_A_t2), nation_X_t,
data.frame(nation_X_t1, nation_X_t2), time1 = 2000, time2 = 2002,
plot.results = TRUE, plot.portfolio = TRUE, psize = region_A_t1)

data(Goettingen)
shiftd(Goettingen$Goettingen2008[2:16], Goettingen[2:16,3:11],
Goettingen$BRD2008[2:16], Goettingen[2:16,13:21],
time1 = 2008, time2 = 2017, industry.names = Goettingen$WA_WZ2008[2:16],
shift.method = "Dunn")
```

shiftp

Shift-share prognosis

Description

Forecasting regional employment growth with the shift-share analysis (Gerfin model)

Usage

```
shiftp(e_ij1, e_ij2, e_i1, e_i2, e_i3, time1, time2, time3,
       industry.names = NULL, print.results = TRUE,
       plot.results = FALSE, plot.colours = NULL, plot.title = NULL,
       plot.portfolio = FALSE, ...)
```

Arguments

e_ij1	a numeric vector with i values containing the employment in i industries in region j at time 1
e_ij2	a numeric vector with i values containing the employment in i industries in region j at time 2
e_i1	a numeric vector with i values containing the total employment in i industries at time 1
e_i2	a numeric vector with i values containing the total employment in i industries at time 2
e_i3	a numeric vector with i values containing the total employment in i industries at time 3 (forecast value for total employment)
time1	start year (single value)
time2	end year of <i>empirical</i> employment data (single value)
time3	year of prognosis (single value)
industry.names	Industry names (e.g. from the relevant statistical classification of economic activities)
print.results	Logical argument that indicates if the function shows the results or not
plot.results	Logical argument that indicates if the results have to be plotted
plot.colours	If plot.results = TRUE: Plot colours
plot.title	If plot.results = TRUE: Plot title
plot.portfolio	Logical argument that indicates if the results have to be plotted in a portfolio matrix additionally
...	Additional arguments for the portfolio plot (see the function portfolio)

Details

The *shift-share analysis* (Dunn 1960) addresses the regional growth (or decline) regarding the overall development in the national economy. The aim of this analysis model is to identify which parts of the regional economic development can be traced back to national trends, effects of the regional industry structure and (positive) regional factors. The growth (or decline) of regional employment consists of three factors: $l_{t+1} - l_t = nps + nds + nts$, where l is the employment in the region at time t and $t + 1$, respectively, and nps is the *net proportionality shift*, nds is the *net differential shift* and nts is the *net total shift*. Other variants are e.g. the shift-share method by Gerfin (Index method), the dynamic shift-share analysis (Barff/Knight 1988) or the extension by Esteban-Marquillas (1972).

As there is more than one way to calculate a Dunn-type *shift-share analysis* and the terms are not used consequently in the regional economic literature, this function and the documentation use the formulae and terms given in Farhauer/Kroell (2013). If `shift.method = "Dunn"`, this function

calculates the *net proportionality shift* (*nps*), the *net differential shift* (*nds*) and the *net total shift* (*nts*) where the last one represents the residuum of (positive) regional factors.

This function calculates an employment prognosis based on a Gerfin shift-share analysis for two years.

Value

A list containing the following objects:

components	A matrix containing the shift-share components related to the chosen method
growth	A matrix containing the industry-specific growth values
prog	A matrix containing the industry-specific prognosis values
method	The chosen method, e.g. "Dunn"

Author(s)

Thomas Wieland

References

- Arcelus, F. J. (1984): "An Extension of Shift-Share Analysis". In: *Growth and Change*, **15**, 1, p. 3-8.
- Barff, R. A./Knight, P. L. (1988): "Dynamic Shift-Share Analysis". In: *Growth and Change*, **19**, 2, p. 1-10.
- Casler, S. D. (1989): "A Theoretical Context for Shift and Share Analysis". In: *Regional Studies*, **23**, 1, p. 43-48.
- Dunn, E. S. Jr. (1960): "A statistical and analytical technique for regional analysis". In: *Papers and Proceedings of the Regional Science Association*, **6**, p. 97-112.
- Esteban-Marquillas, J. M. (1972): "Shift- and share analysis revisited". In: *Regional and Urban Economics*, **2**, 3, p. 249-261.
- Farhauer, O./Kroell, A. (2013): "Standorttheorien: Regional- und Stadtoekonomik in Theorie und Praxis". Wiesbaden : Springer.
- Gerfin, H. (1964): "Gesamtwirtschaftliches Wachstum und regionale Entwicklung". In: *Kyklos*, **17**, 4, p. 565-593.
- Schoenebeck, C. (1996): "Wirtschaftsstruktur und Regionalentwicklung: Theoretische und empirische Befunde fuer die Bundesrepublik Deutschland". *Dortmunder Beitrage zur Raumplanung*, **75**. Dortmund.
- Spiekermann, K./Wegener, M. (2008): "Modelle in der Raumplanung I. 4 - Input-Output-Modelle". Power Point presentation. http://www.spiekermann-wegener.de/mir/pdf/MIR1_4_111108.pdf.

See Also

[portfolio](#), [shiftd](#), [shifti](#), [shift.growth](#)

Examples

```

# Example data from Spiekermann/Wegener 2008:
# two regions, two industries
region1_2000 <- c(1400, 3600)
region1_2006 <- c(1000, 4400)
region2_2000 <- c(1200, 1800)
region2_2006 <- c(1100, 3700)
region3_2000 <- c(1100, 900)
region3_2006 <- c(800, 1000)
# regional values
nation_2000 <- c(3700, 6300)
nation_2006 <- c(2900, 9100)
# national values
nation_2010 <- c(2500, 12500)
# national prognosis values

# Analysis for region 1:
shiftp(region1_2000, region1_2006, nation_2000,
nation_2006, e_i3 = nation_2010,
time1 = 2000, time2 = 2006, time3 = 2010)
# Analysis for region 2:
shiftp(region2_2000, region2_2006, nation_2000,
nation_2006, e_i3 = nation_2010,
time1 = 2000, time2 = 2006, time3 = 2010)
# Analysis for region 3:
shiftp(region3_2000, region3_2006, nation_2000,
nation_2006, e_i3 = nation_2010,
time1 = 2000, time2 = 2006, time3 = 2010)

```

sigmaconv

Analysis of regional sigma convergence for two years using ANOVA

Description

This function provides the analysis of regional economic sigma convergence (decline of deviation) for two years using ANOVA (Analysis of Variance)

Usage

```

sigmaconv(gdp1, time1, gdp2, time2, sigma.measure = "sd",
sigma.log = TRUE, sigma.weighting = NULL, sigma.norm = FALSE,
sigma.issample = FALSE, print.results = FALSE)

```

Arguments

gdp1	A numeric vector containing the GDP per capita (or another economic variable) at time t
time1	A single value of time t (= the initial year)

<code>gdp2</code>	A numeric vector containing the GDP per capita (or another economic variable) at time $t+1$
<code>time2</code>	A single value of time $t+1$
<code>sigma.measure</code>	argument that indicates how the sigma convergence should be measured. The default is <code>output = "sd"</code> , which means that the standard deviation is used. If <code>output = "var"</code> or <code>output = "cv"</code> , the variance or the coefficient of variation is used, respectively.
<code>sigma.log</code>	Logical argument. Per default (<code>sigma.log = TRUE</code>), also in the sigma convergence analysis, the economic variables are transformed by natural logarithm. If the original values should be used, state <code>sigma.log = FALSE</code>
<code>sigma.weighting</code>	If the measure of statistical dispersion in the sigma convergence analysis (coefficient of variation or standard deviation) should be weighted, a weighting vector has to be stated
<code>sigma.norm</code>	Logical argument that indicates if a normalized coefficient of variation should be used instead
<code>sigma.issample</code>	logical argument that indicates if the dataset is a sample or the population (default: <code>is.sample = FALSE</code> , so the denominator of variance is n)
<code>print.results</code>	Logical argument that indicates if the function shows the results or not

Details

From the regional economic perspective (in particular the neoclassical growth theory), regional disparities are expected to decline. This *convergence* can have different meanings: *Sigma convergence* (σ) means a harmonization of regional economic output or income over time, while *beta convergence* (β) means a decline of dispersion because poor regions have a stronger economic growth than rich regions (Capello/Nijkamp 2009). Regardless of the theoretical assumptions of a harmonization in reality, the related analytical framework allows to analyze both types of convergence for cross-sectional data (GDP p.c. or another economic variable, y , for i regions and two points in time, t and $t + T$), or one starting point (t) and the average growth within the following n years ($t + 1, t + 2, \dots, t + n$), respectively. Beta convergence can be calculated either in a linearized OLS regression model or in a nonlinear regression model. When no other variables are integrated in this model, it is called *absolute* beta convergence. Implementing other region-related variables (conditions) into the model leads to *conditional* beta convergence. If there is beta convergence ($\beta < 0$), it is possible to calculate the *speed of convergence*, λ , and the so-called *Half-Life* H , while the latter is the time taken to reduce the disparities by one half (Allington/McCombie 2007, Goecke/Huether 2016). There is *sigma convergence*, when the dispersion of the variable (σ), e.g. calculated as standard deviation or coefficient of variation, reduces from t to $t + T$. This can be measured using ANOVA for two years or trend regression with respect to several years (Furceri 2005, Goecke/Huether 2016).

This function calculates the standard deviation (or variance, coefficient of variation) for the GDP per capita (or another economic variable) for both years and executes an analysis of variance (ANOVA) for these deviation measures (year 1 divided by year 2, F-statistic). If $\sigma_{t1}/\sigma_{t2} > 0$, there is sigma convergence.

Value

Returns a matrix containing the standard deviations, their quotient and the results of the significance test (F-statistic).

Author(s)

Thomas Wieland

References

Allington, N. F. B./McCombie, J. S. L. (2007): "Economic growth and beta-convergence in the East European Transition Economies". In: Arestis, P./Baddely, M./McCombie, J. S. L. (eds.): *Economic Growth. New Directions in Theory and Policy*. Cheltenham: Elgar. p. 200-222.

Capello, R./Nijkamp, P. (2009): "Introduction: regional growth and development theories in the twenty-first century - recent theoretical advances and future challenges". In: Capello, R./Nijkamp, P. (eds.): *Handbook of Regional Growth and Development Theories*. Cheltenham: Elgar. p. 1-16.

Dapena, A. D./Vazquez, E. F./Morollon, F. R. (2016): "The role of spatial scale in regional convergence: the effect of MAUP in the estimation of beta-convergence equations". In: *The Annals of Regional Science*, **56**, 2, p. 473-489.

Furceri, D. (2005): "Beta and sigma-convergence: A mathematical relation of causality". In: *Economics Letters*, **89**, 2, p. 212-215.

Goecke, H./Huether, M. (2016): "Regional Convergence in Europe". In: *Intereconomics*, **51**, 3, p. 165-171.

Young, A. T./Higgins, M. J./Levy, D. (2008): "Sigma Convergence versus Beta Convergence: Evidence from U.S. County-Level Data". In: *Journal of Money, Credit and Banking*, **40**, 5, p. 1083-1093.

See Also

[rca](#), [sigmaconv.t](#), [betaconv.nls](#), [betaconv.speed](#), [cv](#), [sd2](#), [var2](#)

Examples

```
data(G.counties.gdp)
# Loading GDP data for Germany (counties = Landkreise)

sigmaconv (G.counties.gdp$gdppc2010, 2010, G.counties.gdp$gdppc2011, 2011,
sigma.measure = "cv", print.results = TRUE)
# Using the coefficient of variation

sigmaconv (G.counties.gdp$gdppc2010, 2010, G.counties.gdp$gdppc2011, 2011,
sigma.log = TRUE, print.results = TRUE)
# Using the standard deviation with logged GDP per capita
```

sigmaconv.t	<i>Analysis of regional sigma convergence for a time series using trend regression</i>
-------------	--

Description

This function provides the analysis of regional economic sigma convergence (decline of deviation) for a time series using a trend regression

Usage

```
sigmaconv.t(gdp1, time1, gdp2, time2, sigma.measure = "sd", sigma.log = TRUE,
sigma.weighting = NULL, sigma.issample = FALSE,
sigma.plot = FALSE, sigma.plotLSize = 1, sigma.plotLineCol = "black",
sigma.plotRLine = FALSE, sigma.plotRLineCol = "blue",
sigma.Ymin = 0, sigma.plotX = "Time", sigma.plotY = "Variation",
sigma.plotTitle = "Sigma convergence", sigma.bgCol = "gray95", sigma.bgrid = TRUE,
sigma.bgridCol = "white", sigma.bgridSize = 2, sigma.bgridType = "solid",
print.results = FALSE)
```

Arguments

gdp1	A numeric vector containing the GDP per capita (or another economic variable) at time t
time1	A single value of time t (= the initial year)
gdp2	A data frame containing the GDPs per capita (or another economic variable) at time $t+1, t+2, t+3, \dots, t+n$
time2	A single value of time $t+1$
sigma.measure	argument that indicates how the sigma convergence should be measured. The default is <code>output = "sd"</code> , which means that the standard deviation is used. If <code>output = "var"</code> or <code>output = "cv"</code> , the variance or the coefficient of variation is used, respectively.
sigma.log	Logical argument. Per default (<code>sigma.log = TRUE</code>), also in the sigma convergence analysis, the economic variables are transformed by natural logarithm. If the original values should be used, state <code>sigma.log = FALSE</code>
sigma.weighting	If the measure of statistical dispersion in the sigma convergence analysis (coefficient of variation or standard deviation) should be weighted, a weighting vector has to be stated
sigma.issample	Logical argument that indicates if the dataset is a sample or the population (default: <code>is.sample = FALSE</code> , so the denominator of variance is n)
sigma.plot	Logical argument that indicates if a plot of sigma convergence has to be created
sigma.plotLSize	If <code>sigma.plot = TRUE</code> : Line size of the sigma convergence plot

sigma.plotLineCol	If sigma.plot = TRUE: Line color of the sigma convergence plot
sigma.plotRLine	If sigma.plot = TRUE: Logical argument that indicates if a regression line has to be added to the plot
sigma.plotRLineCol	If sigma.plot = TRUE and sigma.plotRLine = TRUE: Color of the regression line
sigma.Ymin	If sigma.plot = TRUE: start value of the Y axis in the plot
sigma.plotX	If sigma.plot = TRUE: Name of the X axis
sigma.plotY	If sigma.plot = TRUE: Name of the Y axis
sigma.plotTitle	If sigma.plot = TRUE: Title of the plot
sigma.bgCol	If sigma.plot = TRUE: Plot background color
sigma.bgrid	If sigma.plot = TRUE: Logical argument that indicates if the plot contains a grid
sigma.bgridCol	If sigma.plot = TRUE and sigma.bgrid = TRUE: Color of the grid
sigma.bgridSize	If sigma.plot = TRUE and sigma.bgrid = TRUE: Size of the grid
sigma.bgridType	If sigma.plot = TRUE and sigma.bgrid = TRUE: Type of the grid
print.results	Logical argument that indicates if the function shows the results or not

Details

From the regional economic perspective (in particular the neoclassical growth theory), regional disparities are expected to decline. This *convergence* can have different meanings: *Sigma convergence* (σ) means a harmonization of regional economic output or income over time, while *beta convergence* (β) means a decline of dispersion because poor regions have a stronger economic growth than rich regions (Capello/Nijkamp 2009). Regardless of the theoretical assumptions of a harmonization in reality, the related analytical framework allows to analyze both types of convergence for cross-sectional data (GDP p.c. or another economic variable, y , for i regions and two points in time, t and $t + T$), or one starting point (t) and the average growth within the following n years ($t + 1, t + 2, \dots, t + n$), respectively. Beta convergence can be calculated either in a linearized OLS regression model or in a nonlinear regression model. When no other variables are integrated in this model, it is called *absolute* beta convergence. Implementing other region-related variables (conditions) into the model leads to *conditional* beta convergence. If there is beta convergence ($\beta < 0$), it is possible to calculate the *speed of convergence*, λ , and the so-called *Half-Life* H , while the latter is the time taken to reduce the disparities by one half (Allington/McCombie 2007, Goecke/Huether 2016). There is *sigma convergence*, when the dispersion of the variable (σ), e.g. calculated as standard deviation or coefficient of variation, reduces from t to $t + T$. This can be measured using ANOVA for two years or trend regression with respect to several years (Furceri 2005, Goecke/Huether 2016).

This function calculates the standard deviation (or variance, coefficient of variation) for all GDPs per capita (or another economic variable) for the given years and executes a trend regression for these deviation measures. If the slope of the trend regression is negative, there is sigma convergence.

Value

Returns a matrix containing the trend regression model and the resulting significance tests (F-statistic, t-statistic).

Author(s)

Thomas Wieland

References

Allington, N. F. B./McCombie, J. S. L. (2007): “Economic growth and beta-convergence in the East European Transition Economies”. In: Arestis, P./Baddely, M./McCombie, J. S. L. (eds.): *Economic Growth. New Directions in Theory and Policy*. Cheltenham: Elgar. p. 200-222.

Capello, R./Nijkamp, P. (2009): “Introduction: regional growth and development theories in the twenty-first century - recent theoretical advances and future challenges”. In: Capello, R./Nijkamp, P. (eds.): *Handbook of Regional Growth and Development Theories*. Cheltenham: Elgar. p. 1-16.

Dapena, A. D./Vazquez, E. F./Morollon, F. R. (2016): “The role of spatial scale in regional convergence: the effect of MAUP in the estimation of beta-convergence equations”. In: *The Annals of Regional Science*, **56**, 2, p. 473-489.

Furceri, D. (2005): “Beta and sigma-convergence: A mathematical relation of causality”. In: *Economics Letters*, **89**, 2, p. 212-215.

Goecke, H./Huether, M. (2016): “Regional Convergence in Europe”. In: *Intereconomics*, **51**, 3, p. 165-171.

Young, A. T./Higgins, M. J./Levy, D. (2008): “Sigma Convergence versus Beta Convergence: Evidence from U.S. County-Level Data”. In: *Journal of Money, Credit and Banking*, **40**, 5, p. 1083-1093.

See Also

[rca](#), [sigmaconv](#), [betaconv.nls](#), [betaconv.speed](#), [cv](#), [sd2](#), [var2](#)

Examples

```
data(G.counties.gdp)
# Loading GDP data for Germany (counties = Landkreise)

# Sigma convergence 2010-2014:
sigmaconv.t (G.counties.gdp$gdppc2010, 2010, G.counties.gdp[65:68], 2014,
sigma.plot = TRUE, print.results = TRUE)
# Using the standard deviation with logged GDP per capita

sigmaconv.t (G.counties.gdp$gdppc2010, 2010, G.counties.gdp[65:68], 2014,
sigma.measure = "cv", sigma.log = FALSE, print.results = TRUE)
# Using the coefficient of variation (GDP per capita not logged)
```

spec

Measures of regional specialization

Description

Calculating three measures of regional specialization (Gini, Krugman, Hoover) for a set of J regions

Usage

```
spec(e_ij, industry.id, region.id, na.rm = TRUE)
```

Arguments

<code>e_ij</code>	a numeric vector with the employment of the industry i in region j
<code>industry.id</code>	a vector containing the IDs of the industries i
<code>region.id</code>	a vector containing the IDs of the regions j
<code>na.rm</code>	logical argument that indicates whether NA values should be excluded before computing results

Details

This function is a convenient wrapper for all functions calculating measures of regional specialization (Gini, Krugman, Hoover)

Value

A matrix with three columns (Gini coefficient, Krugman coefficient, Hoover coefficient) and J rows (one for each regarded region).

Author(s)

Thomas Wieland

References

Farhauer, O./Kroell, A. (2014): "Standorttheorien: Regional- und Stadtoekonomik in Theorie und Praxis". Wiesbaden : Springer.

Schaetzl, L. (2000): "Wirtschaftsgeographie 2: Empirie". Paderborn : Schoeningh.

See Also

[gini.spec](#), [krugman.spec2](#), [hoover](#)

Examples

```
data(G.regions.industries)

spec_j <- spec (e_ij = G.regions.industries$emp_all,
               industry.id = G.regions.industries$ind_code,
               region.id = G.regions.industries$region_code)
```

theil	<i>Theil inequality index</i>
-------	-------------------------------

Description

Calculating the Theil inequality index

Usage

```
theil(x, weighting = NULL, na.rm = TRUE)
```

Arguments

x	a numeric vector
weighting	a numeric weighting vector, e.g. population
na.rm	logical argument that indicates whether NA values should be excluded before computing results

Details

Since there are several Theil measures of inequality, this function uses the formulation from Stoermann (2009).

Value

A single numeric value of the *Theil inequality index* ($0 < TI < 1$).

Author(s)

Thomas Wieland

References

Portnov, B.A./Felsenstein, D. (2010): "On the suitability of income inequality measures for regional analysis: Some evidence from simulation analysis and bootstrapping tests". In: *Socio-Economic Planning Sciences*, **44**, 4, p. 212-219.

Stoermann, W. (2009): "Regionaloekonomik: Theorie und Politik". Muenchen : Oldenbourg.

See Also

[gini](#), [herf](#), [hoover](#)

Examples

```
# Example from Stoermann (2009):  
regincome <- c(10,10,10,20,50)  
theil(regincome)  
# 0.2326302
```

to.dummy

Creating dummy variables

Description

This function creates a dataset of dummy variables based on an input character vector.

Usage

```
to.dummy(x)
```

Arguments

x A character vector

Details

This function transforms a character vector *x* with *c* characteristics to a set of *c* dummy variables whose column names corresponding to these characteristics marked with “_DUMMY”.

Value

A data.frame with dummy variables corresponding to the levels of the input variable.

Note

This function contains code from the authors' package MCI.

Author(s)

Thomas Wieland

References

Greene, W. H. (2012): “Econometric Analysis”. 7th edition. Harlow : Pearson.

Examples

```
charvec <- c("Peter", "Paul", "Peter", "Mary", "Peter", "Paul")  
# Creates a vector with three names (Peter, Paul, Mary)  
to.dummy(charvec)  
# Returns a data frame with 3 dummy variables  
# (Mary_DUMMY, Paul_DUMMY, Peter_DUMMY)
```

var2	<i>Variance (extended)</i>
------	----------------------------

Description

Calculating the variance (var), weighted or non-weighted, for samples or populations

Usage

```
var2(x, is.sample = TRUE, weighting = NULL, wmean = FALSE, na.rm = TRUE)
```

Arguments

x	a numeric vector
is.sample	logical argument that indicates if the dataset is a sample or the population (default: is.sample = TRUE, so the denominator of variance is $n - 1$)
weighting	a numeric vector containing weighting data to compute the weighted standard deviation (instead of the non-weighted sd)
wmean	logical argument that indicates if the weighted mean is used when calculating the weighted standard deviation
na.rm	logical argument that whether NA values should be extracted or not

Details

The function calculates the *variance* (var). Unlike the R base var function, the var2 function allows to choose if the data is treated as sample (denominator of variance is $n - 1$) or not (denominator of variance is n)

From a regional economic perspective, var and sd is closely linked to the concept of *sigma convergence* (σ) which means a harmonization of regional economic output or income over time, while the other type of convergence, *beta convergence* (β), means a decline of dispersion because poor regions have a stronger growth than rich regions (Capello/Nijkamp 2009). The sd allows to summarize regional disparities (e.g. disparities in regional GDP per capita) in one indicator. The coefficient of variation (see the function cv) is more frequently used for this purpose (e.g. Lessmann 2005, Huang/Leung 2009, Siljak 2015). But the sd can also be used for any other types of disparities or dispersion, such as disparities in supply (e.g. density of physicians or grocery stores).

The variance can be weighted by using a second weighting vector. As there is more than one way to weight measures of statistical dispersion, this function uses the formula for the weighted variance (σ_w) from Sheret (1984). The vector x is automatically treated as a sample (such as in the base sd function), so the denominator of variance is $n - 1$, if it is not, set is.sample = FALSE.

Value

Single numeric value. If weighting is specified, the function returns a weighted variance (optionally using a weighted arithmetic mean if wmean = TRUE).

Author(s)

Thomas Wieland

References

- Bahrenberg, G./Giese, E./Mevenkamp, N./Nipper, J. (2010): "Statistische Methoden in der Geographie. Band 1: Univariate und bivariate Statistik". Stuttgart: Borntraeger.
- Capello, R./Nijkamp, P. (2009): "Introduction: regional growth and development theories in the twenty-first century - recent theoretical advances and future challenges". In: Capello, R./Nijkamp, P. (eds.): *Handbook of Regional Growth and Development Theories*. Cheltenham: Elgar. p. 1-16.
- Lessmann, C. (2005): "Regionale Disparitaeten in Deutschland und ausgesuchten OECD-Staaten im Vergleich". *ifo Dresden berichtet*, **3/2005**. https://www.ifo.de/DocDL/ifoedb_2005_3_25-33.pdf.
- Huang, Y./Leung, Y. (2009): "Measuring Regional Inequality: A Comparison of Coefficient of Variation and Hoover Concentration Index". In: *The Open Geography Journal*, **2**, p. 25-34.
- Sheret, M. (1984): "The Coefficient of Variation: Weighting Considerations". In: *Social Indicators Research*, **15**, 3, p. 289-295.
- Siljak, D. (2015): "Real Economic Convergence in Western Europe from 1995 to 2013". In: *International Journal of Business and Economic Development*, **3**, 3, p. 56-67.

See Also

[sd2](#), [cv](#), [gini](#), [herf](#), [hoover](#), [mean2](#), [rca](#)

Examples

```
# Regional disparities / sigma convergence in Germany
data(G.counties.gdp)
# GDP per capita for German counties (Landkreise)
vars <- apply (G.counties.gdp[54:68], MARGIN = 2, FUN = var2)
# Calculating variance for the years 2000-2014
years <- 2000:2014
plot(years, vars, "l", xlab = "year",
      ylab = "Variance of GDP per capita")
# Plot variance over time
```

williamson

Williamson index

Description

Calculating the Williamson index (population-weighted coefficient of variation)

Usage

```
williamson (x, weighting, coefnorm = FALSE, wmean = FALSE, na.rm = TRUE)
```

Arguments

x	a numeric vector
weighting	mandatory: a numeric vector containing weighting data (usually regional population)
coefnorm	logical argument that indicates if the function output is the standardized cv ($0 < v < \infty$) or not ($0 < v < \infty$) (default: coefnorm = FALSE)
wmean	logical argument that indicates if the weighted mean is used when calculating the weighted coefficient of variation
na.rm	logical argument that whether NA values should be extracted or not

Details

The *Williamson index* (Williamson 1965) is a population-weighted coefficient of variation.

The *coefficient of variation*, v , is a dimensionless measure of statistical dispersion ($0 < v < \infty$), based on variance and standard deviation, respectively. The cv (variance, standard deviation) can be weighted by using a second weighting vector. As there is more than one way to weight measures of statistical dispersion, this function uses the formula for the weighted cv (v_w) from Sheret (1984). The cv can be standardized, while this function uses the formula for the standardized cv (v^* , with $0 < v^* < 1$) from Kohn/Oeztuerk (2013). The vector x is automatically treated as a sample (such as in the base sd function), so the denominator of variance is $n - 1$, if it is not, set `is.sample = FALSE`.

Value

Single numeric value. If `coefnorm = FALSE` the function returns the non-standardized cv ($0 < v < \infty$). If `coefnorm = TRUE` the standardized cv ($0 < v^* < 1$) is returned.

Author(s)

Thomas Wieland

References

- Gluschenko, K. (2018): "Measuring regional inequality: to weight or not to weight?" In: *Spatial Economic Analysis*, **13**, 1, p. 36-59.
- Lessmann, C. (2005): "Regionale Disparitaeten in Deutschland und ausgesuchten OECD-Staaten im Vergleich". *ifo Dresden berichtet*, **3/2005**. https://www.ifo.de/DocDL/ifoedb_2005_3_25-33.pdf.
- Huang, Y./Leung, Y. (2009): "Measuring Regional Inequality: A Comparison of Coefficient of Variation and Hoover Concentration Index". In: *The Open Geography Journal*, **2**, p. 25-34.
- Kohn, W./Oeztuerk, R. (2013): "Statistik fuer Oekonomen. Datenanalyse mit R und SPSS". Berlin: Springer.
- Portnov, B.A./Felsenstein, D. (2010): "On the suitability of income inequality measures for regional analysis: Some evidence from simulation analysis and bootstrapping tests". In: *Socio-Economic Planning Sciences*, **44**, 4, p. 212-219.

Sheret, M. (1984): “The Coefficient of Variation: Weighting Considerations”. In: *Social Indicators Research*, **15**, 3, p. 289-295.

Williamson, J. G. (1965): “Regional Inequality and the Process of National Development: A Description of the Patterns”. In: *Economic Development and Cultural Change*, **13**, 4/2, p. 1-84.

See Also

[gini](#), [herf](#), [hoover](#), [cv](#), [disp](#)

Examples

```
data(GoettingenHealth2)
# districts with healthcare providers and population size

williamson((GoettingenHealth2$phys_gen/GoettingenHealth2$pop),
GoettingenHealth2$pop)
```

Index

- atkinson, [4](#), [17](#), [23](#), [68](#)
- Automotive, [5](#)

- betaconv.nls, [6](#), [11](#), [13](#), [100](#), [123](#), [126](#)
- betaconv.ols, [9](#), [9](#), [13](#), [100](#)
- betaconv.speed, [9](#), [11](#), [12](#), [100](#), [123](#), [126](#)

- conc, [14](#)
- converse, [15](#), [64](#), [73](#), [102](#)
- coulter, [5](#), [16](#), [22](#), [23](#), [68](#)
- curvefit, [17](#)
- cv, [5](#), [9](#), [11](#), [13](#), [17](#), [19](#), [22](#), [23](#), [49](#), [56](#), [66](#), [68](#), [91](#), [93](#), [100](#), [123](#), [126](#), [131](#), [133](#)

- dalton, [5](#), [17](#), [21](#), [22](#), [23](#)
- disp, [5](#), [17](#), [22](#), [22](#), [68](#), [133](#)
- dist, [24](#), [27](#), [104](#)
- dist.buf, [23](#), [25](#), [27](#), [64](#), [104](#)
- dist.calc, [25](#), [64](#)
- dist.mat, [24](#), [25](#), [26](#), [64](#), [104](#)
- durpug, [27](#)

- ellison.a, [28](#), [30–32](#), [80](#), [81](#), [88](#)
- ellison.a2, [29](#), [31](#), [32](#), [80](#), [81](#), [88](#)
- ellison.c, [30](#), [32](#), [69–71](#), [80](#), [81](#), [88](#)
- ellison.c2, [31](#), [31](#), [69–71](#), [80](#), [81](#), [88](#)
- EU28.emp, [33](#)

- FK2014_EGC, [34](#)
- Freiburg, [35](#)

- G.counties.gdp, [36](#)
- G.regions.emp, [39](#)
- G.regions.industries, [40](#)
- gifpro, [41](#), [46](#)
- gifpro.tbs, [43](#), [44](#)
- gini, [5](#), [17](#), [21](#), [22](#), [46](#), [52](#), [54](#), [66](#), [68](#), [106](#), [128](#), [131](#), [133](#)
- gini.conc, [15](#), [29–32](#), [49](#), [50](#), [54](#), [56](#), [75–77](#), [79–81](#), [84](#), [88](#), [91](#)
- gini.spec, [28–32](#), [49](#), [52](#), [52](#), [56](#), [75–77](#), [79–81](#), [84](#), [88](#), [91](#), [127](#)
- gini2, [5](#), [17](#), [22](#), [23](#), [55](#)
- Goettingen, [57](#)
- GoettingenHealth1, [59](#)
- GoettingenHealth2, [60](#)
- growth, [61](#)

- hansen, [62](#), [73](#)
- herf, [5](#), [17](#), [21–23](#), [49](#), [56](#), [65](#), [68](#), [91](#), [106](#), [128](#), [131](#), [133](#)
- hoover, [5](#), [15](#), [17](#), [21–23](#), [28](#), [49](#), [56](#), [67](#), [88](#), [91](#), [106](#), [127](#), [128](#), [131](#), [133](#)
- howard.cl, [29–32](#), [68](#), [70](#), [71](#)
- howard.xcl, [29–32](#), [69](#), [69](#)
- howard.xcl2, [29–32](#), [69](#), [70](#), [70](#), [71](#)
- huff, [16](#), [64](#), [71](#), [102](#)

- krugman.conc, [74](#), [76](#), [77](#), [79](#)
- krugman.conc2, [15](#), [75](#), [75](#), [77](#), [79](#)
- krugman.spec, [28](#), [75](#), [76](#), [77](#), [79](#)
- krugman.spec2, [75–77](#), [78](#), [127](#)

- litzenberger, [29–32](#), [79](#), [81](#), [88](#)
- litzenberger2, [29–32](#), [80](#), [80](#)
- lm.beta, [82](#)
- locq, [29–32](#), [75–77](#), [79–81](#), [83](#), [87](#), [88](#)
- locq.growth, [85](#)
- locq2, [29–32](#), [80](#), [81](#), [84](#), [87](#)
- lorenz, [89](#)

- mean2, [92](#), [106](#), [131](#)
- mssd, [93](#)

- portfolio, [43](#), [46](#), [87](#), [94](#), [109](#), [111](#), [113](#), [115](#), [118](#), [120](#)

- rca, [9](#), [11](#), [21](#), [96](#), [106](#), [123](#), [126](#), [131](#)
- REAT (REAT-package), [3](#)
- REAT-package, [3](#)
- reilly, [16](#), [64](#), [73](#), [101](#)

ripley, 103

sd2, 9, 11, 13, 23, 92, 93, 100, 105, 123, 126,
131

shift, 43, 46, 87, 96, 107, 111, 113, 115, 118

shift.growth, 109, 109, 113, 115, 118, 120

shiftd, 43, 46, 87, 96, 109, 111, 111, 120

shiftd, 43, 46, 87, 96, 109, 111, 113, 114,
115, 118, 120

shiftd, 116

shiftp, 118

sigmaconv, 9, 11, 13, 100, 121, 126

sigmaconv.t, 9, 11, 13, 100, 123, 124

spec, 127

theil, 5, 17, 22, 23, 68, 128

to.dummy, 129

var2, 9, 11, 13, 93, 100, 123, 126, 130

williamson, 23, 131