

Package ‘ierd’

June 16, 2026

Title Inverted Exponentiated Rayleigh Distribution Tools

Version 0.1.1

Description Provides the density, distribution function, quantile function, random generation, and visualization tools for the Inverted Exponentiated Rayleigh Distribution.

License MIT + file LICENSE

Encoding UTF-8

RoxygenNote 8.0.0

Imports dplyr, ggplot2, magrittr, tidyr

URL <https://github.com/SudiptaPal0709/ierd>

BugReports <https://github.com/SudiptaPal0709/ierd/issues>

Language en-US

NeedsCompilation no

Author Sudipta Pal [aut, cre]

Maintainer Sudipta Pal <sudiptapal0709@gmail.com>

Repository CRAN

Date/Publication 2026-06-16 20:10:02 UTC

Contents

dierd	2
pierd	2
plot_dierd	3
plot_pierd	4
qierd	5
rierd	6

Index	7
--------------	----------

dierd

Density of the Inverted Exponentiated Rayleigh Distribution

Description

This function computes the probability density function (PDF) of the Inverted Exponentiated Rayleigh distribution.

Usage

```
dierd(x, shape, scale)
```

Arguments

x	A numeric vector of quantiles.
shape	A strictly positive numeric value for the shape parameter (α).
scale	A strictly positive numeric value for the scale parameter (β).

Details

The probability density function is mathematically defined as:

$$f(x) = 2\alpha\beta x^{-3} \exp(-\beta/x^2)(1 - \exp(-\beta/x^2))^{\alpha-1}$$

for $x > 0$, where α is the shape parameter and β is the scale parameter.

Value

A numeric vector of density values evaluated at x.

Examples

```
# Compute the density at various values of x
dierd(x = c(0.5, 1, 1.5, 2), shape = 2, scale = 1)
```

pierd

Cumulative Distribution Function of the Inverted Exponentiated Rayleigh Distribution

Description

This function computes the Cumulative Distribution Function (CDF) of the Inverted Exponentiated Rayleigh distribution.

Usage

```
pierd(t, shape, scale)
```

Arguments

t	A numeric vector of quantiles.
shape	A strictly positive numeric value for the shape parameter (α).
scale	A strictly positive numeric value for the scale parameter (β).

Details

The cumulative distribution function is mathematically defined as:

$$F(t) = 1 - (1 - \exp(-\beta/t^2))^\alpha$$

for $t > 0$, where α is the shape parameter and β is the scale parameter.

Value

A numeric vector of cumulative probabilities evaluated at t.

Examples

```
# Compute the cumulative probabilities at various values of t
pierd(t = c(0.5, 1, 1.5, 2), shape = 2, scale = 1)
```

plot_dierd

Plot Multiple Inverted Exponentiated Rayleigh Densities

Description

This function creates a ggplot2 visualization comparing the dierd PDF across combinations of shape and scale parameters.

Usage

```
plot_dierd(shape, scale, lower = 0.01, upper = 5, paired = FALSE)
```

Arguments

shape	A numeric vector of strictly positive shape parameters.
scale	A numeric vector of strictly positive scale parameters.
lower	A numeric value for the lower bound of the x-axis (default is 0.01).
upper	A numeric value for the upper bound of the x-axis (default is 5).
paired	Logical. If FALSE (default), creates a full grid of all possible shape and scale combinations. If TRUE, pairs the shape and scale vectors element-by-element (vectors must be the same length).

Value

A ggplot object showing the density curves.

Examples

```
# Full grid: 2 shapes * 2 scales = 4 curves
plot_pierd(shape = c(1, 2), scale = c(1, 2))

# Paired: 2 specific combinations = 2 curves
plot_pierd(shape = c(1, 2), scale = c(1, 2), paired = TRUE)
```

plot_pierd

Plot Multiple Inverted Exponentiated Rayleigh CDFs

Description

This function creates a ggplot2 visualization of the pierd CDF across combinations of shape and scale parameters.

Usage

```
plot_pierd(shape, scale, lower = 0.01, upper = 7.5, paired = FALSE)
```

Arguments

shape	A numeric vector of strictly positive shape parameters.
scale	A numeric vector of strictly positive scale parameters.
lower	A numeric value for the lower bound of the x-axis (default is 0.01).
upper	A numeric value for the upper bound of the x-axis (default is 7.5).
paired	Logical. If FALSE (default), creates a full grid of all possible shape and scale combinations. If TRUE, pairs the shape and scale vectors element-by-element (vectors must be the same length).

Value

A ggplot object showing the cumulative distribution curves.

Examples

```
# Full grid: 2 shapes * 2 scales = 4 curves
plot_pierd(shape = c(1, 2), scale = c(1, 2))

# Paired: 2 specific combinations = 2 curves
plot_pierd(shape = c(1, 2), scale = c(1, 2), paired = TRUE)
```

qierd	<i>Quantile Function of the Inverted Exponentiated Rayleigh Distribution</i>
-------	--

Description

This function computes the quantile function (inverse CDF) of the Inverted Exponentiated Rayleigh distribution.

Usage

```
qierd(p, shape, scale)
```

Arguments

p	A numeric vector of probabilities.
shape	A strictly positive numeric value for the shape parameter (α).
scale	A strictly positive numeric value for the scale parameter (β).

Details

The quantile function is mathematically defined as:

$$Q(p) = \sqrt{\frac{-\beta}{\log(1 - (1 - p)^{1/\alpha})}}$$

for $0 \leq p \leq 1$, where α is the shape parameter and β is the scale parameter.

Value

A numeric vector of quantiles evaluated at p.

Examples

```
# Compute the quantiles at various probability values (e.g., quartiles)
qierd(p = c(0.25, 0.5, 0.75), shape = 2, scale = 1)
```

rierd	<i>Random Numbers from the Inverted Exponentiated Rayleigh Distribution</i>
-------	---

Description

This function generates random numbers from an Inverted Exponentiated Rayleigh distribution using inverse transform sampling.

Usage

```
rierd(n, shape, scale)
```

Arguments

n	An integer specifying the number of random values to return.
shape	A strictly positive numeric value for the shape parameter.
scale	A strictly positive numeric value for the scale parameter.

Value

A numeric vector of length n containing the generated random numbers.

Examples

```
# Generate 10 random numbers with shape = 2 and scale = 1  
rierd(n = 10, shape = 2, scale = 1)
```

Index

dierd, 2

pierd, 2

plot_dierd, 3

plot_pierd, 4

qierd, 5

rierd, 6