

Package ‘modeLLtest’

May 9, 2026

Type Package

Title Compare Models with Cross-Validated Log-Likelihood

Version 1.0.5

Date 2025-10-11

Description An implementation of the cross-validated difference in means (CVDm) test by Desmarais and Harden (2014) <[doi:10.1007/s11135-013-9884-7](https://doi.org/10.1007/s11135-013-9884-7)> (see also Harden and Desmarais, 2011 <[doi:10.1177/1532440011408929](https://doi.org/10.1177/1532440011408929)>) and the cross-validated median fit (CVMF) test by Desmarais and Harden (2012) <[doi:10.1093/pan/mpr042](https://doi.org/10.1093/pan/mpr042)>. These tests use leave-one-out cross-validated log-likelihoods to assist in selecting among model estimations. You can also utilize data from Golder (2010) <[doi:10.1177/0010414009341714](https://doi.org/10.1177/0010414009341714)> and Joshi & Mason (2008) <[doi:10.1177/0022343308096155](https://doi.org/10.1177/0022343308096155)> that are included to facilitate examples from real-world analysis.

URL <https://github.com/ShanaScogin/modeLLtest>

License GPL-3

NeedsCompilation yes

BugReports <https://github.com/ShanaScogin/modeLLtest/issues>

Imports stats, quantreg, survival, coxrobust, MASS, Rcpp

Depends R (>= 3.2.3)

Encoding UTF-8

LazyData TRUE

LazyLoad TRUE

Suggests knitr, rmarkdown, testthat

VignetteBuilder knitr

SystemRequirements GNU make

RoxygenNote 7.1.2

LinkingTo Rcpp, RcppArmadillo

Author Shana Scogin [aut, cre] (ORCID:
<https://orcid.org/0000-0002-7801-853X>),
 Sarah Petersen [aut],
 Jeff Harden [aut],
 Bruce A. Desmarais [aut]

Maintainer Shana Scogin <shanarscogin@gmail.com>

Repository CRAN

Date/Publication 2025-10-12 02:40:02 UTC

Contents

cvdm	2
cvdm_object	4
cvll	5
cvlldiff	6
cvlldiff_object	8
cvll_object	8
cvmf	9
cvmf_object	11
govtform	12
modeLLtest	14
nepaldem	14

Index **18**

cvdm	<i>Cross-Validated Difference in Means (CVDM) Test</i>
------	--

Description

Applies cross-validated log-likelihood difference in means test to compare two methods of estimating a formula. The output identifies the more appropriate model.

In choosing between OLS and MR, please cite:

- Harden, J. J., & Desmarais, B. A. (2011). Linear Models with Outliers: Choosing between Conditional-Mean and Conditional-Median Methods. *State Politics & Policy Quarterly*, 11(4), 371-389. doi: [10.1177/1532440011408929](https://doi.org/10.1177/1532440011408929)

For other applications of the CVDM test, please cite:

- Desmarais, B. A., & Harden, J. J. (2014). An Unbiased Model Comparison Test Using Cross-Validation. *Quality & Quantity*, 48(4), 2155-2173. doi: [10.1007/s1113501398847](https://doi.org/10.1007/s1113501398847)

Note that Eqn. 6 of Desmarais & Harden, 2014 has a typo in the denominator. The code here correctly implements Johnson (1978) (doi: [10.1080/01621459.1978.10480051](https://doi.org/10.1080/01621459.1978.10480051)), Eqn. 2.5; however, the fraction is flipped in the paper's equation.

Usage

```
cvdm(
  formula,
  data,
  method1 = c("OLS", "MR", "RLM", "RLM-MM"),
  method2 = c("OLS", "MR", "RLM", "RLM-MM"),
  subset,
  na.action,
  ...
)
```

Arguments

formula	A formula object, with the dependent variable on the left of a ~ operator, and the independent variables on the right.
data	A data frame, list or environment (or object coercible by <code>as.data.frame</code> to a data frame) containing the variables in the model.
method1	A method to estimate the model. Currently takes Ordinary Least Squares ("OLS"), Median Regression ("MR"), Robust Linear Regression ("RLM") using M-estimation, and Robust Linear Regression using MM-estimation ("RLM-MM"). The algorithm method used to compute the fit for the median regression is the modified version of the Barrodale and Roberts algorithm for l1-regression, which is the rq default by R package <code>quantreg</code> . See <code>quantreg</code> rq function documentation for more details. Fitting for the robust regressions is done by iterated re-weighted least squares (IWLS) and is taken from the MASS package rlm function. The MM-estimation is the M-estimation with Tukey's biweight initialized by a specific S-estimate. The M-estimation, which can be achieved in this package with the option "RLM", is the default for the MASS rlm function. See MASS package rlm documentation for details.
method2	A method to estimate the model. Options are same as for method1.
subset	Expression indicating which subset of the rows of data should be used in the fit. All observations are included by default.
na.action	A missing-data filter function, applied to the model.frame, after any subset argument has been used.
...	Optional arguments, currently unsupported.

Details

This function implements the cross-validated difference in means (CVDM) test between two methods of estimating a formula. The function takes a formula and two methods and computes a vector of cross-validated log-likelihoods (CVLLs) for each method using the leave-one-out method. These output test score is the cross-validated Johnson's t-test. A positive test statistic supports the first method and a negative test statistic supports the second. Singular matrices during the leave-one-out cross-validation process are skipped.

Value

An object of class `cvdm` computed by the cross-validated log likelihood difference in means test (CVDM). The object is the Cross-Validated Johnson's t-test. A positive test statistic supports the first method and a negative test statistic supports the second. See [cvdm_object](#) for more details.

References

- Harden, J. J., & Desmarais, B. A. (2011). Linear Models with Outliers: Choosing between Conditional-Mean and Conditional-Median Methods. *State Politics & Policy Quarterly*, 11(4), 371-389. doi: [10.1177/1532440011408929](https://doi.org/10.1177/1532440011408929)
- Desmarais, B. A., & Harden, J. J. (2014). An Unbiased Model Comparison Test Using Cross-Validation. *Quality & Quantity*, 48(4), 2155-2173. doi: [10.1007/s1113501398847](https://doi.org/10.1007/s1113501398847)

Examples

```
set.seed(123456)
b0 <- .2 # True value for the intercept
b1 <- .5 # True value for the slope
n <- 500 # Sample size
X <- runif(n, -1, 1)

Y <- b0 + b1 * X + rnorm(n, 0, 1) # N(0, 1 error)

obj_cvdm <- cvdm(Y ~ X, data.frame(cbind(Y, X)), method1 = "OLS", method2 = "MR")
```

 cvdm_object

Cross-Validated Difference in Means (CVDM) Object

Description

This class of objects is returned by the `cvdm` function to compare two methods of estimating a formula.

Value

The following components must be included in a legitimate `cvdm` object.

<code>best</code>	name of the estimation method favored by the <code>cvdm</code> test.
<code>test_stat</code>	object returned by the bias-corrected Johnson's t-test. A positive test statistic supports method 1 and a negative test statistic supports method 2.
<code>p_value</code>	p-value for the test statistic.
<code>n</code>	number of observations.
<code>df</code>	degrees of freedom.

The object also contain the following: `call`, `x`, and `y`. See [lm](#) documentation for more.

See Also[cvdm](#)

cvll

*Cross-Validated Log Likelihood (CVLL)***Description**

Extracts the leave-one-out cross-validated log-likelihoods from a method of estimating a formula.

Usage

```
cvll(
  formula,
  data,
  method = c("OLS", "MR", "RLM", "RLM-MM"),
  subset,
  na.action,
  ...
)
```

Arguments

formula	A formula object, with the dependent variable on the left of a ~ operator, and the independent variables on the right.
data	A data frame, list or environment (or object coercible by <code>as.data.frame</code> to a data frame) containing the variables in the model.
method	A method to estimate the model. Currently takes Ordinary Least Squares ("OLS"), Median Regression ("MR"), Robust Linear Regression ("RLM") using M-estimation, and Robust Linear Regression using MM-estimation ("RLM-MM"). The algorithm method used to compute the fit for the median regression is the modified version of the Barrodale and Roberts algorithm for l1-regression, which is the rq default by R package <code>quantreg</code> . See <code>quantreg</code> rq function documentation for more details. Fitting for the robust regressions is done by iterated re-weighted least squares (IWLS) and is taken from the MASS package <code>rlm</code> function. The MM-estimation is the M-estimation with Tukey's biweight initialized by a specific S-estimate. The M-estimation, which can be achieved in this package with the option "RLM", is the default for the MASS <code>rlm</code> function. See MASS package <code>rlm</code> documentation for details.
subset	Expression indicating which subset of the rows of data should be used in the fit. All observations are included by default.
na.action	A missing-data filter function, applied to the <code>model.frame</code> , after any subset argument has been used.
...	Optional arguments, currently unsupported.

Details

This function extracts a vector of leave-one-out cross-validated log likelihoods (CVLLs) from a method of estimating a formula. Singular matrices during the leave-one-out cross-validation process are skipped.

Value

An object of class `cvll` computed by the cross-validated log likelihood (CVLL). See `cvdm_object` for more details.

References

- Harden, J. J., & Desmarais, B. A. (2011). Linear Models with Outliers: Choosing between Conditional-Mean and Conditional-Median Methods. *State Politics & Policy Quarterly*, 11(4), 371-389. doi: [10.1177/1532440011408929](https://doi.org/10.1177/1532440011408929)
- Desmarais, B. A., & Harden, J. J. (2014). An Unbiased Model Comparison Test Using Cross-Validation. *Quality & Quantity*, 48(4), 2155-2173. doi: [10.1007/s1113501398847](https://doi.org/10.1007/s1113501398847)

Note that Eqn. 6 of Desmarais & Harden, 2014 has a typo in the denominator. The code here correctly implements Johnson (1978) (doi: [10.1080/01621459.1978.10480051](https://doi.org/10.1080/01621459.1978.10480051)), Eqn. 2.5; however, the fraction is flipped in the paper's equation.

Examples

```
set.seed(123456)
b0 <- .2 # True value for the intercept
b1 <- .5 # True value for the slope
n <- 500 # Sample size
X <- runif(n, -1, 1)

Y <- b0 + b1 * X + rnorm(n, 0, 1) # N(0, 1 error)

obj_cvll <- cvll(Y ~ X, data.frame(cbind(Y, X)), method = "OLS")
```

cvlldiff

Cross-Validated Difference in Means (CVDM) Test with Vector Inputs

Description

Applies cross-validated log-likelihood to test between two methods of estimating a formula. The output identifies the vector from the more appropriate model.

Please cite:

Desmarais, B. A., & Harden, J. J. (2014). An Unbiased Model Comparison Test Using Cross-Validation. *Quality & Quantity*, 48(4), 2155-2173. doi: [10.1007/s1113501398847](https://doi.org/10.1007/s1113501398847)

Usage

```
cvlldiff(vector1, vector2, df)
```

Arguments

vector1	A numeric vector of cross-validated log-likelihoods.
vector2	A numeric vector of cross-validated log-likelihoods.
df	A value of the degrees of freedom in the models.

Details

This function implements the cross-validated difference in means (CVDM) test between two vectors of cross-validated log-likelihoods. A positive test statistic supports the method that produced the first vector and a negative test statistic supports the second.

Value

An object of class `cvlldiff` computed by the cross-validated log likelihood difference in means test (CVDM). The test statistic object is the Cross-Validated Johnson's t-test. A positive test statistic supports the first method and a negative test statistic supports the second. See [cvdm_object](#) for more details.

References

Desmarais, B. A., & Harden, J. J. (2014). An Unbiased Model Comparison Test Using Cross-Validation. *Quality & Quantity*, 48(4), 2155-2173. doi: [10.1007/s1113501398847](https://doi.org/10.1007/s1113501398847) Note that Eqn. 6 of Desmarais & Harden, 2014 has a typo in the denominator. The code here correctly implements Johnson (1978) (doi: [10.1080/01621459.1978.10480051](https://doi.org/10.1080/01621459.1978.10480051)), Eqn. 2.5; however, the fraction is flipped in the paper's equation.

Examples

```
set.seed(123456)
b0 <- .2 # True value for the intercept
b1 <- .5 # True value for the slope
n <- 500 # Sample size
X <- runif(n, -1, 1)

Y <- b0 + b1 * X + rnorm(n, 0, 1) # N(0, 1 error)
cvll_ols <- cvll(Y ~ X, data.frame(cbind(Y, X)), method = "OLS")
cvll_mr <- cvll(Y ~ X, data.frame(cbind(Y, X)), method = "MR")
obj_compare <- cvlldiff(cvll_ols$cvll, cvll_mr$cvll, cvll_ols$df)
```

cvlldiff_object	<i>Cross-Validated Difference in Means (CVDM) Object from General cvlldiff Function</i>
-----------------	---

Description

This class of objects is returned by the [cvlldiff](#) function to compare vectors of cross-validated log-likelihood values.

Value

The following components must be included in a legitimate cvlldiff object.

best	name of the estimation method favored by the cvdm test.
test_stat	object returned by the bias-corrected Johnson's t-test. A positive test statistic supports the method that generated the first vector of cross-validated log-likelihood values and a negative test statistic supports the method that generated the second vector.
p_value	p-value for the test statistic.

See Also

[cvlldiff](#)

cvll_object	<i>Cross-Validated Log-Likelihood (CVLL) Object</i>
-------------	---

Description

This class of objects is returned by the [cvll](#) function.

Value

The following components must be included in a legitimate cvll object.

cvll	vector of cross-validated log-likelihood values using the leave-one-out method.
n	number of observations.
df	degrees of freedom.
method	method of estimation.

The object also contain the following: call, x, and y. See [lm](#) documentation for more.

See Also

[cvll](#)

Description

Applies cross-validated log-likelihood to test between partial likelihood maximization (PLM) and the iteratively reweighted robust (IRR) method of estimation for a given application of the Cox model. For more, see: Desmarais, B. A., & Harden, J. J. (2012). Comparing partial likelihood and robust estimation methods for the Cox regression model. *Political Analysis*, 20(1), 113-135. doi: [10.1093/pan/mpr042](https://doi.org/10.1093/pan/mpr042)

Usage

```
cvmf(
  formula,
  data,
  method = c("exact", "approximate", "efron", "breslow"),
  trunc = 0.95,
  subset,
  na.action,
  f.weight = c("linear", "quadratic", "exponential"),
  weights,
  singular.ok = TRUE
)
```

Arguments

formula	A formula object, with the response on the left of a ~ operator, and the terms on the right. The response must be a survival object as returned by the Surv function from the survival package.
data	A data frame, list or environment (or object coercible by <code>as.data.frame</code> to a data frame) containing the variables in the model or in the subset and the weights argument.
method	A character string specifying the method for tie handling in <code>coxph()</code> . If there are no tied death times all the methods are equivalent. Following the coxph function in the survival package, the Efron approximation is used as the default. The survival package justifies this due to the Efron method being more accurate when dealing with tied death times, and is as efficient computationally than the common Breslow method. The "exact partial likelihood" is equivalent to a 'conditional logistic model, and is appropriate when the times are a small set of discrete values. This argument does not exist in the coxr function in the <code>coxrobust</code> package. For coxr , method is based on a smooth modification of the partial likelihood. See documentation from survival package for more on coxph method and <code>coxrobust</code> package for coxr method.
trunc	A value that determines the trimming level for the robust estimator. The default is 0.95. Roughly, quantile of the sample $T_i \exp(\beta' Z_i)$. It is an argument in the coxr function in the <code>coxrobust</code> package.

subset	Expression indicating which subset of the rows of data should be used in the fit. All observations are included by default.
na.action	A missing-data filter function, applied to the model.frame, after any subset argument has been used.
f.weight	A type of weighting function for <code>coxr</code> in the <code>coxrobust</code> package. The default is quadratic. See <code>coxr</code> documentation for more.
weights	A vector of case weights for <code>coxph</code> in the <code>survival</code> package. See <code>coxph</code> documentation for more.
singular.ok	Logical value indicating how to handle collinearity in the model matrix. If TRUE, the program will automatically skip over columns of the X matrix that are linear combinations of earlier columns. In this case the coefficients for such columns will be NA, and the variance matrix will contain zeros. For ancillary calculations, such as the linear predictor, the missing coefficients are treated as zeros.

Details

This function implements the cross-validated median fit (CVMF) test. The function `cvmf()` tests between the partial likelihood maximization (PLM) and the iteratively reweighted robust (IRR) method of estimation for a given application of the Cox model. The Cox model is a partial parametric model that does not make assumptions about the baseline hazard. It can be estimated via PLM, the standard estimator, or IRR, a robust estimator that identifies and downweights outliers. The choice between the two methods involves a trade-off between bias and efficiency. PLM is more efficient, but biased under specification problems. IRR reduces bias, but results in high variance due to the loss of efficiency. The `cvmf()` function returns an object to identify the preferred estimation method.

See also [coxph](#), [coxr](#), [Surv](#)

Value

An object of class `cvmf` computed by the cross-validated median fit test (CVMF) to test between the PLM and IRR methods of estimating the Cox model. See [cvmf_object](#) for more details.

References

Desmarais, B. A., & Harden, J. J. (2012). Comparing partial likelihood and robust estimation methods for the Cox regression model. *Political Analysis*, 20(1), 113-135. doi: [10.1093/pan/mpr042](https://doi.org/10.1093/pan/mpr042)

Examples

```
set.seed(12345)
x1 <- rnorm(100)
x2 <- rnorm(100)

x2e <- x2 + rnorm(100, 0, 0.5)

y <- rexp(100, exp(x1 + x2))
y <- survival::Surv(y)
```

```

dat <- data.frame(y, x1, x2e)
form <- y ~ x1 + x2e

results <- cvmf(formula = form, data = dat)

```

cvmf_object

*Cross-Validated Median Fit (CVMF) Object***Description**

This class of objects is returned by the `cvmf` function to test between the partial likelihood maximization (PLM) and the iteratively reweighted robust (IRR) method of estimation for a given application of the Cox model.

Value

The following components must be included in a legitimate `cvmf` object.

<code>best</code>	name of the model of estimation favored by the <code>cvmf</code> test.
<code>p</code>	p-value of the binomial test used to test between estimation models.
<code>cvmf</code>	full output of the binomial test used to test between estimation methods. See documentation for <code>binom.test</code> for more information.
<code>coef_names</code>	names of the coefficients.
<code>irr</code>	full output for the iteratively reweighted robust (IRR) method of estimating the Cox model. See documentation for <code>coxr</code> in the package <code>coxrobust</code> for more information.
<code>plm</code>	full output for the partial likelihood maximization (PLM) method of estimating the Cox model. See documentation for <code>coxph</code> in the package <code>survival</code> for more information.
<code>irr_coefs</code>	estimates obtained from IRR method of estimating the Cox model. See documentation for <code>coxr</code> in the package <code>coxrobust</code> for more information.
<code>plm_coefs</code>	estimates obtained from PLM method of estimating the Cox model. See documentation for <code>coxph</code> in the package <code>survival</code> for more information.
<code>cvpl_irr</code>	observation-wise contributions to the log-partial likelihood for IRR method of estimating the Cox model. See Desmarais and Hardin (Political Analysis 20:113-135, 2012) for more about the test and Verweij and Houwelingen (Statistics in Medicine 12(24): 2305–14, 1993) for more about the measure
<code>cvpl_plm</code>	observation-wise contributions to the log-partial likelihood for PLM method of estimating the Cox model. See Desmarais and Hardin (Political Analysis 20:113-135, 2012) for more about the test and Verweij and Houwelingen (Statistics in Medicine 12(24): 2305–14, 1993) for more about the measure

The object also contain the following: `call`, `x`, and `y`.

See Also[cvmf](#)

govtform

*Data from Golder (2010) on government formation in Western Europe***Description**

Data from a study on Western European government formation duration. Data is at the country-level (N = 409). Variable names are taken directly from original dataset. The data is publicly available and has been included here with the endorsement of the author. Please see the original codebook for a more detailed description of the variables.

Usage

```
data(govtform)
```

Format

A data frame with 410 rows and 18 variables. The following are taken from the codebook at doi: [10.7910/DVN/BUWZBA](https://doi.org/10.7910/DVN/BUWZBA).

countryname names of countries used in analysis

country unique number identifying each country

cabinet unique number identifying each country. Begins with country code, followed by cabinets 1 - n

bargainingdays the number of days between either an election or the resignation of the previous government and the day on which the new government is officially inaugurated

datein date on which a government took office. Format is YYMMDD

dateout date on which a government left office. Format is YYMMDD

postelection dichotomous variable that equals 1 if a government is the first to form after an election (more uncertainty) and 0 if it forms in an interelection period (less uncertainty)

nonpartisan dichotomous variable that equals 1 if the government is nonpartisan and 0 otherwise

legislative_parties a fraction representing the number of parties that have won legislative seats. See codebook for more detail

inconclusive the number of inconclusive bargaining rounds prior to a new government successfully forming

cabinetname cabinet name identified by surname of prime minister (followed by a number if the PM presided over more than one cabinet)

singleparty_majority dichotomous variable that equals 1 if a single party controls a majority of the legislative seats, 0 otherwise

polarization measures the level of ideological polarization in the party system. See codebook for more detail

continuation dichotomous variable that equals 1 if the outgoing government or formateur gets the first opportunity to form a new government, 0 otherwise. See codebook for more detail

positive_parl dichotomous variable that equals 1 if a new government requires the explicit support of a legislative majority in order to take office, 0 otherwise. See codebook for more detail

post_legislative_parties interaction term made by multiplying the postelection variable with the legislative_parties variable

post_polariz interaction term made by multiplying the postelection variable with the polarization variable

post_positive interaction term made by multiplying the postelection variable with the positive_parl variable

Source

doi: [10.7910/DVN/BUWZBA](https://doi.org/10.7910/DVN/BUWZBA)

References

Golder, S. N. (2010). Bargaining delays in the government formation process. *Comparative Political Studies*, 43(1), 3-32. doi: [10.1177/0010414009341714](https://doi.org/10.1177/0010414009341714)

Examples

```
data(govtform)

library(survival)
library(coxrobust)
library(modelLtest)

# Survival models with data from Golder (2010)
golde_r_surv <- Surv(govtform$bargainingdays)
golde_r_x <- cbind(govtform$postelection, govtform$legislative_parties,
  govtform$polarization, govtform$positive_parl, govtform$post_legislative_parties,
  govtform$post_polariz, govtform$post_positive, govtform$continuation,
  govtform$singleparty_majority)
colnames(golde_r_x) <- c("govtform$postelection", "govtform$legislative_parties",
  "govtform$polarization", "govtform$positive_parl", "govtform$post_legislative_parties",
  "govtform$post_polariz", "govtform$post_positive", "govtform$continuation",
  "govtform$singleparty_majority")
golde_r_cox <- coxph(golde_r_surv ~ golde_r_x, method = "efron",
  data = govtform)
golde_r_robust <- coxr(golde_r_surv ~ golde_r_x, data = govtform)

# Comparing PLM to IRR methods of estimating the survival model
obj_cvmf_golde_r <- cvmf(golde_r_surv ~ golde_r_x, method = "efron",
  data = govtform)

obj_cvmf_golde_r
```

 modeLLtest

modeLLtest Overview

Description

modeLLtest has three main functions to implement cross validated log likelihood tests. To use this package, decide which specification(s) of a model and distributions you wish compare. The function `cvdm()` compares the fits of one model specification between a median regression and ordinary least squares. The function `cvmf()` compares between the fits of one model specification between two estimations of a Cox model. The function `cvll()` extracts the leave-one-out cross-validated log-likelihoods from a method of estimating a formula.

 nepaldem

Data from Joshi and Mason (2008) on voter turnout in Nepal

Description

Data from a study on the relationship between land tenure and voter turnout in the three rounds of parliamentary elections in Nepal from the restoration of democracy in 1990 to 1999. Data is at the district-level (N = 75). Variable names are taken directly from original dataset. The data is publicly available and has been included here with the endorsement of the authors.

Usage

```
data(nepaldem)
```

Format

A data frame with 76 rows and 73 variables:

sn a column of identifiers. This column is not a variable

district names of the district in Nepal used in analysis

householdsize average size of household in district

total_holding total land holding

noown_single_tenure number of households that own and cultivate land under single tenure

norent_single_ten number of households that rent for service and cultivate land under single tenure

noother_single_ten number of households that cultivate under single tenure and have another set up other than those above

nomore1_ten_hold number of households with more than one tenure

noholding_below1_pa number of households that hold less than 1.0 hectares of land

noholding_2to3_pa number of households that hold 2 to 3 hectares of land
noholding_4to5_pa number of households that hold 4 to 5 hectares of land
noholding_6to9_pa number of households that hold 6 to 9 hectares of land
noholding_10_pa number of households with more than 10 parcels of land
total_ha total hectares of land
total_parcel total parcels of land
no_hold_fixmoney2 subsection of number of households with fixed cash rent
no_hold_fixproduct2 subsection of households with fixed product rent
no_hold_share2 subsection of households participating in sharecropping
no_hold_services2 subsection of households participating in sharecropping
no_hold_mortgage2 subsection of households with a mortgage
no_hold_fixmoney1 subsection of households with fixed cash rent
no_hold_fixproduct1 subsection of households with fixed product rent
no_hold_share1 subsection of households participating in sharecropping
no_hold_services1 subsection of households with rent for service
no_hold_mortgage1 subsection of households with a mortgage
totalhouseholds total number of households
landless number of landless households
totalvoters1991 total number of voters in 1991
totalcastedvote1991 total number of votes cast in 1991
totalvalidvote1991 total number of valid votes in 1991
constituency1991 constituency in 1991
totalcontestants1991 total number of candidates contesting elections in 1991
totalvoters1994 total number of voters in 1994
totalcastedvote1994 total number of votes cast in 1994
totalvalidvote1994 total number of valid votes in 1994
constituency1994 constituency in 1994
totalcontestants1994 total number of candidates contesting elections in 1994
togalvoters1999 total number of voters in 1999
totalcastedvote1999 total number of votes cast in 1999
totalvalidvote1999 total number of valid votes in 1999
constituency1999 constituency in 1999
totalcontestants1999 total number of candidates contesting elections in 1999
pop_2001 population in 2001
hdi_1996 HDI 1996 (index 0 to 1)
per_without_instcredit percent without access to institutional credit
access_insttional_credit access to institutional credit

total_hh_sharecrop total number of households participating in sharecropping

total_hh_fixmoney total number of households with fixed cash rent

total_hh_fixproduct total number of households with fixed product rent

total_hh_service total number of households with rent for service

total_hh_mortgage total number of households with a mortgage

total_killed total number of people killed. This serves as a measure of political violence during the insurgency

percent_regvote1991 election turnout for 1991 as measured by the percentage of registered voters who voted in the national parliamentary election

percent_regvote1994 election turnout for 1994 as measured by the percentage of registered voters who voted in the national parliamentary election

percent_regvote1999 election turnout for 1999 as measured by the percentage of registered voters who voted in the national parliamentary election

per_total_hold_sharecrop percent of sharecropping households

per_total_hold_fixmoney percent of households that have a fixed cash rent

per_total_hold_fixproduct percent of households that have a fixed product rent

per_total_hold_service percent of households that have rent for service

per_total_hold_mortgage percent of households with a mortgage

per_noholding_below1_pa

landless_1000 landless households (in 1,000s)

totoalkilled_1000 total number of people killed (in 1,000s). This serves as a measure of political violence during the insurgency

cast_eth_fract caste and ethnic fractionalization

linguistic_fract linguistic fractionalization

landless_gap landless households (in 1,000s) gap

below1pa_gap percent smallholder households gap

sharecrop_gap percent sharecropping households gap

service_gap percent rent for service households gap

fixmoney_gap percent fixed cash rent households gap

fixprod_gap percent fixed product rent households gap

hdi_gap HDI 1996 (index 0 to 1) gap

ln_pop2001 population in 2001 (logged)

hdi_gap1 HDI 1996 (index 0 to 1) gap (positive values)

Source

[Journal of Peace Research Replication Datasets](#)

References

Joshi, M., & Mason, T. D. (2008). Between democracy and revolution: peasant support for insurgency versus democracy in Nepal. *Journal of Peace Research*, 45(6), 765-782. doi: [10.1177/0022343308096155](https://doi.org/10.1177/0022343308096155)

Examples

```
data(nepaldem)

library(MASS)
library(modelLtest)

# Models from Joshi and Mason (2008)
model_1991 <- rlm(percent_regvote1991 ~ landless_gap +
  below1pa_gap + sharecrop_gap + service_gap + fixmoney_gap +
  fixprod_gap + per_without_instcredit + hdi_gap1 + ln_pop2001 +
  totalcontestants1991 + cast_eth_fract, data = nepaldem)

model_1994 <- rlm(percent_regvote1994 ~ landless_gap +
  below1pa_gap + sharecrop_gap + service_gap + fixmoney_gap +
  fixprod_gap + per_without_instcredit + hdi_gap1 + ln_pop2001 +
  totalcontestants1994 + cast_eth_fract, data = nepaldem)

model_1999a <- rlm(percent_regvote1999 ~ landless_gap +
  below1pa_gap + sharecrop_gap + service_gap + fixmoney_gap +
  fixprod_gap + per_without_instcredit + hdi_gap1 + ln_pop2001 +
  totalcontestants1999 + cast_eth_fract, data = nepaldem)

model_1999b <- rlm(percent_regvote1999 ~ landless_gap +
  below1pa_gap + sharecrop_gap + service_gap + fixmoney_gap +
  fixprod_gap + per_without_instcredit + totoalkilled_1000 +
  hdi_gap1 + ln_pop2001 + totalcontestants1999 + cast_eth_fract,
  data = nepaldem)

# Comparing OLS to RR fit for model_1999b
obj_cvdm_jm <- cvdm(percent_regvote1999 ~ landless_gap +
  below1pa_gap + sharecrop_gap + service_gap + fixmoney_gap +
  fixprod_gap + per_without_instcredit + totoalkilled_1000 +
  hdi_gap1 + ln_pop2001 + totalcontestants1999 + cast_eth_fract,
  data = nepaldem, method1 = "OLS", method2 = "RLM-MM")

obj_cvdm_jm
```

Index

- * **cox**
 - cvmf_object, 11
- * **cvmf**
 - cvmf_object, 11
- * **cvdm**
 - cvdm_object, 4
 - cvll_object, 8
 - cvlldiff_object, 8
- * **cvlldiff**
 - cvlldiff_object, 8
- * **cvll**
 - cvll_object, 8
- * **datasets**
 - govtform, 12
 - nepaldem, 14

binom.test, 11

coxph, 9–11

coxr, 9–11

cvdm, 2, 4, 5

cvdm_object, 4, 4, 6, 7

cvll, 5, 8

cvll_object, 8

cvlldiff, 6, 8

cvlldiff_object, 8

cvmf, 9, 11, 12

cvmf_object, 10, 11

govtform, 12

lm, 4, 8

modelLtest, 14

nepaldem, 14

rlm, 3, 5

rq, 3, 5

Surv, 9, 10